



HTA Austria
Austrian Institute for
Health Technology Assessment
GmbH

Covid-19



HSS/ Horizon Scanning
Living Document **V09 December 2020**



HTA Austria
Austrian Institute for
Health Technology Assessment
GmbH

Covid-19

HSS/ Horizon Scanning
Living Document **V09 December** 2020

Projektteam

Projektleitung: PD Dr. Claudia Wild

Projektbearbeitung

Updates: Mirjana Huic, MD, MSc, PhD

Projektbeteiligung

Kontroll- und Formatierarbeiten: Ozren Sehic, BA; Smiljana Blagojevic, Dipl.-Ing.

Korrespondenz: Claudia Wild, claudia.wild@aihta.at

Umschlagfoto: @Mike Fouque – stock.adobe.com

Dieser Bericht soll folgendermaßen zitiert werden/This report should be referenced as follows:

AIHTA Policy Brief Nr.: 002_V9 2020: Covid-19, HSS/ Horizon Scanning, Living Document December 2020.

Wien: HTA Austria – Austrian Institute for Health Technology Assessment GmbH.

Interessenskonflikt

Alle beteiligten AutorInnen erklären, dass keine Interessenskonflikte im Sinne der Uniform Requirements of Manuscripts Statement of Medical Journal Editors (www.icmje.org) bestehen.

IMPRESSUM

Medieninhaber und Herausgeber:

HTA Austria - Austrian Institute for Health Technology Assessment GmbH
Garnisongasse 7/Top20 | 1090 Wien – Österreich
www.aihta.ac.at

Für den Inhalt verantwortlich:

Priv.-Doz. Dr. phil. Claudia Wild, Geschäftsführung

Die **AIHTA Policy Briefs** erscheinen unregelmäßig und dienen der Veröffentlichung der Forschungsergebnisse des Austrian Institute for Health Technology Assessment.

Die **AIHTA Policy Briefs** erscheinen in geringer Auflage im Druck und werden über den Dokumentenserver „<https://eprints.aihta.at/view/types/policy=5Fbrief.htm>“ der Öffentlichkeit zur Verfügung gestellt.

AIHTA Policy Brief Nr.: 002

ISSN 2710-3234

ISSN online 2710-3242

© 2020 AIHTA – Alle Rechte vorbehalten

Content

Content.....	3
1 Background: policy question and methods.....	7
1.1 Policy Question.....	7
1.2 Methodology.....	7
1.3 Selection of Products for “Vignettes”.....	10
2 Results: Vaccines.....	11
2.1 Moderna Therapeutics—US National Institute of Allergy.....	19
2.2 CanSino Biological Inc. and Beijing Institute of Biotechnology.....	20
2.3 University of Oxford/ Astra Zeneca.....	21
2.4 BioNTech/Fosun Pharma/Pfizer.....	24
2.5 Sinovac Biotech Ltd.....	25
2.6 China National Pharmaceutical Group Corporation (SINOPHARM).....	26
2.7 Gamaleya Research Institute.....	27
2.8 Janssen Pharmaceutical.....	28
2.9 Novavax.....	29
3 Results: Therapeutics.....	31
3.1 Remdesivir (Veklury®).....	33
3.2 Lopinavir + Ritonavir (Kaletra®).....	42
3.3 Favipiravir (Avigan®).....	42
3.4 Darunavir.....	51
3.5 Chloroquine (Resochin®) and.....	53
3.6 Hydroxychloroquine (Plaquenil®).....	53
3.7 Camostat Mesilate (Foipan®).....	53
3.8 APN01/ Recombinant Human Angiotensin-converting Enzyme 2 (rhACE2).....	54
3.9 Tocilizumab (Roactemra®).....	54
3.10 Sarilumab (Kevzara®).....	58
3.11 Interferon beta 1a (SNG001) (Rebif®, Avonex®) and Interferon beta 1b (Betaferon®, Extavia®).....	58
3.12 Convalescent plasma transfusion.....	64
3.13 Plasma derived medicinal products.....	69
3.13.1 REGN-COV2 - combination of two monoclonal antibodies (REGN10933 and REGN10987).....	69
3.13.2 LY-CoV555 - neutralizing IgG1 monoclonal antibody (bamlanivimab) and LY-CoV016 - recombinant fully human monoclonal neutralizing antibody (etesevimab).....	71
3.13.3 AZD7442 - combination of two monoclonal antibodies (AZD8895 + AZD1061).....	73
3.14 Combination therapy – triple combination of interferon beta-1b, lopinavir–ritonavir and ribavirin vs. lopinavir–ritonavir or other triple combination of interferons.....	73
3.15 Solnatide.....	79
3.16 Umifenovir (Arbidol®).....	80
3.17 Dexamethasone and other corticosteroids.....	84
3.18 Anakinra (Kineret®).....	90
3.19 Colchicine.....	90
3.20 Nafamostat (Futhan©).....	93
3.21 Gimsilumab.....	93
3.22 Canakinumab.....	93
3.23 Lenzilumab.....	94
3.24 Vitamin D.....	95
3.25 Baricitinib.....	98
3.26 Molnupiravir.....	99
References.....	100

Figures

Figure 1.2-1: A living mapping of ongoing randomized trials, living systematic reviews with pairwise meta-analyses and network meta-analyses	9
Figure 1.2-2: Global Coronavirus COVID-19 Clinical Trial Tracker - a real-time dashboard of clinical trials for COVID-19	10

Tables

Table 1.2-1: International Sources	8
Table 2-1: Vaccines in the R&D pipeline (Phase 1 - Phase 3 clinical trials, not preclinical stages),	14
Table 3-1: COVID-19 medicines that have received EMA advice.....	31
Table 3-2: Most advanced therapeutics in the R&D pipeline	32
Table 3.1-1: Summary of findings table on Remdesivir vs Standard care /Placebo (4 RCTs: Wang, Beigel, Spinner, SOLIDARITY-Remdesivir	37
Table 3.1-2: Summary of findings table on Remdesivir 5 days vs Remdesivir 10 days (2 RCTs: Goldman, Spinner)	41
Table 3.3-1: Summary of findings table on favipiravir compared to umifenovir (1 RCT: Chen)	45
Table 3.3-2: Summary of findings table on favipiravir compared to baloxavir marboxil (1 RCT: Lou 2020).....	47
Table 3.3-3: Summary of findings table on favipiravir compared to lopinavir + ritonavir or darunavir/cobicistat + umifenovir + interferon-a (1 RCT: Lou 2020).....	48
Table 3.3 4: Summary of findings table on favipiravir compared to standard care (6 RCTs: Lou 2020, Ivashchenko 2020, Dabbous 2020, Balykova 2020, Ruzhentsova 2020, Udwardia 2020	49
Table 3.4-1: Summary of findings table on darunavir/cobicistat compared to standard care (1 RCT: Chen J)	52
Table 3.9-1: Summary of findings table on tocilizumab compared standard care/placebo (6 RCTs: Rosas, Wang, Hermine, Salvarani, Stone, Salama).....	57
Table 3.11-1: Summary of findings table on Interferon β -1a compared to Standard Care for Moderate/Severe/Critical COVID-19 (3 RCTs: Davoudi-Monfared, Rahmani, SOLIDARITY-INF).....	62
Table 3.12-1: Summary of findings table on Convalescent plasma compared to Standard Care for Mild/Moderate/Severe/Critical COVID-19 (4 RCTs: Li, Gharbharan, Avendano-Sola, Agarwal)	66
Table 3.14-1: Summary of findings table on triple combination of interferon beta-1b, lopinavir-ritonavir and ribavirin (1 RCT: Hung).....	74
Table 3.14-1 continued: Summary of findings tables on Novaferon , Lopinavir/Ritonavir and Novaferon + Lopinavir/Ritonavir (1 RCT: Zheng 2020)	76
Table 3.16-1. Summary of findings table, on umifenovir vs standard care (2 RCTs:Yueping, Yethindra)	81
Table 3.17-1: Summary of findings table, on dexamethasone and other corticosteroids (7 RCTs: Horbey, Tomazini, Dequin, REMAP-CAP Investigators, Jeronimo, Corral, Edalatifard).....	87
Table 3.19-1: Summary of findings table on colchicine compared to standard care (3 RCT: Deftereos, Lopes, Salehzadeh).....	91
Table 3.24-1: Summary of findings table on vitamin D compared to standard care (3 RCT:Entrenas Castillo, Rastogi, Murai).....	96

History of Changes	V09 December
08-11/12/2020	Addition chapter on Baricitinib (chapter 3.25) and Molnupiravir (chapter 3.26)
09/12/2020	Methodology (1.2) – no changes
10/12/2020	Update Vaccine (chapter 2)
17/12/2020	Update Remdesivir (chapter 3.1)
17/12/2020	Update Favipiravir (chapter 3.3)
12/12/2020	Darunavir (chapter 3.4) – no changes
13/12/2020	Camostat Mesilate (chapter 3.7) – no changes
13/12/2020	APN01/rhACE2 (chapter 3.8) – no changes
13/12/2020	Tocilizumab (chapter 3.9) – no changes
13/12/2020	Sarilumab (chapter 3.10) – no changes
14/12/2020	Update Interferon beta (chapter 3.11)
17/12/2020	Update Concoalescent plasma (chapter 3.12)
16/12/2020	Update Plasma derived medicinal products (chapter 3.13) – REGN-COV2; LY-CoV555 and LY-CoV016; AZD7422
15/12/2020	Combination therapy (chapter 3.14) – no changes
15/12/2020	Solnatide (chapter 3.15) – no changes
16/12/2020	Update Umifenovir (chapter 3.16)
17/12/2020	Update Dexamethasone and other corticosteroids (chapter 3.17)
17/12/2020	Anakinra (chapter 3.18) - no changes
17/12/2020	Update Colchicine (chapter 3.19)
16/12/2020	Nafamostat (chapter 3.20) – no changes
16/12/2020	Gimsilumab (chapter 3.21) – no changes
16/12/2020	Canakinumab (chapter 3.22) – no changes
16/12/2020	Lenzilumab (chapter 3.23) – no changes
17/12/2020	Update Vitamin D (chapter 3.24)

1 Background: policy question and methods

1.1 Policy Question

On March 30th 2020, a request was raised by the Austrian Ministry of Health (BMASGK), the Health Funds of the Regions and the Federation of Social Insurances to set up a Horizon Scanning system (HSS) for medicines and vaccines. The establishment of a HSS/ Horizon Scanning System for Covid-19 interventions has the intentions of

- a. informing health policy makers at an early stage which interventions (vaccinations and drugs) are currently undergoing clinical trials and
- b. monitoring them over the next few months in order to support evidence-based purchasing, if necessary.

1.2 Methodology

To respond to this request,

1. As a first step an inventory, based on international sources, is built.
2. As a second step, selective searches by means of searches in study registries are carried out for information on clinical studies in humans and the state of research.
3. This information forms the basis for "vignettes" (short descriptions) for those products that are already in an "advanced" stage.
4. Subsequently, the products are monitored with regard to the status of the clinical studies up to approval and finally evaluated for their benefit and harm.

All work steps are conducted in close international (European) cooperation.

- Version 1 (V1, April 2020): inventory + vignettes for most advanced
- Version 2+: monthly monitoring and updates

Ongoing trials are reported in V1, April 2020 - V3, June 2020 of this Document and in the living documents - EUnetHTA (Covid-19 Rolling Collaborative Reviews: <https://eunetha.eu/rcr01-rcrxx/>).

From V4 July, 2020 of this HSS/ Horizon Scanning Document, only completed, terminated, withdrawn and suspended interventional clinical trials from ClinicalTrials.gov and EUdraCT registers are reported. From Version 8 November, 2020 only terminated, withdrawn and suspended interventional clinical trials are reported.

From V5, August 2020 of this HSS/ Horizon Scanning Document only the best available evidence will be presented in.

**März 2020:
Österr. Politik empfiehlt
Aufbau von HSS
zu Covid-19**

**Information zu
* Status F&E
* Evidenz-basierter
Einkauf**

mehrstufige Methodik

**Bestandsaufnahme
selektive Suche
Vignetten
Monitoring**

**internationale/
europ. Zusammenarbeit**

**V1-V3: auch laufende
Studien - Verweis auf
EUnetHTA
V4: nur abgeschlossene
(oder beendete)
Interventionsstudien aus
2 Studienregistern
ab V5: nur mehr best
verfügbare Evidenz**

Table 1.2-1: International Sources

Primary sources	Link
WHO Drugs: Vaccines:	https://www.who.int/teams/blueprint/covid-19 https://www.who.int/blueprint/priority-diseases/key-action/Table_of_therapeutics_Appendix_17022020.pdf?ua=1 https://www.who.int/who-documents-detail/covid-19-candidate-treatments https://www.who.int/who-documents-detail/draft-landscape-of-covid-19-candidate-vaccines
Danish Medicine Agency Drugs: Vaccines:	https://laegemiddelstyrelsen.dk/da/nyheder/temaer/ny-coronavirus-covid-19/~~/media/5B83D25935DF43A38FF823E24604AC36.ashx https://laegemiddelstyrelsen.dk/da/nyheder/temaer/ny-coronavirus-covid-19/~~/media/3A4B7F16D0924DD8BD157BBE17BFED49.ashx
Pang et al. 2020 [1] Drugs: Vaccines:	https://www.mdpi.com/2077-0383/9/3/623 Table 5+6, Table 3+4
SPS HS-report (UK)	unpublished
Secondary sources	
VfA/ Verband Forschender Arzneimittelhersteller Drugs: Vaccines:	https://www.vfa.de/de/arsneimittel-forschung/woran-wir-forschen/therapeutische-medikamente-gegen-die-coronavirusinfektion-covid-19 https://www.vfa.de/de/arsneimittel-forschung/woran-wir-forschen/impfstoffe-zum-schutz-vor-coronavirus-2019-ncov
EMA/ European Medicines Agency Medicines:	https://www.ema.europa.eu/ https://www.ema.europa.eu/en/medicines/medicines-under-evaluation
FDA/US Food and Drug Administration	https://www.fda.gov/emergency-preparedness-and-response/counterterrorism-and-emerging-threats/coronavirus-disease-2019-covid-19
Trial Registries	
US National Library of Medicine European Union Drug Regulating Authorities Clinical Trials Database WHO International Clinical Trials Registry Platform TrialsTracker	https://clinicaltrials.gov/ https://eudract.ema.europa.eu/ https://www.who.int/ictrp/en/ http://Covid-19.trialstracker.net/
Up-to-date information on clinical trials and literature searching resources relating to COVID-19	
Cochrane COVID-19 Study Register 21/04.20	https://covid-19.cochrane.org/
Living mapping of research and a living systematic review	https://covid-nma.com/ https://covid-nma.com/dataviz/
Dynamic meta-analysis of evidences about drug efficacy and safety for COVID19 - meta/Evidence – COVID-19	http://metaevidence.org/COVID19.aspx
CORDITE (CORona Drug InTEractions database)	https://cordite.mathematik.uni-marburg.de/#/
Living listing of interventional clinical trials in Covid-19/2019-nCoV produced by the Anticancer Fund	http://www.redo-project.org/covid19db/ ; http://www.redo-project.org/covid19_db-summaries/
Global Coronavirus COVID-19 Clinical Trial Tracker	https://www.covid-trials.org/
LitCovid	https://www.ncbi.nlm.nih.gov/research/coronavirus/
UK NIHR Innovation Observatory NIHR COVID-19 Studies COVID-19 Therapeutics Dashboard COVID-19: a living systematic map of the evidence	https://www.nihr.ac.uk/covid-studies/ http://www.io.nihr.ac.uk/report/covid-19-therapeutics/ http://epi.ioe.ac.uk/cms/Default.aspx?tabid=3765
WHO COVID-19 Database new search interface	https://www.who.int/emergencies/diseases/novel-coronavirus-2019/global-research-on-novel-coronavirus-2019-ncov
COVID-evidence Database	https://covid-evidence.org/database
Medical Library Association – COVID-19 Literature search strategies	https://www.mlanet.org/page/covid-19-literature-searching

Centre of Evidence Based Dermatology (CEBD) - Coronavirus Dermatology Online Resource	https://www.nottingham.ac.uk/research/groups/cebd/resources/Coronavirus-resource/Coronavirushom
Ovid Expert Searches for COVID-19	http://tools.ovid.com/coronavirus/
EBSCO Covid-19 Portal Literature searching section of portal Information portal	https://covid-19.ebscomedical.com/research https://covid-19.ebscomedical.com/
NIH COVID-19 Treatment Guidelines, 2020.	https://covid19treatmentguidelines.nih.gov/introduction/
Tertiary sources	
NIPHNO	https://www.fhi.no/en/qk/systematic-reviews-hta/map/
INAHTA	http://www.inahta.org/covid-19-inahta-response/
EUnethTA Covid-19 Rolling Collaborative Reviews (RCR)	https://eunetha.eu/rcr01-rcrx/

Several organisations and international teams of researchers are providing up-to-date information through living listing of interventional clinical trials in Covid-19/2019-nCoV and literature resources (Table 1.2-1) [2-4] [2]. A short description of two of such databases is presented below.

“lebende” Dokumente mit up-to-date Informationen

Boutron et al., 2020 [3] are performing a living mapping of ongoing randomized trials, followed by living systematic reviews with pairwise meta-analyses and when possible, network meta-analyses focusing on two main questions: the effectiveness of preventive interventions for COVID-19 and the effectiveness of treatment interventions for COVID-19 (Figure 1.2-1).

Kartierung von laufenden RCTs

COVID-19 NMA


a living mapping of ongoing research.

▼ As of April 24, 2020...

The Covid-19 - Living NMA initiative collected a number of 506 studies of treatments from the ICTRP. 278 of these trials are recruiting patients. Most of the studies are being conducted in Asia (264 trials) with the majority from China (151 trials). Other countries in Europe (160 trials) and North America (92 trials) are rapidly setting up new trials with the majority being conducted in multiple centers (194 trials).

Search:
 Ex: Intervention, treatment, open, Assistance, Pharmazie, EUCTR2020...

▼ Map



▼ HELP

- Make your browser window as wide as possible for a 2-column display.
- Click on the map or any of the graphs to create filters on the data.
- All the filters are applied jointly, refining your selection.
- To select a Registration date, click and drag to create a range.
- At any moment you can click Reset All below to remove the filters.
- Click on the black arrows to open or close any section.
- For any questions or remarks, please contact us.

All trials selected (506) | Reset All


▼ Recruitment status

- Recruiting (278)
- Not recruiting (229)
- Completed (9)

▼ Registration date



▼ Study aim



▼ Disease severity



Figure 1.2-1: A living mapping of ongoing randomized trials, living systematic reviews with pairwise meta-analyses and network meta-analyses

Clinical Trial Tracker real-time dashboard

Thorlund et al., 2020 [4] developed a COVID-19 clinical trials registry to collate all trials related to COVID-19: Global Coronavirus COVID-19 Clinical Trial Tracker. Data is pulled from the International Clinical Trials Registry Platform, including those from the Chinese Clinical Trial Registry, ClinicalTrials.gov, Clinical Research Information Service - Republic of Korea, EU Clinical Trials Register, ISRCTN, Iranian Registry of Clinical Trials, Japan Primary Registries Network, and German Clinical Trials Register (Figure 1.2-2). They also use content aggregator services, such as LitCovid, to ensure that their data acquisition strategy is complete [5].



Figure 1.2-2: Global Coronavirus COVID-19 Clinical Trial Tracker - a real-time dashboard of clinical trials for COVID-19

1.3 Selection of Products for “Vignettes”

The following products have been selected for further investigation (searches in registry databases and description as “vignettes”) for the following reasons:

- most advanced in clinical research in humans
- most often discussed in clinical journals as potential candidates

The full inventory (list) can be found in Part 2 - Appendix A-1: vaccines, A-2, therapeutics, A3-EudraCT registry studies.

Vignetten zu Produkte, in "fortgeschrittenen" Stadien oder

häufig diskutiert/publiziert

2 Results: Vaccines

As of December 12 2020, thirteen COVID-19 candidate vaccines are investigated in phase 3 RCTs:

13 Impfstoffe in Phase 3

1. **Moderna Therapeutics/NIAID** (RNA LNP-encapsulated mRNA vaccine encoding S protein);
2. **CanSino Biological** (Non-Replicating Viral Vector adenovirus Type 5 Vector vaccine that expresses S protein);
3. **University of Oxford/AstraZeneca** (Non-Replicating Viral Vector ChAdOx1 (AZD1222) vaccine);
4. **BioNTech/Fosun Pharma/Pfizer** (RNA 3 LNP-mRNAs vaccine);
5. **Sinovac Biotech** (inactivated vaccine);
6. **Sinopharm/Wuhan Institute of Biological Products** (inactivated vaccine);
7. **Sinopharm/Beijing Institute of Biological Products** (inactivated vaccine)
8. **Gamaleya Research Institute** (Non-Replicating Viral Vector Adeno-based - rAd5, rAd26) vaccine; and
9. **Janssen Pharmaceuticals** (Non-Replicating Viral Vector Ad26COVS1 vaccine); and
10. **Novavax** (Protein Subunit, VLP-recombinant protein nanoparticle vaccine + Matrix M) vaccine;
11. **Bharat Biotech** (inactivated vaccine);
12. **Anhui Zhifei Longcom Biopharmaceutical/Institute of Microbiology, Chinese Academy of Sciences** (Protein Subunit, Adjuvanted recombinant protein RBF.Dimer, expressed in CHO cells);
13. **Medicago Inc.** (VLP, Plant-derived VLP adjuvanted AS03).

For these 13 coronavirus vaccines are investigated in the phase 3 RCTs, the following articles were published with results related to early phases vaccine trials (phase 1, 1/2 or phase 2) or phase 2/3 trials:

1. Two on **Moderna Therapeutics/NIAID** vaccine: a preliminary report with the results from the phase 1 study (NCT04283461) [6] and
2. the results from the expanded phase 1 study (NCT04283461) in older adults [7];
3. Two on **CanSino Biological** vaccine: the results from the phase 1, dose-escalation, open-label, non-randomised, first-in-human trial for adenovirus type-5 vectored COVID-19 vaccine (ChiCTR2000030906/NCT04313127) [8], as well as
4. phase 2, randomised controlled trials (ChiCTR2000031781/NCT04398147) [9];
5. One on **Novavax** vaccine: the results from the phase 1/2 RCT (NCT04368988) [10];
6. Three on **Oxford/Astra Zeneca** vaccine: a preliminary report with the results from phase 1/2 single-blind, RCT (ISRCTN 15281137/NCT04324606/EudraCT 2020-001072-15) [11],

16 Publikationen zu

Phase 1,
1/2 oder
Phase 2
Phase 2/3

Impfstudien

Results: Vaccines

7. pooled interim analysis phase 2/3 trials (ISRCTN89951424, NCT04324606, NCT04400838, and NCT04444674) [12] and
8. phase 2 component of phase 2/3 trial COV002 (ISRCTN90906759, NCT04400838) [13];
9. One with results on **Gamaleya vaccine**, from two open, non-randomised phase 1/2 studies at two hospitals in Russia (NCT04436471 and NCT04437875) [14];
10. Four on **BioNTech/Fosun Pharma/Pfizer** vaccine: Three with results from two phase 1/2 trials on **BNT162b1** vaccine, one in US (NCT04368728/EudraCT 2020-001038-36) [15],
11. and one in Germany (NCT04380701, EudraCT 2020-001038-36) [16] as well as
12. additional safety and immunogenicity results from the US phase 1 trial (NCT04368728/EudraCT 2020-001038-36) [52, 53] and
13. one pivotal RCT efficacy trial on BNT162b2 (NCT04368728) [17].
14. Two related to **Sinopharm vaccine**: results from two double-blind RCTs, phase 1 and phase 2 (ChiCTR2000031809) [18, 19] on Sinopharm/**Wuhan Institute of Biological Products** vaccine and
15. results from phase 1/2 clinical trials (ChiCTR2000032459) [20] on Sinopharm/**Beijing Institute of Biological Products**, BBIBP-CorV vaccine; and
16. One to **Sinovac** vaccine: results from RCT, phase 1/2 clinical trial (NCT04352608) [21].

Approval status:

On 09/07/2020, Medicines Regulatory Authorities published the report related to phase 3 COVID-19 vaccine trials [23]. They stressed the need for large phase 3 clinical trials that enroll many thousands of people, including those with underlying medical conditions, to generate relevant data for the key target populations. Broad agreement was achieved that clinical studies should be designed with stringent success criteria that would allow a convincing demonstration of the efficacy of COVID-19 vaccines.

On November 11, 2020 EMA publishes safety monitoring plan and guidance on risk management planning for COVID-19 vaccines, <https://www.ema.europa.eu/en/news/ema-publishes-safety-monitoring-plan-guidance-risk-management-planning-covid-19-vaccines>.

On October 01, 2020 EMA announced that **EMA's** human medicines committee (CHMP) has started the first '**rolling review**' of University of **Oxford/AstraZeneca** vaccine [16].

On October 06, 2020 EMA's human medicines committee (CHMP) has started a 'rolling review' of data on a BNT162b2 vaccine, which is being developed by **BioNTech** in collaboration with Pfizer [24]. On 01/12/2020 EMA started with evaluation of conditional marketing authorisation application for this vaccine [25]

On 11 December 2020, **FDA** issued the first emergency use authorisation (**EUA**) for a this vaccine for the prevention of COVID-19 caused by SARS-CoV-2 in individuals 16 years and older in US, <https://www.fda.gov/news-events/press-announcements/fda-takes-key-action-fight-against-covid-19-issuing-emergency-use-authorization-first-covid-19>.

Positionspapier der Internationalen Regulatoren

derzeit 4 Impfstoffe in „rolling review“ bei EMA – Zulassung:

**Oxford/AstraZeneca
BioNTech/ Pfizer
Moderna
Janssen**

FDA Notzulassung am 11. Dez 2020

On November 16, 2020 **EMA** announced that EMA's human medicines committee (CHMP) has started a 'rolling review' of data on a mRNA-1273 COVID-19 vaccine, developed by **Moderna** Biotech Spain, S.L. (a subsidiary of Moderna, Inc.), and on 01/12/2020 started with evaluation of conditional marketing authorisation application [26]. On December 17, 2020 FDA announced the meeting to discuss EUA of the Moderna, Inc., COVID-19 vaccine for the prevention of COVID-19 in individuals 18 years and older in US, <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-december-17-2020-meeting-announcement>.

On December 01, 2020 **EMA** announced that EMA's human medicines committee (CHMP) has started a '**rolling review**' of **Janssen-Cilag** International N.V COVID-19 Ad26.COV2.S vaccine [27].

**EMA Zulassungen im
Dezember angekündigt**

Results: Vaccines

Table 2-1: Vaccines in the R&D pipeline (Phase 1 - Phase 3 clinical trials, not preclinical stages), December 10, 2020

Source: DRAFT landscape of COVID-19 candidate vaccines – 10 December-2020 – 52 candidate vaccines in clinical evaluation, <https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccines>

COVID-19 Vaccine developer/manufacturer	Vaccine platform	Type of candidate vaccine	Number of doses	Timing of doses	Route of Administration	Clinical Stage			
						Phase 1	Phase 1/2	Phase 2	Phase 3
Sinovac	Inactivated	Inactivated	2	0,14 days	IM		NCT04383574 NCT04352608 Study Report NCT04551547		NCT04456595 669/UN6.KEP/EC/2020 NCT04582344 NCT04617483
Wuhan Institute of Biological Products/Sinopharm	Inactivated	Inactivated	2	0,21 days	IM		ChiCTR2000031809 Interim Report		ChiCTR2000034780 ChiCTR2000039000 NCT04612972
Beijing Institute of Biological Products/Sinopharm	Inactivated	Inactivated	2	0,21 days	IM		ChiCTR2000032459 Study Report		ChiCTR2000034780 NCT04560881
Bharat Biotech	Inactivated	Whole-Virion Inactivated	2	0, 28 days	IM		CTRI/2020/07/026300 CTRI/2020/09/027674		CTRI/2020/11/028976 NCT04641481
University of Oxford/AstraZeneca	Non-Replicating Viral Vector	ChAdOx1-S	2	0,28 days	IM		PACTR202006922165132 2020-001072-15 NCT04568031 Interim Report	2020-001228-32 Study Report	ISRCTN89951424 NCT04516746 NCT04540393 CTRI/2020/08/027170
CanSino Biological Inc./Beijing Institute of Biotechnology	Non-Replicating Viral Vector	Adenovirus Type 5 Vector	1		IM	ChiCTR2000030906 NCT04568811 Study Report		ChiCTR2000031781 NCT04566770 Study Report	NCT04526990 NCT04540419

Results: Vaccines

Gamaleya Research Institute	Non-Replicating Viral Vector	Adeno-based (rAd26-S+rAd5-S)	2	0, 21 days	IM		NCT04436471 NCT04437875 Study Report	NCT04587219 NCT04640233	NCT04530396 NCT04564716 NCT04642339
Janssen Pharmaceutical Companies	Non-Replicating Viral Vector	Adenovirus Type 26 vector	1 2	0 0, 56 days	IM		NCT04436276 NCT04509947	NCT04535453	NCT04505722 ISRCTN14722499
Novavax	Protein Subunit	Full length recombinant SARS CoV-2 glycoprotein nanoparticle vaccine adjuvanted with Matrix M	2	0, 21 days	IM		NCT04368988 Study Report	NCT04533399 (phase 2b)	2020-004123-16 NCT04611802
Anhui Zhifei Longcom Biopharmaceutical/Institute of Microbiology, Chinese Academy of Sciences	Protein Subunit	Adjuvanted recombinant protein (RBD-Dimer) expressed in CHO cells	3	0, 28, 56 days	IM	NCT04445194 NCT04636333	NCT04550351	NCT04466085	ChiCTR2000040153
Moderna/NIAID	RNA	LNP-encapsulated mRNA	2	0, 28 days	IM	NCT04283461 Interim Report Final Report		NCT04405076	NCT04470427
BioNTech/Fosun Pharma/Pfizer	RNA	3 LNP-mRNAs	2	0, 21 days	IM	NCT04368728 Study Report	2020-001038-36 ChiCTR2000034825 NCT04537949 NCT04588480 Study Report1 Study Report2		NCT04368728
Medicago Inc.	VLP	Plant-derived VLP adjuvanted with AS03.	2	0, 21 days	IM	NCT04450004			NCT04636697
Inovio Pharmaceuticals/ International Vaccine Institute	DNA	DNA plasmid vaccine with electroporation	2	0, 28 days	ID		NCT04447781 NCT04336410	NCT04642638 ChiCTR2000040146	
Beijing Wantai Biological Pharmacy/ Xiamen University	Replicating Viral Vector	Intranasal flu-based-RBD	1		IN	ChiCTR2000037782		ChiCTR2000039715	
West China Hospital, Sichuan University	Protein Subunit	RBD (baculovirus production expressed in Sf9 cells)	2 or 3	0, 28 days and 0, 14, 28 days	IM	ChiCTR2000037518		ChiCTR2000039994	
Curevac	RNA	mRNA	2	0, 28 days	IM	NCT04449276		NCT04515147	

Results: Vaccines

Institute of Medical Biology, Chinese Academy of Medical Sciences	Inactivated	Inactivated	2	0, 28 days	IM	NCT04412538	NCT04470609		
Research Institute for Biological Safety Problems, Rep of Kazakhstan	Inactivated	Inactivated	2	0, 21 days	IM		NCT04530357		
Shenzhen Kangtai Biological Products Co., Ltd.	Inactivated	Inactivated	2		IM	ChiCTR2000038804	ChiCTR2000039462		
Osaka University/ AnGes/ Takara Bio	DNA	DNA plasmid vaccine + Adjuvant	2	0, 14 days	IM		NCT04463472 NCT04527081		
Cadila Healthcare Limited	DNA	DNA plasmid vaccine	3	0, 28, 56 days	ID		CTRI/2020/07/026352		
Genexine Consortium	DNA	DNA Vaccine (GX-19)	2	0, 28 days	IM		NCT04445389		
Kentucky Bioprocessing, Inc	Protein Subunit	RBD-based	2	0, 21 days	IM		NCT04473690		
Sanofi Pasteur/GSK	Protein Subunit	S protein (baculovirus production)	2	0, 21 days	IM		NCT04537208		
Biological E Ltd	Protein Subunit	Adjuvanted protein subunit (RBD)	2	0, 28 days	IM		CTRI/2020/11/029032		
Israel Institute for Biological Research	Replicating Viral Vector	VSV-S	1		IM		NCT04608305		
Arcturus/Duke-NUS	RNA	mRNA			IM		NCT04480957		
SpyBiotech/Serum Institute of India	VLP	RBD-HBsAg VLPs	2	0, 28 days	IM		ACTRN12620000817943		
Symvivo	DNA	bacTRL-Spike	1		Oral	NCT04334980			
Providence Health & Services	DNA	electroporated S protein plasmid DNA vaccine with or without the combination of electroporated IL-12p70 plasmid	2	0, 28 days	ID	NCT04627675			
Codagenix/Serum Institute of India	Live Attenuated Virus	Codon deoptimized live attenuated vaccines	1 or 2	0 or 0,28 days	IN	NCT04619628			
ImmunityBio, Inc. & NantKwest Inc.	Non-Replicating Viral Vector	hAd5 S+N 2nd Generation Human Adenovirus Type 5 Vector (hAd5) Spike (S) + Nucleocapsid (N)	2	0, 21 days	SC	NCT04591717			

Results: Vaccines

ReiThera/LEUKOCARE/Univercells	Non-Replicating Viral Vector	Replication defective Simian Adenovirus (GRAd) encoding S	1		IM	NCT04528641			
CanSino Biological Inc/Institute of Biotechnology, Academy of Military Medical Sciences, PLA of China	Non-Replicating Viral Vector	Ad5-nCoV	2	0, 28 days	IM/mucosal	NCT04552366			
Vaxart	Non-Replicating Viral Vector	Ad5 adjuvanted Oral Vaccine platform	2	0, 28 days	Oral	NCT04563702			
Ludwig-Maximilians - University of Munich	Non-Replicating Viral Vector	MVA-SARS-2-S	2	0, 28 days	IM	NCT04569383			
City of Hope, USA	Replicating Viral Vector	SARS-CoV-2 S and NP genes inserted into a sMVA vector	2	0, 28 days	IM	NCT04639466			
Clover Biopharmaceuticals Inc./GSK/Dynavax	Protein Subunit	Native like Trimeric subunit Spike Protein vaccine	2	0, 21 days	IM	NCT04405908			
Vaxine Pty Ltd/Medytox	Protein Subunit	Recombinant spike protein with Advax™ adjuvant	1		IM	NCT04453852			
University of Queensland/CSL/Seqirus	Protein Subunit	Molecular clamp stabilized Spike protein with MF59 adjuvant	2	0, 28 days	IM	ACTRN12620000674932p ISRCTN51232965			
Medigen Vaccine Biologics Corporation/NIAID/Dynavax	Protein Subunit	S-2P protein + CpG 1018	2	0, 28 days	IM	NCT04487210			
Instituto Finlay de Vacunas, Cuba	Protein Subunit	rRBD produced in CHO-cell chemically conjugate to tetanus toxoid	2	0, 28 days	IM	IFV/COR/06			
Instituto Finlay de Vacunas, Cuba	Protein Subunit	RBD + Adjuvant	2	0, 28 days	IM	IFV/COR/04 IFV/COR/05			
FBRI SRC VB VECTOR, Rospotrebnadzor, Koltsovo	Protein Subunit	Peptide	2	0, 21 days	IM	NCT04527575			
University Hospital Tuebingen	Protein Subunit	SARS-CoV-2 HLA-DR peptides	1		SC	NCT04546841			

Results: Vaccines

COVAXX / United Biomedical Inc. Asia	Protein Subunit	Multitope peptide-based S1-RBD-protein vaccine	2	0, 28 days	IM	NCT04545749			
Chinese Academy of Military Sciences	Protein Subunit	Subunit expressed in CHO cells	2 or 3	0, 14 days or 0,14, 28 days	IM				
Merck Sharp & Dohme/IAVI	Replicating Viral Vector	Replication-competent VSV delivering the SARS-CoV-2 Spike	1		IM	NCT04569786			
Institute Pasteur/Themis/Univ. of Pittsburg CVR/Merck Sharp & Dohme	Replicating Viral Vector	Measles-vector based	1 or 2	0, 28 days	IM	NCT04497298			
Imperial College London	RNA	LNP-nCoVsaRNA	2		IM	SRCTN17072692			
People's Liberation Army (PLA) Academy of Military Sciences/Walvax Biotech.	RNA	mRNA	2	0, 14 or 0, 28 days	IM	ChiCTR2000034112 ChiCTR2000039212			

2.1 Moderna Therapeutics—US National Institute of Allergy

About the vaccine

The **mRNA-1273** vaccine candidate developed by ModernaTX, Inc. in collaboration with NIAID and sponsored by NIAID/CEPI is an LNP-encapsulated mRNA-based vaccine (mRNA-1273) intended for prevention through full-length, perfusion stabilized spike (S) protein of SARS-CoV-2 that is the key into the human cell [28].

**mRNA-1273
collab mit NIAID/CEPI**

Estimated timeline for approval

Phase 1 trial with 45 healthy participants (NCT04283461) is ongoing. Participants are split to 3 groups where they receive two injections of low (25 mcg), medium (100 mcg) or high doses (250 mcg) of mRNA-1273 and are monitored for any AEs and immune response [29]. The Phase I safety study should be completed by June 2021.

**Phase 1:
45 gesunde Erwachsene
Juni 2021**

A **phase 2a**, randomized, observer-blind, placebo controlled, dose-confirmation study to evaluate the safety, reactogenicity, and immunogenicity of mRNA-1273 vaccine in adults aged 18 years and older (NCT04405076) is underway. This Phase 2 study should be completed by August 2021.

**Phase 2a:
bis August 2021**

The randomized, **phase 3**, 1:1 placebo-controlled trial is currently ongoing (NCT04470427). It is expected to include approximately 30,000 participants enrolled in the U.S.

**Phase 3 Studienprotokoll
RCT mt ca 30.000
Teilnehmer*innen**

Results of publications

A preliminary report with the results from the above mentioned **phase 1** study was published [6]. After the first vaccination, antibody responses were higher with higher dose (day 29 enzyme-linked immunosorbent assay anti-S-2P antibody geometric mean titer [GMT], 40,227 in the 25- μ g group, 109,209 in the 100- μ g group, and 213,526 in the 250- μ g group). After the second vaccination, the titers increased (day 57 GMT, 299,751, 782,719, and 1,192,154, respectively). Systemic adverse events were more common after the second vaccination, particularly with the highest dose, and three participants (21%) in the 250- μ g dose group reported one or more severe adverse events.

**vorläufige Publikation der
Phase 1 Studie**

**höhere Dosis,
höhere Titer**

Anderson et al. 2020 [7] published results from the above mentioned phase 1 trial in healthy adults, which was expanded to include 40 older adults, who were stratified according to age (56 to 70 years or ≥ 71 years). All the participants were assigned sequentially to receive two doses of either 25 μ g or 100 μ g of vaccine administered 28 days apart. Solicited adverse events were predominantly mild or moderate in severity. Binding-antibody responses increased rapidly after the first immunization. The 100- μ g dose induced higher binding- and neutralizing-antibody titers than the 25- μ g dose, which supports the use of the 100- μ g dose in a phase 3 vaccine trial.

**Okt. 2020:
Publikation der
Phase 1 Studie**

**unterschiedliche
Dosierung in
verschiedenen
Altersgruppen**

On November 30, 2020, Moderna, Inc. **announced results** from primary efficacy analysis of the **phase 3** COVE study enrolled more than 30,000 participants ages 18 and older in the U.S. Primary analysis was based on 196 cases, of which 185 cases of COVID-19 were observed in the placebo group versus 11 cases observed in the mRNA-1273 group, a point estimate of vaccine efficacy of 94.1%. A secondary endpoint analyzed severe cases of COVID-19 and included 30 severe cases in this analysis. All 30 cases occurred in the placebo group and none in the mRNA-1273 vaccinated group; one COVID-19-related death occurred in the placebo group.

**Nov. 2020:
COVE, 30.000
Teilnehmer*innen
94,1% Wirksamkeit
basierend auf primärer
Datenanalyse
196 Infektionen
185 in KG: 11 in IG**

2.2 CanSino Biological Inc. and Beijing Institute of Biotechnology

About the vaccine

The **AD5-nCoV** vaccine candidate developed by CanSino Biologics Inc. and the Beijing Institute of Biotechnology is a replication-defective adenovirus type 5 that expresses SARS-CoV-2 spike proteins. The platform (non-replicating viral vector) of AD5-nCoV was originally used for an Ebola vaccine (AD5-EBOV) [30, 31].

AD5-nCoV

Estimated timeline for approval

The first clinical, **phase 1** trial (ChiCTR2000030906/ NCT04313127) with 108 healthy adults is a single-centre dose-escalation study to test both the safety and tolerability of AD5-nCoV injections in three intervention groups using different dosages (low, medium and high). The primary endpoint of the trial is adverse reactions up to seven days post-vaccination. The study is estimated to be completed in December 2022 [32]. A RCT, **phase 2**, double-blinded, placebo-controlled, parallel, three groups trial (ChiCTR2000031781/NCT04398147), aims to evaluate vaccine safety and immunogenicity in healthy adults aged above 18 years. Two intervention groups are using middle or low dose of novel vaccine, and the third group is using placebo. This RCT will be conducted from 2020-04-12 to 2021-01-31.

**Phase 1:
108 gesunde Erwachsene
Dezember 2020**

**Phase 2:
Jänner 2021**

Two new **phase 3** RCTs are registered: a global multicenter, randomized, double-blind, placebo-controlled, adaptive designed clinical trial, to evaluate the efficacy, safety and immunogenicity of Recombinant Novel Coronavirus Vaccine (Adenovirus Type 5 Vector) in adults 18 years old and above, planned to enrol 40,000 participants in Pakistan (NCT04526990), and on 500 participants in Russian federation (NCT04540419). Estimated completion dates are December, 2021 and July, 2021, respectively [33].

**2 neue Phase 3 RCTs
registriert:
40.000 in Pakistan
500 Russland**

bis 2021

Results of publications

The results from **phase 1** study were published (ChiCTR2000030906/NCT04313127) [8]. 108 participants were recruited and received the low dose (n=36), middle dose (n=36), or high dose (n=36) of the vaccine (all were included in the analysis). At least one adverse reaction within the first 7 days after the vaccination was reported in 30 (83%) participants in the low dose group, 30 (83%) participants in the middle dose group, and 27 (75%) participants in the high dose group.

**1 veröffentlichte Phase 1
Studie:
108 Studienteil-
nehmer*innen
erhalten unterschiedliche
Dosierungen**

The results from the above mentioned **phase 2** RCT were published also [9]; 508 eligible participants were randomly assigned to receive the vaccine (1×10^{11} viral particles $n=253$; 5×10^{10} viral particles $n=129$) or placebo ($n=126$). In the 1×10^{11} and 5×10^{10} viral particles dose groups, the RBD-specific ELISA antibodies peaked at 656.5 (95% CI 575.2–749.2) and 571.0 (467.6–697.3), with seroconversion rates at 96% (95% CI 93–98) and 97% (92–99), respectively, at day 28. Both doses of the vaccine induced significant neutralising antibody responses to live SARS-CoV-2, with GMTs of 19.5 (95% CI 16.8–22.7) and 18.3 (14.4–23.3) in participants receiving 1×10^{11} and 5×10^{10} viral particles, respectively. Severe adverse reactions were reported by 24 (9%) participants in the 1×10^{11} viral particles dose group and one (1%) participant in the 5×10^{10} viral particles dose group.

**Publikation der
Phase 2 Studie (RCT)**

508 Teilnehmer*innen

96%/ 97%
**Serokonversionsrate bei
2 unterschiedlichen
Dosierungen**

2.3 University of Oxford/ Astra Zeneca

About the vaccine

The **ChAdOx1 nCoV-19** (AZD1222, AstraZeneca licensed from Oxford University) vaccine candidate developed by the Jenner Institute at Oxford University is based on a non-replicating viral vector. A chimpanzee adenovirus platform is hereby used [34, 35]. The vaccine candidate uses a genetically modified safe adenovirus that may cause a cold-like illness. The intended prevention is through the modified adenovirus producing Spike proteins, eventually leading to the formation of antibodies to the coronavirus's Spike proteins [34].

ChAdOx1 nCoV-19

Estimated timeline for approval

Currently, the first clinical **phase 1/2** single-blinded, placebo-controlled, multi-centre randomised controlled trial to test efficacy, safety and immunogenicity of ChAdOx1 nCoV-19 in 510 healthy adults is ongoing (ISRCTN 15281137/NCT04324606/EudraCT 2020-001072-15). The primary endpoints are number of virologically confirmed symptomatic cases/symptomatic cases of COVID-19 (efficacy) and occurrence of serious adverse events (safety), measured within six months and an optional follow-up visit is offered at day 364. The study is estimated to be completed in May 2021 [36].

Phase 1/2:
510 gesunde Erwachsene

bis Mai 2021

Phase 2b/3 study (EudraCT 2020-001228-32/NCT04400838) is ongoing; the primary endpoint is virologically confirmed (PCR positive) symptomatic COVID-19 infection.

Phase 2b/3 :
laufend

Phase 3 RCT (ISRCTN89951424) is ongoing in Brazil and South Africa, with another country in Africa set to follow, as well as a trial in the US (NCT04516746) [37]. Participants are randomly allocated to receive the investigational vaccine or a well-established meningitis vaccine. Volunteers will be followed for 12 months, and they will be tested for COVID-19 if they develop any symptoms which may represent COVID-19 disease[38]. The study is estimated to be completed in July 2021.

Phase 3 RCT
Brazilien, Südafrika, USA
12-Monate Follow-Up

Ende Juli 2021

A preliminary report with the results from **phase 1/2 RCT** (ISRCTN 15281137/NCT04324606/EudraCT 2020-001072-15) was published [11]. 1077 participants were enrolled and assigned to receive either ChAdOx1 nCoV-19 (n=543) or MenACWY (n=534), ten of whom were enrolled in the non-randomised ChAdOx1 nCoV-19 prime-boost group. Local and systemic reactions were more common in the ChAdOx1 nCoV-19 group (all $p < 0.05$). There were no serious adverse events related to ChAdOx1 nCoV-19. In the ChAdOx1 nCoV-19 group, spike-specific T-cell responses peaked on day 14 (median 856 spot-forming cells per million peripheral blood mononuclear cells, IQR 493–1802; $n=43$). Anti-spike IgG responses rose by day 28 (median 157 ELISA units [EU], 96–317; $n=127$), and were boosted following a second dose (639 EU, 360–792; $n=10$). Neutralising antibody responses against SARS-CoV-2 were detected in 32 (91%) of 35 participants after a single dose when measured in MNA80 and in 35 (100%) participants when measured in PRNT50. After a booster dose, all participants had neutralising activity (nine of nine in MNA 80 at day 42 and ten of ten in Marburg VN on day 56). Neutralising antibody responses correlated strongly with antibody levels measured by ELISA ($R^2=0.67$ by Marburg VN; $p < 0.001$).

Voysey et al. 2020 [12] published results from a pooled interim analysis of four ongoing blinded, randomised, controlled, **phase 2/3 trials** done across the UK, Brazil, and South Africa (ISRCTN89951424, NCT04324606, NCT04400838, and NCT04444674). Participants aged 18 years and older were randomly assigned (1:1) to ChAdOx1 nCoV-19 vaccine or control (meningococcal group A, C, W, and Y conjugate vaccine or saline). Participants in the ChAdOx1 nCoV-19 group received two doses; a subset in the UK trial received a half dose as their first dose (low dose) and a standard dose as their second dose (LD/SD cohort). The primary efficacy analysis included symptomatic COVID-19 in seronegative participants with a nucleic acid amplification test-positive swab more than 14 days after a second dose of vaccine. 23,848 participants were enrolled and 11,636 participants (7548 in the UK, 4088 in Brazil) were included in the interim primary efficacy analysis. In participants who received two standard doses, vaccine efficacy was 62.1% (95% CI 41.0–75.7; 27 [0.6%] of 4440 in the ChAdOx1 nCoV-19 group vs 71 [1.6%] of 4455 in the control group) and in participants who received a low dose followed by a standard dose, efficacy was 90.0% (67.4–97.0; three [0.2%] of 1367 vs 30 [2.2%] of 1374; p -interaction=0.010). Overall vaccine efficacy across both groups was 70.4% (95.8% CI 54.8–80.6; 30 [0.5%] of 5807 vs 101 [1.7%] of 5829).

From 21 days after the first dose, there were ten cases hospitalised for COVID-19, all in the control arm; two were classified as severe COVID-19, including one death. There were 74,341 person-months of safety follow-up (median 3.4 months, IQR 1.3–4.8): 175 severe adverse events occurred in 168 participants, 84 events in the ChAdOx1 nCoV-19 group and 91 in the control group. Three cases of transverse myelitis were initially reported as suspected unexpected serious adverse reactions, with two in the ChAdOx1 nCoV-19 vaccine study arm, triggering a study pause for careful review in each case. Independent clinical review of these cases has indicated that one in the experimental group and one in the control group are unlikely to be related to study interventions, but a relationship remained possible in the third case. Careful monitoring of safety, including neurological events, continues in the trials. The vaccine can be stored and distributed at 2–8°C.

In summary, ChAdOx1 nCoV-19 has an acceptable safety profile and is efficacious against symptomatic COVID-19, with no hospital admissions or

**vorläufige Publikation
Phase 1/2:
1.077 Teilnehmer*innen**

**Antikörper-Response bei
91% bis 100% der
Teilnehmer*innen**

**Phase 2/3 Interimanalyse
basierend auf 11.636
Teilnehmer*innen**

**2 Standard-Dosen:
62,1% Wirksamkeit**

**1 niedrige Dosis
+ 1 Standard Dosis:
90% Wirksamkeit**

**zusammen: 70,4%
1**

**hospitalisierte
Patient*innen: nur in KG**

**NW: gleich verteilt in
KG vs. IG**

**3 Fälle von
Transverser Myelitis
(2 in IG, 1 KG)**

**unwahrscheinlich, dass
mit Impfung assoziiert
Monitoring !**

**gute Verträglichkeit,
Aufbewahrung 2–8°C**

severe cases reported in the ChAdOx1 nCoV-19 arm. The vaccine can be stored and distributed at 2–8°C, making it particularly suitable for global distribution.

Ramasamy et al. 2020 [13] published results from the **phase 2** component of a single-blind, randomised, controlled, phase 2/3 trial -COV002 (ISRCTN90906759, NCT04400838), healthy adults aged 18 years and older were enrolled at two UK clinical research facilities, in an age-escalation manner, into **18–55 years**, **56–69 years**, and **70 years** and older immunogenicity subgroups. The specific objectives of this report were to assess the safety and humoral and cellular immunogenicity of a single-dose and two-dose schedule in adults older than 55 years. 560 participants were enrolled: 160 aged 18–55 years (100 assigned to ChAdOx1 nCoV-19, 60 assigned to MenACWY), 160 aged 56–69 years (120 assigned to ChAdOx1 nCoV-19: 40 assigned to MenACWY), and 240 aged 70 years and older (200 assigned to ChAdOx1 nCoV-19: 40 assigned to MenACWY). **Local and systemic reactions** were more common in participants given ChAdOx1 nCoV-19 than in those given the control vaccine, and similar in nature to those previously reported (injection-site pain, feeling feverish, muscle ache, headache), but were less common in older adults (aged ≥ 56 years) than younger adults. In those receiving two standard doses of ChAdOx1 nCoV-19, after the prime vaccination local reactions were reported in 43 (88%) of 49 participants in the 18–55 years group, 22 (73%) of 30 in the 56–69 years group, and 30 (61%) of 49 in the 70 years and older group, and systemic reactions in 42 (86%) participants in the 18–55 years group, 23 (77%) in the 56–69 years group, and 32 (65%) in the 70 years and older group.

As of Oct 26, 2020, 13 **serious adverse** events occurred during the study period, none of which were considered to be related to either study vaccine. In participants who received two doses of vaccine, median anti-spike SARS-CoV-2 IgG responses 28 days after the boost dose were similar across the three age cohorts (standard-dose groups: 18–55 years, 20 713 arbitrary units [AU]/mL [IQR 13 898–33 550], n=39; 56–69 years, 16170 AU/mL [10233–40353], n=26; and ≥ 70 years 17561 AU/mL [9705–37 796], n=47; p=0.68). Neutralising antibody titres after a boost dose were similar across all age groups (median MNA80 at day 42 in the standard-dose groups: 18–55 years, 193 [IQR 113–238], n=39; 56–69 years, 144 [119–347], n=20; and ≥ 70 years, 161 [73–323], n=47; p=0.40). By 14 days after the boost dose, 208 (>99%) of 209 boosted participants had neutralising antibody responses. T-cell responses peaked at day 14 after a single standard dose of ChAdOx1 nCoV-19 (18–55 years: median 1187 spot-forming cells [SFCs] per million peripheral blood mononuclear cells [IQR 841–2428], n=24; 56–69 years: 797 SFCs [383–1817], n=29; and ≥ 70 years: 977 SFCs [458–1914], n=48).

In summary, in this clinical trial of a vaccine against SARS-CoV-2 tested in an older adult population (aged 18–55 years, 56–69 years, and ≥ 70 years), the vaccine was safe and well tolerated, with reduced reactogenicity in older adults. Antibody responses against the SARS-CoV-2 spike protein were induced in all age groups and were boosted and maintained at 28 days after booster vaccination, including in the 70 years and older group. Cellular immune responses were also induced in all age and dose groups, peaking at day 14 after vaccination.

**Phase 2 in 3 Kohorten
18-55J, 56-69J, >70J
RCT
560 Teilnehmer*innen**

Studie zur Sicherheit

**lokale und systemische
Reaktionen**

häufiger in IG als in KG

**schwerwiegende
Nebenwirkungen:
keine, die auf Impfung
zurückzuführen ist**

**gut verträglich, gute
Immunantworten in allen
3 Kohorten**

2.4 BioNTech/Fosun Pharma/Pfizer

About the vaccine

The **BNT-162** vaccine candidate developed by BioNTech in collaboration with Fosun Pharma and Pfizer is an mRNA platform-based vaccine expressing codon-optimized undisclosed SARS-CoV-2 protein(s) encapsulated in 80-nm ionizable cationic lipid/ phosphatidylcholine/ cholesterol/ polyethylene glycol–lipid nanoparticles [39].

Estimated timeline for approval

A **phase 1/2**, randomized, placebo-controlled, triple-blind, dose-finding, and vaccine candidate-selection study in healthy adults in the US as well as in Germany [40] (**NCT04368728**/EudraCT 2020-001038-36). The study evaluates the safety, tolerability, immunogenicity, and potential efficacy of up to 4 different SARS-CoV-2 RNA vaccine candidates against (COVID-19 BNT162a1, BNT162b1, BNT162b2, and BNT162c2): as a 2-dose or single-dose schedule; at up to 3 different dose levels; in 3 age groups (18 to 55 years of age, 65 to 85 years of age, and 18 to 85 years of age). The study consists of 3 stages: Stage 1: to identify preferred vaccine candidate(s), dose level(s), number of doses, and schedule of administration (with the first 15 participants at each dose level of each vaccine candidate comprising a sentinel cohort); Stage 2: an expanded-cohort stage; and Stage 3; a final candidate/dose large-scale stage. Study NCT04380701 is located in Germany.

Phase 2/3 RCT is ongoing (**NCT04368728**/EudraCT 2020-002641-42) with aim to describe the safety, tolerability, immunogenicity and efficacy of RNA vaccine candidate against COVID-19 in healthy adults (Argentina, Brazil, South Africa, Turkey, US). The candidate selected for evaluation in Phase 2/3 is BNT162b2 (mid-dose). Estimated number of participants is 43998, and completion study date December 2022 [9].

Results of publications

Mulligan et al. 2020 [15] published results from above mentioned **phase 1/2** ongoing study among 45 healthy adults (18–55 years of age) in US, who were randomized to receive 2 doses—separated by 21 days—of 10 µg, 30 µg or 100 µg of BNT162b1 (**NCT04368728**/EudraCT 2020-001038-36). Local reactions and systemic events were dose-dependent, generally mild to moderate, and transient. A second vaccination with 100 µg was not administered because of the increased reactogenicity and a lack of meaningfully increased immunogenicity after a single dose compared with the 30-µg dose. RBD-binding IgG concentrations and SARS-CoV-2 neutralizing titres in sera increased with dose level and after a second dose.

Sahin et al. 2020 published results from a second, non-randomised open-label **phase 1/2** trial in healthy adults, 18-55 years of age in Germany (**NCT04380701**, EudraCT 2020-001038-36) [16], providing a detailed characterisation of antibody and T-cell immune responses elicited by BNT162b1 vaccination. Two doses of 1 to 50 µg of BNT162b1 elicited robust CD4+ and CD8+ T-cell responses and strong antibody responses, with RBD-binding IgG concentrations clearly above those in a COVID-19 human convalescent sample (HCS) panel. Day 43 SARS-CoV-2 serum neutralising geometric mean titers were 0.7-fold (1 µg) to 3.5-fold (50 µg) those of the HCS panel. Immune sera broadly neutralised pseudoviruses with diverse SARS-

BNT-162

**Phase 1 / 2
mehrstufiges
Studiendesign**

**Phase 1/2
(Deutschland)**

November 2022

**Phase 2/3 RCT
läuft derzeit**

**Publikation der
Phase 1 / 2**

**unterschiedliche
Dosierungen**

**weitere Phase
1 / 2 Studie publiziert**

18-55 J

CoV-2 spike variants. Most participants had T helper type 1 (TH1) skewed T cell immune responses with RBD-specific CD8+ and CD4+ T-cell expansion. Interferon (IFN) γ was produced by a high fraction of RBD-specific CD8+ and CD4+ T cells.

Walsh et al. 2020 [41, 42] recently reported, as preprint, additional safety and immunogenicity data from the US **phase 1** trial that supported selection of the vaccine candidate advanced to a pivotal phase 2/3 safety and efficacy evaluation: a direct comparison between BTN126b1 and BTN162b2 (**NCT04368728**) in healthy adults 18–55 and 65–85 years of age. In both younger and older adults, the 2 vaccine candidates elicited similar dose dependent SARS-CoV-2–neutralizing geometric mean titers (GMTs), comparable to or higher than the GMT of a panel of SARS-CoV-2 convalescent sera. BNT162b2 was associated with less systemic reactogenicity, particularly in older adults.

Polack et al. 2020 published results from the **phase 2/3 part** of a global phase 1/2/3, ongoing multinational, placebo-controlled, observer-blinded, pivotal efficacy trial (**NCT04368728**) [17], with randomly assigned persons 16 years of age or older in a 1:1 ratio to receive two doses, 21 days apart, of either placebo or the BNT162b2 vaccine candidate (30 μ g per dose). 43,448 received injections: 21,720 with BNT162b2 and 21,728 with placebo. There were 8 cases of Covid-19 with onset at least 7 days after the second dose among participants assigned to receive BNT162b2 and 162 cases among those assigned to placebo; BNT162b2 was **95% effective** in preventing Covid-19 (95% credible interval, 90.3 to 97.6). Similar vaccine efficacy (generally 90 to 100%) was observed across subgroups defined by age, sex, race, ethnicity, baseline body-mass index, and the presence of coexisting conditions. Among 10 cases of severe Covid-19 with onset after the first dose, 9 occurred in placebo recipients and 1 in a BNT162b2 recipient. The **safety profile** of BNT162b2 was characterized by short-term, mild-to-moderate pain at the injection site, fatigue, and headache. The incidence of serious adverse events was low and was similar in the vaccine and placebo groups.

Publikation zu Sicherheitsdaten zur Auswahl von Kandidaten für Phase 2 / 3

**Phase 2/3
43.448 Teilnehmer*innen**

**8 IG vs. 162 KG
Infektionen**

95% Wirksamkeit

**nur milde bis moderate
Newbenwirkungen**

gut verträglich

2.5 Sinovac Biotech Ltd.

About the vaccine

Sinovac Life Sciences Co., Ltd. is the developer of CoronaVac, an inactivated COVID-19 vaccine candidate, and will be the marketing authorization holder of CoronaVac in China with a vaccine production license from China National Medical Products Administration (NMPA).

CoronaVac

Estimated timeline for approval

The **phase 1 and 2** trials started on April 16, 2020 in Jiangsu Province, China: a group of healthy adults aged 18-59 years old were vaccinated with a 0, 14 day schedule. According to Sinovac announcement, preliminary phase I/II results showed that there was no serious adverse event after vaccinating a total of 743 volunteers in the trials, demonstrating a good safety profile for the vaccine candidate. Over 90% seroconversion was observed in the phase II clinical trial 14 days after completion of a two-dose vaccination at day 0 and day 14. A Phase II study on elderly adults is being conducted which will be

**Phase 1/2 :
743 Teilnehmer*innen**

followed by child and adolescent groups. The phase II trial is expected to be completed at the end of 2020 [43].

A **phase 1/2** RCT on 552 healthy volunteers in China (NCT04551547) aims to evaluate the safety and immunogenicity of the experimental vaccine in healthy children and adolescents aged 3-17 years. Estimated study completion date is September 2021.

Phase 3 RCT (NCT04456595) aims to assess efficacy and safety of the Adsorbed COVID-19 (inactivated) vaccine in health care professionals in Brazil. Estimated number of participants is 8870. The study is double-blind placebo-controlled trial with participants randomly allocated 1:1 to placebo and vaccine arms. The immunization schedule is two doses intramuscular injections (deltoid) with a 14-days interval. All participants will be followed up to 12 months. Interim preliminary efficacy analysis can be triggered by reaching the target number of 150 cases [33]. The study is estimated to be completed in October 2021.

Results of publications

Zhang et al. 2020 published results from randomised, double-blind, placebo-controlled, **phase 1/2** clinical trial (NCT04352608) [21], in healthy adults aged 18–59 years received experimental vaccine: for the phase 1 trial was manufactured using a cell factory process (CellSTACK Cell Culture Chamber 10, Corning, Wujiang, China) , whereas those for the phase 2 trial were produced through a bioreactor process (ReadyToProcess WAVE 25, GE, Umea, Sweden). 144 participants were enrolled in the phase 1 trial, and 600 participants were enrolled in the phase 2 trial. 743 participants received at least one dose of investigational product (n=143 for phase 1 and n=600 for phase 2; safety population). In summary, in this first in-human study of CoronaVac, a phase 1/ 2 study design was used to screen the safety of two doses and two vaccination schedules in a dose-escalation study in a small cohort before expanding the study to a larger cohort to explore the immunogenicity of the vaccine in healthy adults. The immune response in the phase 2 study was substantially higher than in the phase 1 study, which might be due to the difference in preparation process of vaccine batches used in phase 1 and 2 resulting in a higher proportion of intact spike protein on the purified inactivated SARS-CoV-2 virions in the vaccine used in phase 2 than that used in phase 1. Data from this study support the approval of emergency use of CoronaVac in China, and three phase 3 clinical trials that are ongoing in Brazil, Indonesia, and Turkey. The **3 µg dose** of CoronaVac is the suggested dose for efficacy assessment in **future phase 3 trials**. Data from this study support the approval of emergency use of CoronaVac in China, and three phase 3 clinical trials that are ongoing in Brazil, Indonesia, and Turkey.

**Sinovac: Phase 3
RCT in Brazilien
8.870, nur
Gesundsheitspersonal
12-Monate Follow.Up
Oktober 2021**

Phase 1/ 2

**144 Phase 1
600 Phase 2**

**Phase 1: Dosisfindung
Phase 2: Sicherheit**

Phase 3: mit 3 µg Dosis

2.6 China National Pharmaceutical Group Corporation (SINOPHARM)

About the vaccine

The China National Pharmaceutical Group Corporation (SINOPHARM), the state-owned Chinese company, developed a β-propiolactone–inactivated whole-virus vaccine against COVID-19 jointly by the Beijing Institute of Biological Products and the Wuhan Institute of Biological Products under SINOPHARM [18].

inactivated

Estimated timeline for approval

A **phase 3** double-blind, placebo controlled RCT has been initiated (ChiCTR2000034780), to evaluate the protective efficacy of inactivated SARS-CoV-2 Vaccine (Vero Cell) after full course of immunization in preventing diseases caused by the SARS-CoV-2 in healthy subjects aged 18 years old and above. It is currently underway in Abu Dhabi and United Arab Emirates. The study is estimated to be completed in July 2021.

Phase 3 initiiert

Juli 2021

A **phase 3**, randomized, double blind, placebo parallel-controlled clinical trial to evaluate the efficacy, immunogenicity and safety of this vaccine in Argentina, in 3000 healthy participants aged between 18 and 85 years old, is underway also (NCT04560881). The study is estimated to be completed in December 2021.

Dezember 2021

Results of publications

In interim analysis of Xia et al. 2020 [55, 56], related to safety and immunogenicity of an investigational inactivated whole-virus COVID-19 vaccine in China reported results from two double-blind RCTs, **phase 1 and phase 2** (ChiCTR2000031809). The experimental group received a β -propiolactone-inactivated whole-virus vaccine against COVID-19, developed by **Wuhan Institute of Biological Products**. The placebo group contained only sterile phosphate buffered saline and alum adjuvant.

**Phase 1 und
Phase 2 RCTs**

In the **phase 1 RCT**, 96 participants were assigned to 1 of the 3 dose groups (2.5, 5, and 10 μ g/dose) and an aluminum hydroxide (alum) adjuvant-only group (n = 24 in each group), and received 3 intramuscular injections at days 0, 28, and 56. In the **phase 2 RCT** trial, 224 adults were randomized to 5 μ g/dose in 2 schedule groups (injections on days 0 and 14 [n = 84] vs alum only [n = 28], and days 0 and 21 [n = 84] vs alum only [n = 28]). Xia et al. 2020 [20] recently published evidence for the safety and immunogenicity of a SARS-CoV-2 vaccine candidate developed by **China National Biotec Group** and the **Beijing Institute of Biological Products** (BBIBP-CorV), which was tested in randomised, double-blind, placebo controlled **phase 1/2** clinical trials in healthy individuals aged 18 years and older (ChiCTR2000032459). In the phase 1 dose-escalating trial, the vaccine was given at a two-dose schedule at three different concentrations (2 μ g, 4 μ g, and 8 μ g per dose) and was well tolerated in both age groups (18–59 years and ≥ 60 years). The early **phase 2** trial of the BBIBP-CorV vaccine in healthy adults aged 18–59 years assessed the effect of shortening the interval between two doses from 28 days to 14 days or 21 days on the vaccine's immunogenicity. The 4 μ g dose of the vaccine was the most immunogenic when given at the 21-day interval (neutralising antibody titre 283), but its immunogenicity significantly decreased when the interval was reduced to 14 days (neutralising antibody titre 170), suggesting that the interval cannot be shorter than 3 weeks [20, 44].

**Phase 1:
96 Teilnehmer*innen –
3 unterschiedliche
Dosierungen**

**Phase 2:
224 Teilnehmer*innen – 2
unterschiedliche
Zeitpläne**

**Endpunkte: Antikörper
Response**

**weitere Intervalle –
bessere Immunantwort**

2.7 Gamaleya Research Institute

About the vaccine

Vaccine Gam-COVID-Vac, adenoviral-based vaccine against SARS-CoV-2, a solution for intramuscular injection, is a heterologous COVID-19 vaccine consisting of two components, a recombinant adenovirus type 26 (rAd26) vector and a recombinant adenovirus type 5 (rAd5) vector, both carrying the gene for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spike

Gam-COVID-Vac

glycoprotein (rAd26-S and rAd5-S). The trials are sponsored by **Gamaleya Research Institute of Epidemiology and Microbiology, Health Ministry of the Russian Federation** in collaboration with Acellena Contract Drug Research and Development.

Estimated timeline for approval

Based on results of two open, non-randomised **phase 1/2** studies, presented below, according to recent press release, Russian COVID-19 vaccine, called Sputnik V, is the first in the world received national regulatory approval, and was approved for public use even ahead of its Phase III trial.

Phase 3 randomised controlled trial is now underway (NCT04530396). The trial will include 40000 volunteers, with estimated study completion date in May 2021. Phase 3 randomised controlled trial is underway (NCT04564716) in Belarus also, with estimated enrollment of 100 participants.

Results of publications

Two **phase 1/2** studies on healthy adult volunteers (men and women) aged 18–60 years are reported as completed (NCT04436471 and NCT04437875) [33]. In phase 1 of each study, administered intramuscularly on day 0 either one dose of rAd26-S or one dose of rAd5-S and assessed the safety of the two components for 28 days. In phase 2 of the study, which began no earlier than 5 days after phase 1 vaccination, administered intramuscularly a prime-boost vaccination, with rAd26-S given on day 0 and rAd5-S on day 21.

76 participants were enrolled to the two studies (38 in each study). In each study, nine volunteers received rAd26-S in phase 1, nine received rAd5-S in phase 1, and 20 received rAd26-S and rAd5-S in phase 2. Both vaccine formulations were safe and well tolerated. Most adverse events were mild and no serious adverse events were detected. All participants produced antibodies to SARS-CoV-2 glycoprotein. At day 42, receptor binding domain-specific IgG titres were 14 703 with the frozen formulation and 11 143 with the lyophilised formulation, and neutralising antibodies were 49.25 with the frozen formulation and 45.95 with the lyophilised formulation, with a seroconversion rate of 100% [14].

Phase 1/ 2 als Sputnik zugelassen ohne RCT

Phase 3 RCT_ läuft bis Mai 2021

Phase 1/ 2 Studien in Russland

76 Teilnehmer*innen

wenig Nebenwirkungen, Antikörper

2.8 Janssen Pharmaceutical

About the vaccine

The Janssen Pharmaceutical Companies of Johnson & Johnson developed the investigational vaccine (also known as Ad.26.COV2.S), a recombinant vector vaccine that uses a human adenovirus to express the SARS-CoV-2 spike protein in cells.

Estimated timeline for approval

Janssen Pharmaceutical registered **phase 3**, randomised controlled trial (NCT04505722) to demonstrate the efficacy of Ad26.COV2.S in the prevention of molecularly confirmed moderate to severe/critical COVID-19, compared to placebo, in SARS-CoV-2 adult participants. Estimated enrollment is 60,000 participants, with study completion day in March 2023.

Ad.26.COV2.S

Phase 3 RCT mit 60.000 Teilnehmer*innen

März 2023

Results of publications

Sadoff et al. 2020 [45] reported, as preprint, interim results of a **phase 1/2**, double-blind, randomized, placebo-controlled trial related to safety and immunogenicity of the Ad26.COV2.S COVID-19 vaccine candidate (NCT04436276) in healthy adults. Ad26.COV2.S was administered at a dose level of 5x10¹⁰ or 1x10¹¹ viral particles (vp) per vaccination, either as a single dose or as a two-dose schedule spaced by 56 days in healthy adults (18-55 years old; cohort 1a & 1b; n= 402 and healthy elderly >65 years old; cohort 3; n=394). In cohorts 1 and 3 solicited local adverse events were observed in 58% and 27% of participants, respectively. Solicited systemic adverse events were reported in 64% and 36% of participants, respectively.

Phase 1/2
2 Dosierungen
2 Intervalle
3 Kohorten

2.9 Novavax

About the vaccine

The Novavax COVID-19 vaccine being developed by Novavax and co-sponsored by CEPI [46] is a recombinant protein nanoparticle technology platform that is to generate antigens derived from the coronavirus spike (S) protein [47]. Matrix-M™ is Novavax patented saponin-based adjuvant that has the potential to boost the immune system by stimulating the entry of antigen-presenting cells into the injection site and enhancing antigen presentation in local lymph nodes, boosting immune responses [48, 49].

CEPI
Matrix-M™

Estimated timeline for approval

The **phase 1/2**, randomized, placebo-controlled, triple-blind, parallel assignment clinical trial (NCT04368988) in 131 healthy adults aims to evaluate the immunogenicity and safety of SARS-CoV-2 rS nanoparticle vaccine with or without Matrix-M adjuvant in healthy participants ≥ 18 to 59 years of age [33, 50-52]. This RCT will be conducted from May 15, 2020 to July 31, 2021. Estimated Primary Completion Date is December 31, 2020.

Phase 1:
131 gesunde Erwachsene
Juli 2021

A **phase 2b** RCT trial (NCT04533399) started also, to evaluate the effectiveness and safety in South Africans adults; 2904 participants are planned to enrolled, with estimated primary completion date in November 2021 [33].

Phase 2b RCT
2.904 Südafrika
bis 2021

A **phase 3** RCT (EUdraCT 2020-004123-16) is ongoing, in healthy adults in the UK. Main aim is to demonstrate the efficacy of SARS-CoV-2 rS with Matrix-M1 adjuvant in the prevention of virologically confirmed (by polymerase chain reaction [PCR]) to SARS-CoV-2, symptomatic COVID-19, when given as a 2-dose vaccination regimen, as compared to placebo, in serologically negative (to SARS-CoV-2) adult participants. 9000 participants are planned to enrolled.

Phase 3
9.000 Teilnehmer*innen
in UK

Results of publications

A results from above mentioned randomized, placebo-controlled, **phase 1–2 trial** to evaluate the safety and immunogenicity of the rSARS-CoV-2 vaccine (in 5-µg and 25-µg doses, with or without Matrix-M1 adjuvant, and with observers unaware of trial-group assignments) in 131 healthy adults were published [10]. In phase 1, vaccination comprised two intramuscular

Publikation der Phase 1/2
keine schwerwiegenden
NW beobachtet

Results: Vaccines

injections, 21 days apart. After randomization, 83 participants were assigned to receive the vaccine with adjuvant and 25 without adjuvant, and 23 participants were assigned to receive placebo. No serious adverse events were noted. Unsolicited adverse events were mild in most participants; there were no severe adverse events. The two-dose 5- μ g adjuvanted regimen induced geometric mean anti-spike IgG (63,160 ELISA units) and neutralization (3906) responses that exceeded geometric mean responses in convalescent serum from mostly symptomatic Covid-19 patients (8344 and 983, respectively).

3 Results: Therapeutics

EMA is providing guidance to assist developers of potential COVID-19 medicines, to prepare for eventual applications for marketing authorisation. This includes scientific advice, as well as informal consultation with the COVID-19 EMA pandemic Task Force (COVID-ETF). The outcome of any consultation or advice from EMA is not binding on developers. COVID-19 medicines that have received EMA advice can be found in Table 3-1a below, <https://www.ema.europa.eu/en/human-regulatory/overview/public-health-threats/coronavirus-disease-covid-19/treatments-vaccines-covid-19>.

**EMA scientific advice
für viele unterschiedliche
Medikamente**

Table 3--1: COVID-19 medicines that have received EMA advice

Product	Developer	Therapeutic class/drug type	Development stage at time of guidance
VIR-7831, VIR-7832	Vir Biotechnology/GSK	Antiviral (monoclonal antibody)	Clinical phase
UNI911	Union Therapeutics	Antiviral	Clinical phase
Tocilizumab	Roche	Immunomodulator	Clinical phase
SNG-001	Synargein	Immunomodulator	Clinical phase
Siltuximab	EUSApharma	Immunomodulator	Clinical phase
Sarilumab	Sanofi Aventis	Immunomodulator	Clinical phase
Remdesivir	Gilead	Antiviral	Clinical phase
RBT-9	Renibus Therapeutics Inc	Antiviral	Clinical phase
Ravulizumab	Alexion	Other therapeutics	Clinical phase
Otilimab	GSK	Immunomodulator	Clinical phase
Meplazumab	Jiangsu Pacific Meinuoke Biophar.	Antiviral (mAb)	Clinical phase
Mavrilimumab	Kiniksa Pharmaceuticals	Immunomodulator	Clinical phase
Gimsilumab	Roivant	Immunomodulator	Clinical phase
Favipiravir	Glenmark Pharmaceuticals Ltd	Antiviral	Clinical phase
Emapalumab and anakinra	Swedish Orphan Biovitrum AB	Immunomodulator	Clinical phase
Eculizumab	Alexion	Immunomodulator	Clinical phase
Danoprevir	Ascleptis Pharmaceuticals Co Ltd	Antiviral	Clinical phase
Copper chloride	ACOM srl	Antiviral	Clinical phase
Chloroquine and hydroxychloroquine cyclops DPI	PurelMS	Other therapeutics	Clinical phase
Chloroquine	Oxford University	Other therapeutics	Clinical phase
CD24Fc	Oncoimmune Inc	Immunomodulator	Clinical phase
Baricitinib	Eli Lilly	Immunomodulator	Clinical phase
Apremilast	Amgen Europe BV	Immunomodulator	Clinical phase
APN01	Apeiron Biologics	Immunomodulator	Clinical phase
Anti-SARS-CoV-2 polyclonal hyperimmune immunoglobulin	Alliance hyperimmune project (Biotest AG, Bio Products Laboratory, LFB, Octapharma, CSL Behring and Takeda)	Antiviral	Clinical phase
Acalabrutinib	Acerta Pharma BV	Immunomodulator	Clinical phase

In this document we present information for some therapies in development

Table 3 -2: Most advanced therapeutics in the R&D pipeline

Drug	Mechanism of operation	Approval Status Withdrawn, suspended or terminated
Remdesivir (Veklury®)	Antiviral agent	Approved by EMA (conditional marketing authorisation) and FDA (marketing authorisation) 2 RCTs (suspended and terminated)
Favipiravir (Avigan, T-705)	Antiviral agent	No withdrawn, suspended or terminated studies found
Darunavir (Prezista®)	Antiviral agent	No withdrawn, suspended or terminated studies found
Camostat Mesilate (Foipan®)	Antiviral cell-entry inhibitor	No withdrawn, suspended or terminated studies found
APN01 (rhACE2)	Antiviral cell-entry inhibitor	1 RCT – Withdrawn
Tocilizumab (RoActemra®)	Monoclonal antibody	1 RCT withdrawn, 4 RCTs terminated
Sarilumab (Kevzara®)	Monoclonal antibody	1 RCT suspended, 1 RCTs terminated
Interferon beta 1a (SNG001) and 1b	Interferon	1 RCT suspended
Convalescent Plasma	Convalescent Plasma	1 RCT terminated, 1 RCT withdrawn
Plasma derived medicinal products: REGN-COV2; LY-CoV555 (bamlanivimab); LY-CoV016 (etesevimab); AZD7442	Neutralizing monoclonal antibodies	No withdrawn, suspended or terminated studies found
Solnatide	Synthetic peptide	No withdrawn, suspended or terminated studies found
Umifenovir (Arbidol®)	Antiviral agent	No withdrawn, suspended or terminated studies found
Dexamethasone and other corticosteroids	Glucocorticoid	EMA CHMP positive opinion on dexamethasone 2 RCTs terminated, 1 RCT suspended, 1 RCT withdrawn
Anakinra (Kyneret®)	Interleukin 1 receptor antagonist	1 RCT suspended, 1 RCT terminated
Colchicine	An alkaloid, with anti-gout and anti-inflammatory activities	No withdrawn, suspended or terminated studies found
Nafamostat (Futhan©)	Trypsin-like serine protease inhibitor	No withdrawn, suspended or terminated studies found
Gimsilumab	Human monoclonal antibody	No withdrawn, suspended or terminated studies found
Canakinumab	Human monoclonal antibody	No withdrawn, suspended or terminated studies found
Lenzilumab	Recombinant monoclonal antibody	No withdrawn, suspended or terminated studies found
Vitamin D	Vitamin	No withdrawn, suspended or terminated studies found
Baricitinib	Inhibitor of Janus kinase (JAK)1 and JAK2	No withdrawn, suspended or terminated studies found
Molnupiravir	Pro-drug of the nucleoside analogue N4-hydroxycytidine (NHC)	No withdrawn, suspended or terminated studies found

3.1 Remdesivir (Veklury®)

About the drug under consideration

Remdesivir (Veklury) is an antiviral medicine for systemic use which received a **conditional marketing authorisation** in EU in July, 2020 [53-55], https://ec.europa.eu/commission/presscorner/detail/en/mex_20_1266..

Remdesivir (Veklury) is **indicated** for the treatment of coronavirus disease 2019 (COVID-19) in adults and adolescents (aged 12 years and older with body weight at least 40 kg) with pneumonia requiring supplemental oxygen. The drug is for administration by intravenous infusion after further dilution. The **recommended dosage** of remdesivir in patients 12 years of age and older and weighing at least 40 kg is: Day 1 – single loading dose of remdesivir 200 mg given by intravenous infusion, Day 2 onwards – 100 mg given once daily by intravenous infusion. The total **duration of treatment** should be at least 5 days and not more than 10 days. **Concomitant use of remdesivir with chloroquine phosphate or hydroxychloroquine sulphate is not recommended** due to antagonism observed in vitro.

The **most common adverse reaction** in healthy volunteers is increased transaminases (14%). The most common adverse reaction in patients with COVID-19 is nausea (4%) [56].

Remdesivir (Veklury) is subject to **additional monitoring for safety**. Due to a conditional marketing authorisation, Marketing Authorisation Holder (MAH) should complete some **measures to confirm the efficacy and safety within different timeframe** [63].

On October 02, 2020 EMA announced that EMA's safety committee (PRAC) has started a review of a safety signal to assess reports of acute kidney injury in some patients with COVID-19 taking Veklury (remdesivir) [57].

On October 22, 2020 the **U.S. Food and Drug Administration approved** remdesivir for use in adult and pediatric patients 12 years of age and older and weighing at least 40 kilograms (about 88 pounds) for the treatment of **COVID-19 requiring hospitalization**.

The **FDA** recently issued an emergency use authorization (EUA) for the Janus kinase inhibitor **baricitinib** to be used **in combination with remdesivir** in patients with COVID-19 who require oxygen or ventilatory support [58].

Recently, the new **WHO living guidance** on remdesivir for COVID-19 was published [59]. The WHO panel made a conditional recommendation against the use of remdesivir in hospitalized patients with COVID-19, regardless of disease severity, with new information and recommendations on remdesivir after publication of results from the WHO SOLIDARITY trial [60]. The recommendation on remdesivir was informed by results from a systematic review and network meta-analysis (NMA) that pooled data from four randomized trials with 7333 participants hospitalized for COVID-19. The resulting GRADE evidence summary suggested that remdesivir has possibly no effect on mortality (odds ratio 0.90, 95% confidence interval [CI] 0.70 - 1.12; absolute effect estimate 10 fewer deaths per 1000 patients, 95% CI from 29 fewer - 11 more deaths per 1000 patients; low certainty evidence); and possibly no effect on the other important outcomes identified by the panel, with similar low to very low certainty of evidence. The panel judged the

**erstes zugelassenes
antivirales Medikament
gegen Coronavirus:
conditional marketing
authorisation**

**indiziert für Patient*innen
≥ 12 Jahre mit
Lungenentzündung,
Sauerstoff-unterstützt
Verabreichung iv
5-10 Tage**

Nebenwirkungen

**Okt 2020:
EMA Sicherheitsanalyse**

**FDA Zulassung im
Okt 2020**

**FDA Notzulassung für
Kombinationstherapie
Remdesivir + Baricitinib**

**WHO empfiehlt
Remdesivir nicht,
unabhängig von
Patientenpopulation
basierend auf Ergebnisse
aus SOLIDARITY**

overall credibility of subgroup analyses assessing differences in mortality by severity of illness to be insufficient to make subgroup recommendations.

US COVID-19 **Treatment Guidelines Panel** issued new recommendations on remdesivir treatment for patients with COVID-19 (as of December 3, 2020) [61]:

Remdesivir, an antiviral agent, is currently the only drug that is approved by the FDA for the treatment of COVID-19. It is recommended for use in hospitalised patients who require supplemental oxygen. However, it is not routinely recommended for patients who require mechanical ventilation due to the lack of data showing benefit at this advanced stage of the disease.

Gilead Sciences Inc. said it plans to start human trials of an inhaled version of its anti-Covid-19 drug remdesivir. An inhaled version, through a nebulizer, could allow Gilead to give the drug to a broader group of patients, including those with milder symptomatic cases who don't need to be hospitalized, <https://www.pharmacist.com/article/gilead-begin-human-testing-inhaled-version-covid-19-drug-remdesivir>.

Withdrawn, suspended or terminated studies

The two phase 3 randomised controlled trials (RCT) to evaluate intravenous RVD in patients with 2019-nCoV, initiated in the beginning of February in China, are suspended (NCT04252664) or terminated (NCT04257656) (the epidemic of COVID-19 has been controlled well in China, and no eligible patients can be enrolled further).

Results of publications

Wang Y et al. 2020 [62] published results of the first randomised, double-blind, placebo-controlled, multicentre trial, conducted in China (NCT04257656), on intravenous remdesivir in adults admitted to hospital with severe COVID-19. The study was terminated before attaining the prespecified sample size (237 of the intended 453 patients were enrolled) because the outbreak of COVID-19 was brought under control in China. Remdesivir treatment was not associated with a statistically significant difference in time to clinical improvement (hazard ratio 1.23 [95% CI 0.87–1.75]); duration of invasive mechanical ventilation; viral load; adverse events.

Beigel et al. 2020 [63] reported results from double-blind, randomized, placebo-controlled trial of intravenous remdesivir in 1062 adults hospitalized with Covid-19 (541 assigned to remdesivir and 521 to placebo) (NCT04280705). Remdesivir group had a median recovery time of 10 days (95% confidence interval [CI], 9 to 11) vs 15 days (95% CI, 13 to 18) among placebo group (rate ratio for recovery, 1.29; 95% CI, 1.12 to 1.49; P<0.001, by a log-rank test). The rate ratio for recovery was largest among patients with a baseline ordinal score of 5 (rate ratio for recovery, 1.45; 95% CI, 1.18 to 1.79). The Kaplan–Meier estimates of mortality were 6.7% with remdesivir vs 11.9% in placebo group by day 15 (hazard ratio, 0.55; 95% CI, 0.36 to 0.83); 11.4% with remdesivir vs 15.2% with placebo by day 29 (hazard ratio, 0.73; 95% CI, 0.52 to 1.03). The between group differences in mortality varied considerably according to baseline severity, with the statistically significant difference seen among patients with a baseline ordinal score of 5 (hazard ratio, 0.30; 95% CI, 0.14 to 0.64). Serious adverse events were reported in 131 of the 532 patients who received remdesivir (24.6%) and in 163 of the 516 patients who received placebo (31.6%). There were 47 serious respiratory failure adverse events in

US COVID-19 Treatment Guidelines

Empfehlung: nicht routinemäßig

Vorhaben von Gilead: Darreichungsform mittels Inhalator

in ClinicalTrials.gov & EUdraCT keine weiteren beendeten Studien

Ergebnisse der Studien:

Wang (Hubei/ China): frühzeitig beendet wegen Mangel an Pts.

keine Unterschiede bei klinischer Verbesserung, invasiver Beatmung

Beigel (USA) 1.062 Pts. kürzere Dauer zur Erholung

Unterschiede bei Baseline-Schwergrad erschweren die Interpretation der Mortalitätsdaten

the remdesivir group (8.8% of patients), including acute respiratory failure and the need for endotracheal intubation, and 80 in the placebo group (15.5% of patients). No deaths were considered by the investigators to be related to treatment assignment.

Goldman et al. 2020 [64] published the results from the randomized, open-label, phase 3 trial involving 397 hospitalized patients with confirmed SARS-CoV-2 infection, oxygen saturation of 94% or less while they were breathing ambient air, and radiologic evidence of pneumonia (NCT04292899), to receive intravenous remdesivir for either 5 days or 10 days. Trial did not show a significant difference between a 5-day course and a 10-day course of remdesivir. -The most common adverse events were nausea (9% of patients), worsening respiratory failure (8%), elevated alanine aminotransferase level (7%), and constipation (7%). The absence of a control group in this study did not permit an overall assessment of the efficacy of remdesivir.

Spinner et al. 2020 [65] published results from a randomised, open-label, phase 3 trial (NCT04292730) performed on 596 hospitalised patients with moderate COVID-19 pneumonia (pulmonary infiltrates and room-air oxygen saturation >94%). Patients were randomized in a 1:1:1 ratio to receive a 10-day course of remdesivir (n = 197), a 5-day course of remdesivir (n = 199), or standard care (n = 200). On day 11, patients in the 5-day remdesivir group had statistically significantly higher odds of a better clinical status distribution vs standard care (odds ratio, 1.65; 95% CI, 1.09-2.48; p=0.02), but the difference was of uncertain clinical importance. The clinical status distribution on day 11 between the 10-day remdesivir and standard care groups was not significantly different (p=0.18 by Wilcoxon rank sum test).

There were no significant differences between the 5-day or 10-day remdesivir groups and standard care for any of the exploratory end points—time to 2-point or greater improvement in clinical status, time to 1-point or greater improvement in clinical status, time to recovery, time to modified recovery, and time to discontinuation of oxygen support, duration of oxygen therapy or hospitalization and all-cause mortality at day 28. The difference in AEs proportions between the 5-day remdesivir group and standard care was not statistically significant (4.8%; 95% CI, -5.2% to 14.7%; p=0.36), but the difference between the 10-day remdesivir group and standard care was significant (12.0%; 95% CI, 1.6%-21.8%; p=0.02). Nausea (10% vs 3%), hypokalemia (6% vs 2%), and headache (5% vs 3%) were more frequent among remdesivir-treated patients compared with standard care. Serious adverse events were less common in the remdesivir groups, but the difference was not statistically significant.

Interim results from the **WHO SOLIDARITY trial (ISRCTN83971151, NCT04315948)**, large, international, adaptive, open-label, randomized controlled trial to evaluate remdesivir, lopinavir/ritonavir, interferon beta-1a and hydroxychloroquine treatment for COVID-19, were published, with 2750 patients allocated to remdesivir [60, 66]. Death rate ratio was not statistically significant different between remdesivir and standard care; RR=0.95 (0.81-1.11, p=0.50; 301/2743 active vs 303/2708 control). The same was true for the outcomes: initiation of ventilation and hospitalisation duration, and other three investigation treatment.

**Goldman (USA, IT, SP...)
RCT, open-label
397 Pts.**

**Vergleich von 5 vs. 10
Tagen RDV**

**primärer Endpunkt:
klinischer Status
am Tag 14**

**Spinner (USA, Europa,
Asien)**

**5-Tage vs
10-Tage vs.
SOC**

**596 Pts
kein signifikanter
Unterschied zwischen
5 vs. 10 Tage vs. SOC**

**AE signifikanter
Unterschied zwischen 10
Tage vs. SOC zu
Ungunsten von
Remdesivir
SAE häufiger in
SOC Gruppe**

WHO SOLIDARITY

**kein Unterschied
bei Mortalität
kein Unterschied bei
anderen Endpunkten**

Based on the **living synthesis** of currently available scientific evidence from **4 RCTs** (Wang, Beigel, Spinner and SOLIDARITY-Remdesivir), on remdesivir compared with standard care/placebo, presented in recently published EUnetHTA Rapid Collaborative Review document [67], current scientific conclusions were listed: According to the results of four RCTs with moderate certainty of evidence, remdesivir has no effect on mortality in COVID-19 patients compared to standard treatment; According to the results of three RCTs, remdesivir decreases the incidence of WHO progression score level 6 or above (moderate certainty of evidence), as well as the WHO progression score level 7 or above D14-D28 (high certainty of evidence), compared to standard treatment; According to the results of one RCT with very low certainty of evidence, remdesivir has no effect on viral clearance, compared to standard treatment; According to the results of three RCTs with moderate certainty of evidence, remdesivir increases the number of discharged patients within 28 days compared to standard treatment; According to low certainty of evidence, remdesivir has no effect on outcomes mechanical ventilation (4 RCTs); time to clinical improvement (3 RCTs); duration of ventilation (2RCTs); duration of hospitalisation (3 RCTs) and serious adverse events leading to discontinuation (3 RCTs), compared to standard treatment; According to the results of two RCTs with high certainty of evidence, remdesivir does not increase adverse events compared to standard treatment; According to the results of three RCTs with moderate certainty of evidence, remdesivir decreases the number of patients with SAEs compared to standard treatment.

Details can be found in the Summary of findings Table 3.1-1.

The Living Systematic Review with Meta-Analysis (MA), related to Remdesivir 5 days vs Remdesivir 10 days (2 RCTs, Spinner and Goldman) and the Summary of findings table (https://covid-nma.com/living_data/index.php) are presented in Table 3.1-2.

**EUnetHTA Bericht
zu 4 RCTs (Dez 2020):**

**kein Unterschied:
all-cause mortality**

**Unterschied bei klinischer
Verbesserung und bei
Nebenwirkungen**

Results: Therapeutics

*Table 3.1-1: Summary of findings table on **Remdesivir vs Standard care /Placebo** (4 RCTs: Wang, Beigel, Spinner, SOLIDARITY-Remdesivir)
Patient or population: Mild/Moderate/Severe/Critical COVID-19*

Setting: Worldwide

Intervention: Remdesivir

Comparison: Standard Care/Placebo

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect difference (95% CI)	Number of participants (studies)	Certainty of evidence ^e (GRADE)	Comments
	Risk with Standard care ^a	Risk with Remdesivir					
All-cause Mortality^b	112 per 1.000	101 per 1.000 (82 to 125)	RR 0.90 (0.73 to 1.11)	11 fewer per 1.000 (from 30 fewer to 12 more)	7345 (4 RCTs) Spinner, 2020; SOLIDARITY 2020; Beigel, 2020; Wang, 2020[68][68]	⊕⊕⊕○ MODERATE	Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of events
Clinical improvement D14-D28^b	759 per 1.000	805 per 1.000 (751 to 858)	RR 1.06 (0.99 to 1.13)	46 more per 1.000 (from 8 fewer to 99 more)	832 (2 RCTs) Spinner, 2020; Wang, 2020	⊕⊕⊕○ MODERATE	Imprecision downgraded by 1 level: due to low number of events and/or participants
WHO progression score (level 6 or above) D14-D28^b	193 per 1.000	131 per 1.000 (106 to 164)	RR 0.68 (0.55 to 0.85)	62 fewer per 1.000 (from 87 fewer to 29 fewer)	1894 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕⊕○ MODERATE	Risk of bias downgraded by 1 level: some concerns due to deviation from intended intervention and outcome measurement
WHO progression score level 7 or above D14-28^b	178 per 1.000	124 per 1.000 (100 to 156)	RR 0.70 (0.56 to 0.88)	53 fewer per 1.000 (from 78 fewer to 21 fewer)	1894 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕⊕⊕ HIGH	
Viral negative conversion D7^b	492 per 1.000	502 per 1.000 (374 to 679)	RR 1.02 (0.76 to 1.38)	10 more per 1.000 (from 118 fewer to 187 more)	196 (1 RCT) Wang, 2020	⊕○○○ VERY LOW	Risk of bias downgraded by 1 level: some concerns with missing data

Results: Therapeutics

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect difference (95% CI)	Number of participants (studies)	Certainty of evidence ^e (GRADE)	Comments
	Risk with Standard care ^a	Risk with Remdesivir					
							Indirectness downgraded by 1 level: despite a multicenter design this is a single study from a single country, therefore results in this population might not be generalizable to other settings Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of events
Adverse events^b	583 per 1.000	542 per 1.000 (496 to 589)	RR 0.93 (0.85 to 1.01)	41 fewer per 1.000 (from 87 fewer to 6 more)	1894 (2 RCTs) Wang, 2020; Beigel, 2020;	⊕⊕⊕⊕ HIGH	Presume that the adverse event rates, and the corresponding relative risks, are similar across diverse settings; therefore not downgraded for indirectness
Serious adverse events^b	40 per 1.000	24 per 1.000 (15 to 38)	RR 0.60 (0.38 to 0.96)	16 fewer per 1.000 (from 25 fewer to 2 fewer)	1894 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕⊕○ MODERATE	Presume that the adverse event rates, and the corresponding relative risks, are similar across diverse settings; therefore not downgraded for indirectness Imprecision downgraded by 1 level: few events and a wide confidence interval consistent with the possibility of a benefit and the possibility of no effect.

Results: Therapeutics

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect difference (95% CI)	Number of participants (studies)	Certainty of evidence ^e (GRADE)	Comments
	Risk with Standard care ^a	Risk with Remdesivir					
Serious adverse events leading to discontinuation^c	15 per 1.000	15 per 1000	OR 1.00 (0.37 - 3.83)	0 fewer per 1.000 (from 9 fewer to 40 more)	1894 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕○○ Low	Very serious imprecision
Mechanical ventilation^c	105 per 1000	95 per 1000	OR: 0.89 (0.76 - 1.03)	10 fewer per 1000 (from 23 fewer to 3 more)	6549 (4 RCTs) Spinner, 2020; SOLIDARITY, 2020; Beigel, 2020; Wang, 2020	⊕⊕○○ Low	Due to serious risk of bias and serious imprecision
Duration of ventilation^c	14.7 Days mean	13.4 Days mean	Measured by: Scale: lower better	Difference: MD 1.3 lower (from 4.1 lower to 1.5 higher)	440 (2 RCTs) Wang, 2020; Beigel, 2020;	⊕⊕○○ Low	Due to very serious imprecision
Time to clinical improvement^c	11.0 Days mean	9.0 Days mean	Measured by: Scale: lower better	Difference: MD 2.0 lower (from 4.2 lower to 0.9 higher)	1882 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕○○ Low	Due to serious imprecision and serious indirectness
Duration of hospitalization^c	12.8 Days mean	12.3 Days mean	Measured by: Scale: lower better	Difference: MD 0.5 lower (from 3.3 lower to 2.3 higher)	1882 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕○○ Low	Due to serious imprecision and serious indirectness

Results: Therapeutics

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect difference (95% CI)	Number of participants (studies)	Certainty of evidence ^e (GRADE)	Comments
	Risk with Standard care ^a	Risk with Remdesivir					
Number of patients discharged within 28 days^d	478 per 1,000	540 per 1,000 (488 to 593)	RR 1.13 (1.02 to 1.24)	62 more per 1,000 (from 10 more to 115 more)	1894 (3 RCTs) Beigel, 2020; Spinner, 2020; Wang, 2020	⊕⊕⊕○ MODERATE	Downgraded of one level for high risk of performance bias in two studies and unclear risk of selection, attrition and reporting bias in one study

Source: [67] [65] [60] [63] [62]

a Background risk in the control group is based on the observed risk in the studies; b outcome data and GRADE assessment from Covid-nma.com, https://covid-nma.com/living_data/index.php (The evidence profile and summary of findings table were updated on November 17th, 2020); c Outcome data and GRADE assessment from WHO guideline [59] d Outcome data and GRADE assessment from the department of Epidemiology Lazio Regional Health Service (DEPLazio), Italy, <http://deplazio.net/farmacicovid/index.html>; e GRADE Working Group grades of evidence: High certainty=we are very confident that the real effect is close to that of the estimated effect; Moderate certainty=we are moderately confident in the effect estimation: the real effect may be close to the estimated effect, but there is a possibility that it is substantially different; Low certainty=our confidence in the effect estimation is limited: the real effect may be substantially different from the estimated effect; Very Low certainty=we have very little confidence in estimating the effect: the actual effect is likely to be substantially different from the estimated one.

Abbreviations: CI= confidence interval; RR=relative risk; OR=odds ratio.

GRADE Working Group grades of evidence: **High certainty:** We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Results: Therapeutics

Table 3.1-2: Summary of findings table on **Remdesivir 5 days vs Remdesivir 10 days** (2 RCTs: Goldman, Spinner) - https://covid-nma.com/living_data/index.php

Remdesivir 5 days compared to Remdesivir 10 days for Mild/Moderate/Critical/Severe Covid-19

Patient or population: Mild/Moderate/Critical/Severe Covid-19

Setting: Worldwide

Intervention: Remdesivir 5 days

Comparison: Remdesivir 10 days

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Remdesivir 10 days	Risk with Remdesivir 5 days				
Incidence of viral negative conversion D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
Incidence of clinical improvement D7	368 per 1,000	438 per 1,000 (371 to 515)	RR 1.19 (1.01 to 1.40)	798 (2 RCTs) ^b	⊕⊕○○ LOW ^{e,g}	
Incidence of clinical improvement D14-28	708 per 1,000	750 per 1,000 (616 to 920)	RR 1.06 (0.87 to 1.30)	798 (2 RCTs) ^b	⊕○○○ VERY LOW ^{c,d,f}	
Incidence of WHO progression score (level 6 or above) D14-28	174 per 1,000	109 per 1,000 (78 to 153)	RR 0.63 (0.45 to 0.88)	798 (2 RCTs) ^b	⊕⊕○○ LOW ^{e,g}	
Incidence of WHO progression score (level 7 or above) D14-28	146 per 1,000	85 per 1,000 (58 to 124)	RR 0.58 (0.40 to 0.85)	798 (2 RCTs) ^b	⊕⊕○○ LOW ^{e,g}	
All-cause mortality D14-28	60 per 1,000	45 per 1,000 (29 to 81)	RR 0.74 (0.41 to 1.34)	798 (2 RCTs) ^b	⊕⊕○○ LOW ^{e,g}	
Adverse events	650 per 1,000	604 per 1,000 (546 to 669)	RR 0.93 (0.84 to 1.03)	798 (2 RCTs) ^b	⊕⊕⊕○ MODERATE ^e	
Serious adverse events	196 per 1,000	126 per 1,000 (92 to 171)	RR 0.64 (0.47 to 0.87)	798 (2 RCTs) ^b	⊕⊕○○ LOW ^{e,g}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

a. Last update: September 18, 2020; b. Spinner CD, 2020; Goldman JD, 2020; c. Risk of bias downgraded by 1 level: some concerns due to concerns during the randomization process, deviation from intended intervention and outcome measurement; d. Imprecision downgraded by 1 level: due to low number of events and/or participants; e. Inconsistency downgraded by 1 level: I²= 79.3%
f. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm; g. Risk of bias downgraded by 1 level: some concerns due to concerns during the randomization process and deviation from intended intervention

3.2 Lopinavir + Ritonavir (Kaletra®)

Due to the lack of effectiveness of lopinavir/ritonavir in treating adults hospitalized with COVID-19 patients and the decisions to stop enrolling participants to the lopinavir/ritonavir (Kaletra) arms of the RECOVERY, SOLIDARITY and DISCOVERY studies in adults hospitalized with COVID-19, our reporting related to lopinavir/ritonavir was stopped also.

Last reporting V6/September 2020:

https://eprints.aihta.at/1234/50/Policy_Brief_002_Update_09.2020.pdf

wegen erwiesenem Mangel an Wirksamkeit wurde Beobachtung beendet

3.3 Favipiravir (Avigan®)

About the drug under consideration

Favipiravir (Avigan®), an antiviral drug, is a new type of RNA-dependent RNA polymerase (RdRp) inhibitor [69, 70].

antivirales Medikament

Favipiravir (Avigan®) has not been approved by the European Medicines Agency (EMA) or the American Food and Drug Administration (FDA) for COVID-19.

The US COVID-19 Treatment Guidelines Panel **recommends against** using the **Lopinavir/ritonavir (AI) or other HIV protease inhibitors (AIII)**, except in a clinical trial, because of unfavorable pharmacodynamics and because clinical trials have not demonstrated a clinical benefit in patients with COVID-19 [61].

Empfehlungen des US COVID-19 Treatment Guidelines Panel GEGEN jegliche HIV Protease Inhibitoren

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated RCTs were found in two clinical trial registers (ClinicalTrials.gov and EUdraCT).

Results of publications

Chen C et al. 2020 [71] published results (as preprint) on a RCT (ChiCTR2000030254) related to efficacy and safety of favipiravir, **in comparison with umifenovir**. Summary of findings table on favipiravir compared to umifenovir (1 RCT: Chen) is presented in Table 3.3-1.

1 Publikation zu RCT Vergleich mit Umifenovir

Lou Y et al. 2020, published as preprint results of exploratory RCT with 3 arms (ChiCTR2000029544) [72] related to the efficacy and safety of favipiravir **in comparison with baloxavir marboxil, and lopinavir + ritonavir or darunavir/cobicistat + umifenovir + interferon-a** in hospitalized adult patients with COVID-19. The percentage of patients who turned viral negative after 14-day treatment was 70%, 77%, and 100% in the baloxavir, favipiravir, and control group respectively, with the medians of time from randomization to clinical improvement was 14, 14 and 15 days, respectively.

1 weitere Publikation Vergleich mit Baloxavir marboxil

Summary of findings table on favipiravir compared to baloxavir marboxil is presented in Table 3.3-2 and favipiravir compared to lopinavir + ritonavir or darunavir/cobicistat + umifenovir + interferon-a (1 RCT: Lou 2020) [69] is presented in Table 3.3-3.

Interim results from an adaptive, multicenter, open label, randomized, phase 2/3 clinical trial (NCT04434248) of favipiravir (AVIFAVIR) versus standard of care (SOC) in 60 hospitalized patients with moderate COVID-19 pneumonia were published (three treatment groups: AVIFAVIR 1600/600 mg, AVIFAVIR 1800/800 mg, or SOC). AVIFAVIR enabled SARS-CoV-2 viral clearance in 62.5% of patients within 4 days, and was safe and well-tolerated. Based on these interim results, the Russian Ministry of Health granted a conditional marketing authorization to AVIFAVIR, which makes it the only approved oral drug for treatment of moderate COVID-19 to date [73].

AVIFAVIR
Phase 2/3 RCT bei
moderater Covid-19
Erkrankung

interim Auswertung
orale Verabreichung in
Russland „conditional“
zugelassen

Dabbous HM et al. 2020 published results, as preprint, from open-label, phase 3 RCT, comparing favipiravir vs standard care (hydroxychloroquine plus oseltamivir) in 100 patients with mild to moderate COVID-19 in Egypt (NCT04349241) [74]. No statistically significant difference was found related to time to PCR negativity ($p=0.7$). Four patients in favipiravir group had increase in liver transaminase, and 20 patients in standard care group (hydroxychloroquine plus oseltamivir) developed heartburn and nausea. One patient died in hydroxychloroquine plus oseltamivir group after acute myocarditis resulted in acute heart failure.

Phase 3 RCT (Ägypten)
kein Unterschied

Balykova et al. 2020 [75] published results from a RCT in 200 hospitalised patients with COVID-19 showed a significant advantage of favipiravir therapy compared with standard therapy in terms of the rate of improvement in clinical status (on average by 4 days), the speed and frequency of recovery on the 10 day of therapy (no clinical signs of the disease in the study and control groups were observed in 44 and 10% of patients, respectively), the frequency of achieving the viral clearance on the 10th day of therapy (98 and 78% in the study and control groups, respectively) ($p=0.00003$). Favipiravir therapy was accompanied by a significant improvement in lung condition according to CT data, improved laboratory parameters and normalization of oxygen saturation levels. Favipiravir therapy was characterized by a favorable safety profile. In the main group, no aggravation of the course of the disease or serious adverse events related to the drug were recorded.

RCT
200 hospitalisierte
Patient*innen

raschere klinische
Verbesserung (-4 Tage),
insb. der Lunge

akzeptables
Sicherheitsprofil

Ruzhentsova et al. 2020 [76] published results as preprint from open-labeled, randomized, active-controlled multicenter trial (NCT04501783) of an oral dosage form of favipiravir in out- and hospitalized patients with mild to moderate COVID-19 in 10 clinical centers in Russia. 190 Patients were randomly assigned (in a 2:1 ratio) to receive either favipiravir (1800 mg BID on day 1, followed by 800 mg BID for up to 9 days), or standard of care (SOC) treatment (umifenovir + intranasal interferon alpha-2b, or hydroxychloroquine) for up to 10 days. The median time to clinical improvement was 6.0 (IQR 4.0; 9.3) days in favipiravir group and 10.0 (IQR 5.0; 21.0) days in SOC group; the median difference was 4 days (HR 1.63; 95% CI 1.14-2.34, $p=0.007$). The statistically significant difference in the median time to viral clearance was observed only in the hospitalized cohort of patients: 3.0 (IQR 3.0; 3.0) vs. 5.0 (IQR 4.5; 5.5), respectively (HR 2.11; 95% CI 1.04-4.31; $p=0.038$). However, the rate of viral elimination on Day 5 in the favipiravir group was significantly higher in the whole population: 81.2% vs. 67.9% respectively (RR 1.22; 95% CI 1.00-1.48; $p=0.022$). The rate of clinical improvement on Day 7 in the favipiravir group was 1.5-fold higher compared to SOC: 52.7% vs. 35.8% (RR 1.50; 95% CI 1.02-2.22; $p=0.020$). Favipiravir was well tolerated: most of the adverse events (AE) were mild. Any AEs were reported in 74.1% of patients in the favipiravir group vs. 60.0% in the SOC group; the most common adverse reactions were asymptomatic hyperuricemia,

RCT
190 Patient*innen
milde oder moderate
Erkrankung

ambulante oder
hospitalisiert

Vergleich mit SOC
(umifenovir + intranasal
interferon alpha-2b, or
hydroxychloroquine)

raschere Reduktion der
Viruslast und
klinische Verbesserung
mit favipiravir

akzeptables
Sicherheitsprofil

transient elevation of ALT & AST, and gastrointestinal disorders (diarrhea, nausea, abdominal pain).

Udwadia et al. 2020 [77] published results from randomized, open-label, parallel-arm, multicenter, phase 3 trial (CTRI/2020/05/025114), in adults with mild to moderate COVID-19 in India. 150 patients were randomized to favipiravir (n=75) or control (n=75). Median time to cessation of viral shedding was 5 days (95% CI: 4 days, 7 days) versus 7 days (95% CI: 5 days, 8 days), p=0.129, and median time to clinical cure was 3 days (95% CI: 3 days, 4 days) versus 5 days (95% CI: 4 days, 6 days), p=0.030, for favipiravir and control respectively. Adverse events were observed in 36% of favipiravir and 8% of control patients. One control patient died due to worsening disease.

RCT
150 Patient*innen
milde oder moderate
Erkrankung

raschere Reduktion der
Viruslast und
klinische Verbesserung
mit favipiravir

Data related to **Summary of findings table on favipiravir compared to standard care** (6 RCTs: Lou 2020, Ivashchenko 2020, Dabbous 2020, Balykova 2020, Ruzhentsova 2020, Udwadia 2020) could be found in Table 3.3-4 below. Only one clinical outcome was statistically significant different in favour of favipiravir, based on results from 3 RCTs - Clinical improvement D7, RR 1.58 (1.15 to 2.16), with low certainty of evidence.

Doi et al. 2020 published results from RCT (Japan Registry of Clinical Trials **jRCTs041190120**), related to early versus late favipiravir in hospitalised patients with COVID-19 [78]. 88 patients were randomly assigned at a 1:1 ratio to **early or late favipiravir therapy** (the same regimen starting on day 6 instead of day 1). Viral clearance occurred within 6 days in 66.7% and 56.1% of the early and late treatment groups (adjusted hazard ratio [aHR], 1.42; 95% confidence interval [95% CI], 0.76–2.62). Of 30 patients who had a fever ($\geq 37.5^{\circ}\text{C}$) on day 1, time to defervescence was 2.1 days and 3.2 days in the early and late treatment groups (aHR, 1.88; 95%CI, 0.81–4.35). During therapy, 84.1% developed transient hyperuricemia. Neither disease progression nor death occurred to any of the patients in either treatment group during the 28-day participation.

Okt 2020:
RCT mit 89 Pts.
Japan
Vergleich von früher und
später Favipiravir Therapie
bei hospitalisierten Pts.

kein Unterschied

Zhao H et al. 2020, published results from RCT in moderate to critical COVID-19 patients in China, comparing **favipiravir to tocilizumab and favipiravir plus tocilizumab (ChiCTR2000030096, NCT04310228)** [79]. Patients were randomly assigned (3:1:1) to a 14-day combination of favipiravir combined with tocilizumab (combination group), favipiravir, and tocilizumab. The cumulative lung lesion remission rate at day 14 was significantly higher in the combination group as compared with favipiravir group (p = 0.019, HR 2.66 95% CI [1.08 to 6.53]); a significant difference between tocilizumab and favipiravir found also (p = 0.034, HR 3.16, 95% CI 0.62 to 16.10). There was no significant difference between the combination group and the tocilizumab group (p = 0.575, HR 1.28 95%CI 0.39 to 4.23). Combined therapy can also significantly relieve clinical symptoms and help blood routine to return to normal. No serious adverse events were reported.

RCT
19 Patient*innen
favipiravir + tocilizumab vs.
favipiravir vs. tocilizumab

Kombinationstherapie
von Vorteil

Table 3.3-1: Summary of findings table on **favipiravir compared to umifenovir** (1 RCT: Chen) - https://covid-nma.com/living_data/index.php

Summary of findings:						
Favipiravir compared to Umifenovir for COVID-19						
Patient or population: COVID-19						
Setting: Worldwide						
Intervention: Favipiravir						
Comparison: Umifenovir						
Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Umifenovir	Risk with Favipiravir				
Incidence viral negative conversion D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement - not reported	-	-	-	-	-	outcome not yet measured or reported
Incidence of clinical recovery D7	517 per 1.000	594 per 1.000 (470 to 744)	RR 1.15 (0.91 to 1.44)	240 (1 RCT)	⊕○○○ VERY LOW ^{a,b,c}	
Incidence of WHO progression score (level 6 or above) - not reported	-	-	-	-	-	outcome not yet measured or reported
Incidence of WHO progression score (level 7 or above) - not reported	-	-	-	-	-	outcome not yet measured or reported
All-cause mortality D7				240 (1 RCT)	⊕○○○ VERY LOW ^{b,d,e}	zero events in both groups
Adverse events D7	275 per 1.000	358 per 1.000 (245 to 523)	RR 1.30 (0.89 to 1.90)	240 (1 RCT)	⊕⊕○○ LOW ^{a,c,f}	

Results: Therapeutics

Serious adverse events D7	240 (1 RCT)	⊕○○○ VERY LOW ^{a,d,f}	zero events in both groups
---------------------------	----------------	--------------------------------------	-------------------------------

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

- a. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, deviations from intended interventions and outcome measurement
- b. Indirectness downgraded by 1 level: single study from a single country, therefore results in this population might not be generalizable to other settings
- c. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants
- d. Imprecision downgraded by 2 levels: no events in both groups and low number of participants
- e. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization and deviations from intended interventions
- f. We presume that the adverse event rates, and the corresponding relative risks, is similar across diverse settings; therefore not downgraded for indirectness

Results: Therapeutics

Table 3.3-2: Summary of findings table on **favipiravir compared to baloxavir marboxil** (1 RCT: Lou 2020) [69] - https://covid-nma.com/living_data/index.php

Favipiravir compared to Baloxavir marboxil for Mild/COVID-19

Patient or population: Mild/COVID-19

Setting: Worldwide

Intervention: Favipiravir

Comparison: Baloxavir marboxil

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Baloxavir marboxil	Risk with Favipiravir				
Incidence viral negative conversion D7	600 per 1.000	402 per 1.000 (162 to 996)	RR 0.67 (0.27 to 1.66)	20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,c}	
Incidence clinical Improvement D7	100 per 1.000	200 per 1.000 (21 to 1.000)	RR 2.00 (0.21 to 18.69)	20 (1 RCT)	⊕○○○ VERY LOW ^{b,c,d}	
Incidence clinical Improvement D14-D28	600 per 1.000	498 per 1.000 (222 to 1.000)	RR 0.83 (0.37 to 1.85)	20 (1 RCT)	⊕○○○ VERY LOW ^{b,c,d}	
Incidence of WHO progression score (level 6 or above D14-D28)	100 per 1.000	33 per 1.000 (2 to 732)	RR 0.33 (0.02 to 7.32)	20 (1 RCT)	⊕○○○ VERY LOW ^{b,c,d}	
Incidence of WHO progression score (level 7 or above D14-D28)	100 per 1.000	33 per 1.000 (2 to 732)	RR 0.33 (0.02 to 7.32)	20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,c}	
All-cause mortality D7	-	-	-	20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,e}	zero events in both groups
All-cause mortality D14-D28	-	-	-	20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,e}	zero events in both groups
Adverse events - not reported	-	-	-	-	-	outcome not yet measured or reported
Serious adverse events D14-D28	600 per 1.000	402 per 1.000 (162 to 996)	RR 0.67 (0.27 to 1.66)	20 (1 RCT)	⊕⊕○○ LOW ^{d,f,g}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: **High certainty:** We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviations from intended interventions and selection of the reported results; b. Indirectness downgraded by 1 level: single study from a single institution, therefore results in this population might not be generalizable to other settings; c. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants; d. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviations from intended interventions, measurement of the outcome and selection of the reported results; e. Imprecision downgraded by 2 levels: no events in both groups and very low number of participants; f. Indirectness not downgraded: we presume that adverse event rate is not specific to a certain setting; g. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants

Results: Therapeutics

Table 3.3-3: Summary of findings table on favipiravir compared to lopinavir + ritonavir or darunavir/cobicistat + umifenovir + interferon-a (1 RCT: Lou 2020) [69] - https://covid-nma.com/living_data/index.php

Favipiravir compared to Lopinavir + Ritonavir or Darunavir/Cobicistat + Umifenovir + Interferon-a for Mild/COVID-19

Patient or population: Mild/COVID-19

Setting: Worldwide

Intervention: Favipiravir

Comparison: Lopinavir + Ritonavir or Darunavir/Cobicistat + Umifenovir + Interferon-a

Outcomes	Anticipated absolute effects* (95% CI) Risk with Lopinavir + Ritonavir or Darunavir/Cobicistat + Umifenovir + Interferon-a	Risk with Favipiravir	Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
Incidence viral negative conversion D7	500 per 1.000	400 per 1.000 (150 to 1.000)	RR 0.80 (0.30 to 2.13)	20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,c}	
Incidence clinical Improvement D7	100 per 1.000	200 per 1.000 (21 to 1.000)	RR 2.00 (0.21 to 18.69)	20 (1 RCT)	⊕○○○ VERY LOW ^{b,c,d}	
Incidence clinical Improvement D14-D28	500 per 1.000	500 per 1.000 (210 to 1.000)	RR 1.00 (0.42 to 2.40)	20 (1 RCT)	⊕○○○ VERY LOW ^{b,c,d}	
Incidence of WHO progression score (level 6 or above D14-D28)				20 (1 RCT)	⊕○○○ VERY LOW ^{b,d,e}	zero events in both groups
Incidence of WHO progression score (level 7 or above D14-D28)				20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,e}	zero events in both groups
All-cause mortality D7				20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,e}	zero events in both groups
All-cause mortality D14-D28				20 (1 RCT)	⊕○○○ VERY LOW ^{a,b,e}	zero events in both groups
Adverse events - not reported	-	-	-	-	-	outcome not yet measured or reported
Serious adverse events D14-D28	400 per 1.000	400 per 1.000 (136 to 1.000)	RR 1.00 (0.34 to 2.93)	20 (1 RCT)	⊕⊕○○ LOW ^{d,f,g}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: **High certainty:** We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviations from intended interventions and selection of the reported results; b. Indirectness downgraded by 1 level: single study from a single institution, therefore results in this population might not be generalizable to other settings; c. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants; d. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviations from intended interventions, measurement of the outcome and selection of the reported results; e. Imprecision downgraded by 2 levels: no events in both groups and very low number of participants; f. Indirectness not downgraded: we presume that adverse event rate is not specific to a certain setting; g. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants

Results: Therapeutics

Table 3.3-4: Summary of findings table on **favipiravir compared to standard care** (6 RCTs: Lou 2020, Ivashchenko 2020, Dabbous 2020, Balykova 2020, Ruzhentsova 2020, Udwardia 2020) - https://covid-nma.com/living_data/index.php

Favipiravir compared to Standard care for Mild/Moderate/Unclear COVID-19

Patient or population: Mild/Moderate/Unclear COVID-19

Setting: Worldwide

Intervention: Favipiravir

Comparison: Standard care

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard care	Risk with Favipiravir				
Viral negative conversion D3	455 per 1,000	555 per 1,000 (450 to 602)	RR 1.22 (0.99 to 1.50)	318 (3 RCTs) ^b	⊕⊕○○ LOW ^{c,d}	
Viral negative conversion D7	688 per 1,000	750 per 1,000 (653 to 867)	RR 1.09 (0.95 to 1.26)	677 (6 RCTs) ^e	⊕○○○ VERY LOW ^{d,f,g}	
Clinical improvement D7	221 per 1,000	349 per 1,000 (254 to 477)	RR 1.58 (1.15 to 2.16)	379 (3 RCTs) ^g	⊕⊕○○ LOW ^{h,i}	
Clinical improvement D14-28	885 per 1,000	895 per 1,000 (868 to 931)	RR 1.00 (0.97 to 1.04)	379 (4 RCTs) ^j	⊕⊕○○ LOW ^{h,i}	
WHO progression score (level 6 or above) D7	100 per 1,000	300 per 1,000 (37 to 1,000)	RR 3.00 (0.37 to 24.17)	20 (1 RCT) ^k	⊕○○○ VERY LOW ^{l,m,n}	
WHO progression score (level 6 or above) D14-28	0 per 1,000 (0 to 0)	0 per 1,000 (0 to 0)	not estimable	220 (2 RCTs) ^o	⊕○○○ VERY LOW ^{l,p}	zero events in both groups
WHO progression score (level 7 or above) D7	0 per 1,000 (0 to 0)	0 per 1,000 (0 to 0)	not estimable	20 (1 RCT) ^k	⊕○○○ VERY LOW ^{l,p,q}	zero events in both groups
WHO progression score (level 7 or above) D14-28	0 per 1,000 (0 to 0)	0 per 1,000 (0 to 0)	not estimable	220 (2 RCTs) ^o	⊕○○○ VERY LOW ^{l,q}	zero events in both groups
All-cause mortality D7	6 per 1,000	2 per 1,000 (0 to 50)	RR 0.33 (0.01 to 7.99)	320 (3 RCTs) ^r	⊕○○○ VERY LOW ^{l,q}	zero events in the intervention group
All-cause mortality D14-28	6 per 1,000	2 per 1,000 (0 to 44)	RR 0.32 (0.01 to 7.82)	360 (3 RCTs) ^o	⊕○○○ VERY LOW ^{l,q}	zero events in the intervention group
Adverse events	289 per 1,000	455 per 1,000 (259 to 800)	RR 1.53 (0.87 to 2.69)	559 (3 RCTs) ^o	⊕○○○ VERY LOW ^{l,s,t}	
Serious adverse events	22 per 1,000	26 per 1,000 (10 to 64)	RR 1.20 (0.48 to 2.99)	519 (4 RCTs) ^o	⊕○○○ VERY LOW ^{l,u}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
 CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence
 High certainty: We are very confident that the true effect lies close to that of the estimate of the effect.
 Moderate certainty: We are moderately confident in the effect estimate. The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.
 Low certainty: Our confidence in the effect estimate is limited. The true effect may be substantially different from the estimate of the effect.
 Very low certainty: We have very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of effect.

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect. **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.**Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect.**Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Results: Therapeutics

Explanations: a. Last update: December 4, 2020; b. Lou Y, 2020; Ruzhentsova T, 2020; Udwardia Z, 2020; c. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviation from intended intervention and selection of reported results; d. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for no effect; e. Balykova L, 2020; Dabbous HM, 2020; Ivashchenko AA, 2020; Lou Y, 2020; Ruzhentsova T, 2020; Udwardia Z, 2020; f. Inconsistency downgraded by 1 level: $I^2=50.4\%$; g. Balykova L, 2020; Lou Y, 2020; Ruzhentsova T, 2020; h. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviation from intended intervention, outcome measurement and selection of reported results; i. Imprecision downgraded by 1 level: due to low number of events and/or participants; j. Ivashchenko AA, 2020; Lou Y, 2020; Ruzhentsova T, 2020; Udwardia Z, 2020; k. Lou Y, 202; l. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviations from intended intervention and outcome measurement; m. Indirectness downgraded by 1 level: single study from a single institution, therefore results in this population might not be generalizable to other settings; n. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants; o. Balykova L, 2020; Lou Y, 2020; p. Imprecision downgraded by 2 levels: no events in both groups and low number of participants; q. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization and deviation from intended intervention; r. Balykova L, 2020; Dabbous HM, 2020; Lou Y, 2020; s. Inconsistency downgraded by 1 level: $I^2=78.9\%$; t. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for no effect and the possibility for harm

3.4 Darunavir

About the drug under consideration

Darunavir is an antiviral agent from the group of human immunodeficiency virus (HIV) protease inhibitors for the treatment of HIV-1 infections. Darunavir is combined with a pharmacokinetic booster such as ritonavir or cobicistat [80].

Darunavir (Prezista®) has not been approved by the European Medicines Agency (EMA) or the American Food and Drug Administration (FDA) for COVID-19.

The US COVID-19 Treatment Guidelines Panel recommends **against** using the **Lopinavir/ritonavir (AI) or other HIV protease inhibitors (AIII)**, except in a clinical trial, because of unfavorable pharmacodynamics and because clinical trials have not demonstrated a clinical benefit in patients with COVID-19 [61].

Withdrawn, suspended or terminated studies

The search in two clinical trial registers (ClinicalTrials.gov and EudraCT) yielded no suspended, withdrawn or terminated RCTs in COVID-19.

Results of publications

Chen J et al. 2020 [81] published results from single-center, randomized, open-label trial (**NCT04252274**) which aimed to evaluate the antiviral activity and safety of darunavir/cobicistat (DRV/c) in treating mild COVID-19 patients. Participants were randomized to receive DRV/c for 5 days on the top of interferon alpha 2b inhaling or interferon alpha 2b inhaling alone. DRV/c did not increase the proportion of negative conversion vs standard of care alone: the proportion of negative PCR results at day 7 was 46.7% (7/15) and 60.0% (9/15) in the DRV/c and control groups ($p = 0.72$), respectively. The viral clearance rate at day 3 was 20% (3/15) in both study groups, while the number increased to 26.7% (4/15) in the DRV/c group and remained 20% (3/15) in the control group at day 5. Fourteen days after randomization, 1 participant in the DRV/c group progressed to critical illness and discontinued DRV/c, while all the patients in the control group were stable ($p=1.0$). The frequencies of adverse events in the two groups were comparable. The findings are presented in Table 3.4-1.

antivirales Medikament

**als HIV Medikament
zugelassen
EMA 2007**

**Empfehlungen des US
COVID-19 Treatment
Guidelines Panel GEGEN
jegliche HIV Protease
Inhibitoren**

**keine weiteren Studien in
ClinicalTrials.gov and
EudraCT als
abgeschlossen oder
beendet registriert**

Publikation zu RCT

**bei milder Covid-19
Erkrankung
DRV+IFN vs. IFN
kein Unterschied**

Results: Therapeutics

Table 3.4-1: Summary of findings table on **darunavir/cobicistat compared to standard care** (1 RCT: Chen J) - https://covid-nma.com/living_data/index.php [81]

Darunavir/cobistat compared to Standard Care for Moderate COVID-19

Patient or population: Moderate COVID-19

Setting: Worldwide

Intervention: Darunavir/cobistat

Comparison: Standard Care

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	№ of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard Care	Risk with Darunavir/cobicistat				
Incidence of viral negative conversion D7	600 per 1.000	468 per 1.000 (234 to 924)	RR 0.78 (0.39 to 1.54)	30 (1 RCT)	⊕○○○ VERY LOW a,b,c	
Clinical improvement - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical recovery - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO progression score (level 6 or above) - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO progression score (level 7 or above D7)	0 per 1.000	0 per 1.000 (0 to 0)	RR 3.00 (0.13 to 68.26)	30 (1 RCT)	⊕○○○ VERY LOW a,b,d	zero events in control group
All-cause mortality D14-D28				30 (1 RCT)	⊕○○○ VERY LOW a,b,e	zero events in both groups
Adverse events - not reported	-	-	-	-	-	outcome not yet measured or reported
Serious adverse events D14-D28				30 (1 RCT)	⊕○○○ VERY LOW e,f,g	zero events in both groups
*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval; RR: Risk ratio						

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Risk of bias downgraded by 1 level: some concerns or high risk due to concerns during the randomization process, deviations from intended interventions and selection of the reported results; b. Indirectness downgraded by 1 level: single study from a single institution, therefore results in this population might not be generalizable to other settings; c. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants; d. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants e. Imprecision downgraded by 2 levels: no events in both groups and very low number of participants; f. Risk of bias downgraded by 2 levels: some concerns or high risk due to concerns during the randomization process, deviation from intended intervention, missing data and selection of reported results; g. We presume that the adverse event rates, and the corresponding relative risks, is similar across diverse settings, therefore not downgraded for indirectness

3.5 Chloroquine (Resochin®) and

3.6 Hydroxychloroquine (Plaquenil®)

Due to the lack of effectiveness of chloroquine and hydroxychloroquine in treating COVID-19 patients; in the light of serious adverse effects as well as the decisions to stop enrolling participants to the hydroxychloroquine arm of the RECOVERY and SOLIDARITY trials, the reporting related to these two pharmaceuticals was stopped also.

Last reporting V4/ July:

https://eprints.aihta.at/1234/10/Policy_Brief_002_Update_07.2020.pdf

wegen erwiesenem Mangel an Wirksamkeit wurde Beobachtung beendet

3.7 Camostat Mesilate (Foipan®)

About the drug under consideration

Camostat Mesilate (Foipan®) is classified as a so-called serine protease inhibitor, blocking several pancreatic and plasmatic enzymes like trypsin, thrombin and plasmin [82]. Studies showed effects on the cell-entry mechanism of coronaviruses (e.g. SARS-CoV and SARS-CoV-2) in in-vitro human cells [83, 84] as well as in pathogenic mice-models [85] by inhibiting the enzyme Transmembrane protease, serine 2 (TMPRSS2).

Camostat Mesilate (Foipan®) ist not approved for any anti-viral use (FDA, EMA).

It is one of the drugs for which the German Federal Ministry of Health initiated centralized procurement in April 2020 for the treatment of infected and seriously ill COVID-19 patients in Germany (<https://www.abda.de>). Up to August 1, 2020, 35 to 60 Covid-19 patients have been treated with the centrally procured medicinal product Foipan (Camostat) as part of an individual medical treatment. There was no obligation for the treating physicians to collect data in a registry [86].

**Protease-Inhibitor bei Entzündung der Bauchspeicheldrüse
Zulassung: Japan, Südkorea**

**nicht EMA, FDA
FDA: Orphan Drug Designation seit 2011
vom dt. BMG für schwere Erkrankungen zentral eingekauft**

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated studies were found in ClinicalTrials.gov and EUdraCT registers.

in ClinicalTrials.gov and EUdraCT keine klinischen Studien registriert

Results of publications

Until now no scientific publication on a RCT of Camostat Mesilate (Foipan®) in Covid-19 patients could be identified.

3.8 APN01/ Recombinant Human Angiotensin-converting Enzyme 2 (rhACE2)

Drug under consideration

APN01 is a recombinant human Angiotensin Converting Enzyme 2 (rhACE2) developed by Apeiron Biologics under Phase 2 clinical development in ALI (Acute Lung Injury) and PAH (Pulmonal arterial hypertension) [87], [88], [89].

The therapy with APN01 is currently not approved by the European Medicine Agency (EMA) and Food and Drug Administration (FDA) for COVID-19.

Withdrawn, suspended or terminated studies

One RCT number NCT04287686 is visible as withdrawn (without CDE Approval).

Results of publications

No relevant finished publications or finished trials assessing the efficacy and safety could be identified. First results, related to a phase 2/3 study of rhACE2 in 200 hospitalised patients with COVID-19, with primary composite outcome – All-cause mortality or invasive mechanical ventilation can be expected on the 10th of November 2020 (NCT04335136) [90].

aus SARS-Forschung hervorgegangen

**keine Zulassung
1 Studie (Phase 2 RCT), vor Rekrutierung**

in ClinicalTrials.gov and EUdraCT keine abgeschlossene, aber eine zurückgezogene Studie registriert

keine Publikationen zu klinischen Studien

3.9 Tocilizumab (Roactemra®)

Drug under consideration

Tocilizumab (*RoActemra*) is a human monoclonal antibody that specifically binds to soluble and membrane-bound interleukin (IL)-6 receptors (IL-6R α), and inhibits IL-6-mediated signalling [91].

Tocilizumab is being investigated as a possible treatment for patients with moderate to severe or critical COVID-19. The therapy is currently not approved by the European Medicine Agency (EMA) and Food and Drug Administration (FDA) for COVID-19.

The US COVID-19 Treatment Guidelines Panel **recommend against** anti-IL-6 receptor monoclonal antibodies (e.g., sarilumab, tocilizumab) or anti-IL-6 monoclonal antibody (siltuximab) (BI) for the treatment of COVID-19 [61], except in a clinical trial.

Withdrawn, suspended or terminated studies

One withdrawn RCT (NCT04361552, in US, abandoned due to drug billing issues) and four terminated RCTs were found in ClinicalTrials.gov and EudraCT registers: NCT04346355, in Italy, based on interim analysis for futility and given an enrolment rate almost nil; RCT on 129 patients in Brazil compared tocilizumab vs best supportive care NCT04403685 (TOCIBRAS) due to safety issue; RCT NCT04322773, TOCOVID trial, due to changed clinical conditions and too few patients available; RCT NCT04335071 (CORON-ACT) in Switzerland because dexamethasone was included in the

Interleukin-6-Rezeptor für rheumatoide Arthritis zugelassen (EMA)

Empfehlungen des US COVID-19 Treatment Guidelines Panel: insuffiziente Datenlage, nur in klinischen Studien

**1 beendeter RCT,
1 zurückgezogener (admin Gründe),
1 abgebrochener (Mangel an Rekrutierung)**

standard care and planned number of patients was not possible to recruit in the planned study period).

Results of publications

Rosas et al. 2020 [92] reported results from the phase 3, RCT - **COVACTA** (NCT04320615, EUdraCT 2020-001154-22) as preprint: 452 patients with severe COVID-19 pneumonia were randomized; the modified-intention-to-treat population included 294 tocilizumab-treated and 144 placebo-treated patients. Clinical status at day 28 was not statistically significantly improved for tocilizumab versus placebo ($p=0.36$). Median (95% CI) ordinal scale values at day 28: 1.0 (1.0 to 1.0) for tocilizumab and 2.0 (1.0 to 4.0) for placebo (odds ratio, 1.19 [0.81 to 1.76]). There was no difference in mortality at day 28 between tocilizumab (19.7%) and placebo (19.4%) (difference, 0.3% [95% CI, -7.6 to 8.2]; nominal $p=0.94$). Median time to hospital discharge was 8 days shorter with tocilizumab than placebo (20.0 and 28.0, respectively; nominal $p=0.037$; hazard ratio 1.35 [95% CI 1.02 to 1.79]). Median duration of ICU stay was 5.8 days shorter with tocilizumab than placebo (9.8 and 15.5, respectively; nominal $p=0.045$). In the safety population, serious adverse events occurred in 34.9% of 295 patients in the tocilizumab arm and 38.5% of 143 in the placebo arm.

Wang et al. 2020 [93] reported, as preprint, results from a small randomized, controlled, open-label, multicenter trial at 6 hospitals in Anhui and Hubei (**ChiCTR2000029765**). 65 moderate to severe patients were enrolled and randomly assigned to a treatment group (33 to tocilizumab and 32 to the controls). The cure rate in tocilizumab group was higher than that in the controls but not significant (94.12% vs 87.10%, $p=0.4133$). Adverse events were recorded in 20 (58.82%) of 34 tocilizumab recipients versus 4 (12.90%) of 31 in the controls. No serious adverse events were reported in tocilizumab group.

Salama et al. 2020 [94], reported as preprint, results from the phase III **EMPACTA** study (NCT04372186) (389 patients in the United States, South Africa, Kenya, Brazil, Mexico and Peru), showing that patients with COVID-19 associated pneumonia who received tocilizumab plus standard of care were 44% less likely to progress to mechanical ventilation or death compared to patients who received placebo plus standard of care (log-rank p -value = 0.0348; HR [95% CI] = 0.56 [0.32, 0.97]). The cumulative proportion of patients who progressed to mechanical ventilation or death by day 28 was 12.2% in tocilizumab arm versus 19.3% in the placebo arm. Key secondary outcomes (difference in time to hospital discharge or “ready for discharge” to day 28; difference in time to improvement in ordinal clinical status to day 28; time to clinical failure to day 28 and mortality by day 28) were not statistically significant different between groups. At day 28, incidence of infections was 10% and 11% in the tocilizumab and placebo arms, respectively, and the incidence of serious infections was 5.0% and 6.3% in tocilizumab and placebo arms, respectively. The most common adverse events in patients who received tocilizumab were constipation (5.6%), anxiety (5.2%), and headache (3.2%).

COVACTA
4RCT, 52 Pts
schwere Erkrankung

kein Unterschied bei
Mortalität, aber kürzer
Zeit zur Erholung

Wang (China)
65 Pts
schwere Erkrankung

EMPACTA
389 Pts
RCT (US, SA, Kenya,
Brasilien, Mexiko, Peru)
schwere Erkrankung

Vorteil bei Verhinderung
im Fortschreiten der
Erkrankung

bei weiteren Endpunkten:
kein Unterschied

Hermine et al. 2020 [95] published the results from multicentre **CORIMUNO-TOCI-1 RCT (NCT04331808)**, which included 131 moderate to severe COVID-19 patients (63 treated with tocilizumab, others in usual care group) in France, with follow-up through 28 days. In the TCZ group, 12 patients had a WHO-CPS score greater than 5 at day 4 vs 19 in the UC group (median posterior absolute risk difference [ARD] -9.0%; 90% credible interval [CrI], -21.0 to 3.1), with a posterior probability of negative ARD of 89.0% not achieving the 95% predefined efficacy threshold. At day 14, 12% (95% CI -28% to 4%) fewer patients needed noninvasive ventilation (NIV) or mechanical ventilation (MV) or died in the TCZ group than in the UC group (24% vs 36%, median posterior hazard ratio [HR] 0.58; 90% CrI, 0.33-1.00), with a posterior probability of HR less than 1 of 95.0%, achieving the predefined efficacy threshold. The HR for MV or death was 0.58 (90% CrI, 0.30 to 1.09). At day 28, 7 patients had died in the TCZ group and 8 in the UC group (adjusted HR, 0.92; 95% CI 0.33-2.53). Serious adverse events occurred in 20 (32%) patients in the TCZ group and 29 (43%) in the UC group (p=0.21).

CORIMUNO-TOCI-1
131 Pts.
moderate bis
schwere Erkrankung

Vorteil bei Bedarf nach
Beatmung
kein Unterschied bei
Mortalität

Salvarani et al. 2020 [96] published results from multicentre RCT (**RCT-TCZ-COVID-19 (NCT04346355)**) conducted on 126 severe COVID-19 patients in Italy (60 received tocilizumab). Seventeen patients of 60 (28.3%) in the tocilizumab arm and 17 of 63 (27.0%) in the standard care group showed clinical worsening within 14 days since randomization (rate ratio, 1.05; 95% CI, 0.59-1.86). Two patients in the experimental group and 1 in the control group died before 30 days from randomization, and 6 and 5 patients were intubated in the 2 groups, respectively. The trial was prematurely interrupted after an interim analysis for futility.

RCT-TCZ-COVID-19
126 Pts
schwere Erkrankung

kein Unterscheid,
frühzeitiger
Studienabbruch

Stone et al. 2020 [97] published results from multicentre RCT (**NCT04356937**) conducted on 243 moderate to severe COVID-19 patients in US (161 received tocilizumab). The hazard ratio for intubation or death in the tocilizumab group vs placebo group was 0.83 (95% confidence interval [CI], 0.38 to 1.81; p=0.64), and the hazard ratio for disease worsening was 1.11 (95% CI, 0.59 to 2.10; p=0.73). At 14 days, 18.0% of the patients in the tocilizumab group and 14.9% of the patients in the placebo group had worsening of disease. The median time to discontinuation of supplemental oxygen was 5.0 days (95% CI, 3.8 to 7.6) in the tocilizumab group vs 4.9 days (95% CI, 3.8 to 7.8) in the placebo group (p=0.69). At 14 days, 24.6% of the patients in the tocilizumab group and 21.2% of the patients in the placebo group were still receiving supplemental oxygen. Patients who received tocilizumab had fewer serious infections than patients who received placebo.

RCT 243
moderate bis
schwere Erkrankung

keine oder kaum
Unterschiede in einigen
Endpunkten

Tocilizumab continues to be evaluated in the **RECOVERY trial**. Because over 850 patients randomised to tocilizumab versus standard of care (almost twice the size of the COVACTA trial) will provide critical data to confirm or refute the COVACTA results [98].

Tocilizumab auch in
RECOVERY
850 Pts

Meta-analysis with Summary of findings table on tocilizumab compared to standard of care (related to **6 RCTs**) is presented in Table 3.9-1. In all outcomes presented, including All-cause mortality, there was no statistically significant difference in risk ratio between tocilizumab and standard care/placebo group.

Results: Therapeutics

Table 3.9-1: Summary of findings table on **tocilizumab compared standard care/placebo** (6 RCTs: Rosas, Wang, Hermine, Salvarani, Stone, Salama)

Tocilizumab compared to Standard care/Placebo for Mild/Moderate/Severe/Critical COVID-19

Patient or population: Mild/Moderate/Severe/Critical COVID-19

Setting: Worldwide

Intervention: Tocilizumab

Comparison: Standard care/Placebo

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard care/Placebo	Risk with Tocilizumab				
Incidence of viral negative conversion D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement D7 - not measured	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement D14-28	897 per 1,000	906 per 1,000 (843 to 977)	RR 1.01 (0.94 to 1.09)	366 (2 RCTs) ^b	⊕⊕⊕⊕ LOW ^{g,h}	
Incidence of WHO progression score (level 6 or above) D7	537 per 1,000	493 per 1,000 (419 to 564)	RR 0.99 (0.78 to 1.05)	582 (2 RCTs) ^e	⊕⊕⊕⊕ MODERATE ^d	
Incidence of WHO progression score (level 6 or above) D14-D28	381 per 1,000	308 per 1,000 (247 to 392)	RR 0.81 (0.65 to 1.03)	582 (2 RCTs) ^e	⊕⊕⊕⊕ MODERATE ^d	
Incidence of WHO progression score (level 7 or above) D7	389 per 1,000	347 per 1,000 (247 to 467)	RR 0.87 (0.62 to 1.22)	582 (2 RCTs) ^e	⊕⊕⊕⊕ MODERATE ^d	
Incidence of WHO progression score (level 7 or above) D14-D28	312 per 1,000	278 per 1,000 (209 to 371)	RR 0.89 (0.67 to 1.19)	582 (2 RCTs) ^e	⊕⊕⊕⊕ MODERATE ^d	
All-cause mortality D7	73 per 1,000	83 per 1,000 (42 to 164)	RR 1.14 (0.58 to 2.25)	452 (1 RCT) ^f	⊕⊕⊕⊕ LOW ^g	
All-cause mortality D14-D28	104 per 1,000	114 per 1,000 (83 to 156)	RR 1.1 (0.8 to 1.5)	1327 (5 RCTs) ^h	⊕⊕⊕⊕ LOW ^{g,i}	
Adverse events	124 per 1,000	241 per 1,000 (71 to 825)	RR 1.94 (0.57 to 6.64)	431 (3 RCTs) ^j	⊕⊕⊕⊕ VERY LOW ^{g,k,l}	
Serious adverse events	442 per 1,000	384 per 1,000 (322 to 464)	RR 0.87 (0.73 to 1.05)	1392 (6 RCTs) ^m	⊕⊕⊕⊕ LOW ^{g,n}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Last update: November 6, 2020; b. Stone JH, 2020; Salvarani C, 2020; c. Risk of bias downgraded by 1 level: some concerns or high risk regarding deviations from intended interventions, randomization, and outcome measurement; d. Imprecision downgraded by 1 level: due to low number of events and participants; e. Hermine O, 2020; Rosas I, 2020; f. Rosas I, 2020; g. Imprecision downgraded by 2 levels: due to low number of events and a wide confidence interval consistent with the possibility for benefit and the possibility for harm; h. Stone JH, 2020; Hermine O, 2020; Rosas I, 2020; Salama C, 2020; Salvarani C, 2020; i. Risk of bias downgraded by 1 level: some concerns or high risk regarding deviations from intended interventions and randomization; j. Stone JH, 2020; Wang D, 2020; Salvarani C, 2020; k. Inconsistency downgraded by 1 level: unexplained statistical heterogeneity (I-sq = 83.5%); l. We presume that the adverse event rates, and the corresponding relative risks, are similar across diverse setting

3.10 Sarilumab (Kevzara®)

Drug under consideration

Sarilumab (*Kevzara*) is a human monoclonal antibody that specifically binds to soluble and membrane-bound interleukin (IL)-6 receptors (IL-6R α), and inhibits IL-6-mediated signalling [99]. It is being investigated as a possible treatment for patients with moderate to severe or critical COVID-19. The therapy is currently not approved by the European Medicine Agency (EMA) and Food and Drug Administration (FDA) for COVID-19.

The US COVID-19 Treatment Guidelines Panel **recommend against** anti-IL-6 receptor monoclonal antibodies (e.g., sarilumab, tocilizumab) or anti-IL-6 monoclonal antibody (siltuximab) (BI) for the treatment of COVID-19 [61], except in a clinical trial.

Withdrawn, suspended or terminated studies

One RCT found as suspended, NCT04341870 - CORIMUNO-VIRO Trial (DSMB recommendation (futility)). One RCT found as terminated, NCT04322773 (TOCIVID) in Denmark, due to changed clinical conditions and too few patients available).

Results of publications

On July 03, 2020 in press release related to sarilumab RCT conducted in US, <https://www.clinicaltrialsarena.com/news/kevzara-us-covid19-trial-data/>, Sanofi and Regeneron Pharmaceuticals have reported that this phase III clinical trial of sarilumab, compared 400mg dose of the drug plus best supportive care to best supportive care alone, failed to meet its primary and key secondary endpoints in 194 critically ill Covid-19 patients who required mechanical ventilation in the US. In the primary analysis arm, adverse events were reported in 80% of patients treated with sarilumab and 77% of those on placebo. Serious adverse events in at least 3% of patients, more frequent among sarilumab patients, were multi-organ dysfunction syndrome and hypotension. Based on the data, the companies have halted this US-based trial, including a second cohort of patients who were on a higher 800mg dose of the drug. The trial being conducted outside of the US is continuing, in hospitalised patients with severe and critical Covid-19 using a different dosing regimen.

Interleukin-6-Rezeptor für rheumatoide Arthritis zugelassen (EMA)

Covid-10: bei erhöhten IL-6-Spiegeln

Empfehlung des US COVID-19 Treatment Guidelines Panel: insuffiziente Datenlage, nur in klinischen Studien

Juli 2020: Pressemeldung zu RCT mit 194 Pts

kein Unterschied mehr SAE in Sarilumab Gruppe

3.11 Interferon beta 1a (SNG001) (Rebif®, Avonex®) and Interferon beta 1b (Betaferon®, Extavia®)

About the drug under consideration

Interferon beta-1a (INFb) is a cytokine in the interferon family used to treat relapsing multiple sclerosis (MS). Finding of studies in patients with MERS-CoV have led to exploration of treatment with INFb in COVID-19 [100].

INFb Präparate bei Multipler Sklerose zugelassen (EMA)

Two pharmaceuticals which the active substance Interferon beta-1a are commercially available: Rebif® and Avonex®. They are used to slow the progression of disability and reduce the number of relapses in MS. Rebif is approved by the European Medicines Agency (EMA) since 1998 and by the American Food and Drug Administration (FDA) since 2002. Avonex is approved by EMA since 1997 and by the FDA since 1996. Both drugs are approved for the treatment of relapsing forms of multiple sclerosis (MS), in cases of clinically isolated syndromes, as well as relapsing remitting disease, and active secondary progressive disease in adults.

**Interferon beta-1a: Rebif®
Avonex®
seit 1997/1998 zugelassen**

nicht für Covid-19

Two pharmaceuticals, with the active substance Interferon beta-1b, are commercially available in EU: Betaferon® and Extavia® to treat adults with multiple sclerosis (MS) [101, 102]. Betaferon® is approved by the European Medicines Agency (EMA) since 1995. Extavia® is approved by EMA since 2008. Interferon beta-1a and beta-1b are not approved for COVID-19 patients treatment.

**Interferon beta-1b:
Betaferon® and Extavia®
seit 1995/2008 zugelassen
nicht für Covid-19**

The US COVID-19 Treatment Guidelines Panel [61] **recommends against** use of the **interferons (alfa or beta)** for the treatment of **severely or critically ill** patients with COVID-19, except in the context of a clinical trial (AIII).

**Empfehlung des US
COVID-19 Treatment
Guidelines Panel: nur in
klinischen Studien**

There are **insufficient data** for the Panel to recommend **either for or against** the use of the **Interferon-beta** for the treatment of early (i.e., <7 days from symptom onset) **mild and moderate** COVID-19.

Withdrawn, suspended or terminated studies

One RCT was found as suspended, NCT04469491 (COV-NI), on interferon beta 1b by nebulization in France (in anticipation for Data and Safety Monitoring Board).

Results of publications

The results from the first randomised controlled trial on triple combination of interferon beta-1b, lopinavir–ritonavir and ribavirin, in comparison with lopinavir–ritonavir (NCT04276688) are presented in Section 3.14 of this report [103].

**Kombinationstherapie:
Ergebnisse in 3.14**

Results from **Huang et al. 2020 (ChiCTR2000029387)** [104] related to Ribavirin Plus Interferon-Alpha, Lopinavir/Ritonavir Plus Interferon-Alpha, and Ribavirin Plus Lopinavir/Ritonavir Plus Interferon-Alpha in Patients With Mild to Moderate COVID-19 were presented in Section 3.14 of this report.

**August 2020:
2 RCTs publiziert
1 RCT zu Kombinations-
therapie in 3.14**

Esquivel-Moynelo et al. 2020 [105] presented the results from a RCT for efficacy and safety evaluation of subcutaneous IFN- α 2b and IFN γ administration in 79 patients positive to SARS-CoV-2. Patients were randomly assigned in a 1:1 ratio to receive either, subcutaneous treatment with a combination of 3.0 MIU IFN- α 2b and 0.5 MIU IFN- γ , twice a week for two weeks, or thrice a week intramuscular injection of 3.0 MIU IFN- α 2b. Additionally, all patients received lopinavir-ritonavir 200/50 mg every 12 h and chloroquine 250 mg every 12 h (standard of care). None of the patients developed severe COVID-19 during the study or the epidemiological follow-up for 21 more days.

**1 RCT
79 Pts.
Kombinationstherapie IFN
(unterscheidliche
Dosierungen) + Kaletra**

**79 symptomatische/
asymptomatische Pts.**

Monk et al. 2020 published results from randomised, double-blind, placebo-controlled, phase 2 pilot trial at nine UK sites (NCT04385095) [106]. 101 COVID-19 hospitalized adult patients were randomly assigned (1:1) to receive **inhaled nebulised interferon beta-1a** (SNG001) (6 MIU) or placebo by

**1 RCT
101 Pts
inhaltiertes INF**

inhalation via a mouthpiece daily for 14 days. 66 (67%) patients required oxygen supplementation at baseline: 29 in the placebo group and 37 in the SNG001 group. Patients receiving SNG001 had greater odds of improvement on the OSCI scale (odds ratio 2.32 [95% CI 1.07–5.04]; $p=0.033$) on day 15 or 16 and were more likely than those receiving placebo to recover to an OSCI score of 1 (no limitation of activities) during treatment (hazard ratio 2.19 [95% CI 1.03–4.69]; $p=0.043$). No significant difference was found between treatment groups in the odds of hospital discharge by day 28: 39 (81%) of 48 patients had been discharged in the nebulised interferon beta-1a group compared with 36 (75%) of 48 in the placebo group (OR 1.84 [95% CI 0.64–5.29]; $p=0.26$). There was no significant difference between treatment groups in the odds of intubation or the time to intubation or death. SNG001 was well tolerated: the most frequently reported treatment-emergent adverse event was headache (seven [15%] patients in the SNG001 group and five [10%] in the placebo group). There were three deaths in the placebo group and none in the SNG001 group.

Davoudi-Monfared et al. 2020 published results related to the RCT on Interferon beta-1a treatment ($n=46$) vs the **standard of care** ($n=46$), in 92 patients with severe COVID-19 in Iran (**IRCT20100228003449N28**) [107]. Finally 81 patients (42 in the IFN and 39 in the control group) completed the study. Time to the clinical response was not significantly different between the IFN and the control groups (9.7 +/- 5.8 vs. 8.3 +/- 4.9 days respectively, $P=0.95$). On day 14, 66.7% vs. 43.6% of patients in the IFN group and the control group were discharged, respectively (OR= 2.5; 95% CI: 1.05- 6.37). The 28-day overall mortality was significantly lower in the IFN then the control group (19% vs. 43.6% respectively, $p= 0.015$). Early administration significantly reduced mortality (OR=13.5; 95% CI: 1.5-118).

Rahmani et al. 2020 [108] published the results of RCT evaluated efficacy and safety of interferon (IFN) β -1b in the treatment of 80 patients with severe COVID-19 (**IRCT20100228003449N27**). Patients in the IFN group received IFN β -1b (250 mcg subcutaneously every other day for two consecutive weeks) along with the national protocol medications while in the control group, patients received only the national protocol medications (lopinavir/ritonavir or atazanavir/ritonavir plus hydroxychloroquine for 7–10 days). 33 patients in each group completed the study. Time to clinical improvement in the IFN group was significantly shorter than the control group ([9(6–10) vs. 11(9–15) days respectively, $p = 0.002$, HR = 2.30; 95% CI: 1.33–3.39]). At day 14, the percentage of discharged patients was 78.79% and 54.55% in the IFN and control groups respectively (OR = 3.09; 95% CI: 1.05–9.11, $p = 0.03$). ICU admission rate in the control group was significantly higher than the IFN group (66.66% vs. 42.42%, $p = 0.04$). The duration of hospitalization and ICU stay were not significantly different between the groups. All-cause 28-day mortality was 6.06% and 18.18% in the IFN and control groups respectively ($p = 0.12$).

In **SOLIDARITY (INF)** RCT (**ISRCTN83971151**) results on comparisons of subcutaneous interferon beta-1a vs standard care in patients with mild to critical COVID-19 admitted to 405 centers in 30 countries were published as preprint [60, 66]. In 11,266 adults were randomized, with 2750 allocated remdesivir, 954 hydroxychloroquine, 1411 lopinavir, 651 interferon plus lopinavir, 1412 only interferon, and 4088 no study drug. Death rate ratio for interferon was not statistically significant different in comparison with control group: RR=1.16 (0.96-1.39, $p=0.11$; 243/2050 vs 216/2050) (or 1.12, 0.83-1.51, without lopinavir co-administration). The same is true for outcomes Initiation of ventilation or Hospitalisation duration.

Vorteil bei klinischen Verbesserungen, nicht aber bei Dauer des Spitalsaufenthalts

**RCT (Iran)
92 Pts**

Reduktion der 28-Tages Mortalität insb. bei früher Verabreichung von IFN

**RCT (Iran)
80 Pts
Zeit zur klinischen Verbesserung signifikant kürzer mit IFN, weniger ICU Einweisungen**

nicht aber Dauer der Hospitalisierung und in ICU

**SOLIDARITY
651 Pts INF + lopinavir,
1.412 Pts. nur INF**

keine Unterschiede bei den Endpunkten

Summary of Findings table related to **meta-analysis** on results of **3 RCTs** (Davoudi-Monfared, Rahmani, SOLIDARITY-INF), on comparisons of interferon beta-1a vs standard of care in patients with moderate/severe/critical COVID-19 patients, is presented in Table 3.11-1. In summary, according to the very low certainty of evidence, WHO progression score level 6 or above D14-D28; WHO progression score level 7 or above D14-D28; All-cause mortality D7 were all statistically significant better in favour of interferon beta-1a, but not outcome All-cause mortality D14-28: RR 0.68 (95%CI 0.32 to 1.45).

**sehr niedrige Evidenz:
bei Mortalität signifikante
Vorteile**

Results: Therapeutics

Table 3.11-1: Summary of findings table on **Interferon β-1a compared to Standard Care for Moderate/Severe/Critical COVID-19** (3 RCTs: Davoudi-Monfared, Rahmani, SOLIDARITY-INF) – https://covid-nma.com/living_data/index.php

Interferon β compared to Standard Care for Moderate/Severe/Critical COVID-19

Patient or population: Moderate/Severe/Critical COVID-19

Setting: Worldwide

Intervention: Interferon β

Comparison: Standard Care

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (events)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard Care	Risk with Interferon β				
Viral negative conversion - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO progression score level 6 or above D7	293 per 1,000	149 per 1,000 (59 to 375)	RR 0.51 (0.30 to 1.38)	165 (2 RCTs) ^b	⊖○○○ VERY LOW ^{1aA}	
WHO progression score level 6 or above D14-D28	268 per 1,000	123 per 1,000 (54 to 241)	RR 0.46 (0.24 to 0.90)	165 (2 RCTs) ^b	⊖○○○ VERY LOW ^{1aF}	
WHO progression score level 7 or above D7	256 per 1,000	149 per 1,000 (79 to 277)	RR 0.58 (0.31 to 1.08)	165 (2 RCTs) ^b	⊖○○○ VERY LOW ^{1aA}	
WHO progression score level 7 or above D14-D28	268 per 1,000	123 per 1,000 (54 to 241)	RR 0.46 (0.24 to 0.90)	165 (2 RCTs) ^b	⊖○○○ VERY LOW ^{1aF}	
All-cause mortality D7	134 per 1,000	15 per 1,000 (1 to 122)	RR 0.11 (0.01 to 0.91)	165 (2 RCTs) ^b	⊖○○○ VERY LOW ^{1aF}	
All-cause mortality D14-D28	112 per 1,000	76 per 1,000 (36 to 163)	RR 0.68 (0.32 to 1.45)	4265 (3 RCTs) ^c	⊖○○○ VERY LOW ^{1aA}	
Adverse events - not reported	-	-	-	-	-	outcome not yet measured or reported
Serious adverse events - not reported	-	-	-	-	-	outcome not yet measured or reported

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
CI, Confidence interval; RR, Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Last update: November 10, 2020; b. Davoudi-Monfared E, 2020; Rahmani H, 2020; c. Risk of bias downgraded by 2 levels: some concerns regarding adequate randomization, outcome measurement and selection of reported results, and high risk regarding deviations from intended interventions and missing data; d. Indirectness downgraded by 1 level: studies from a single country, therefore results in this population might not be generalizable to other settings; e. Imprecision downgraded by 2 levels: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants and events; f. Imprecision downgraded by 1 level: due to low number of events and/or participants; g. Risk of bias downgraded by 2 levels: some concerns regarding adequate randomization and selection of reported results, and high risk regarding deviations from intended interventions and missing data; h. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for no effect and low number of participants and events; i. Davoudi-Monfared

Results: Therapeutics

E, 2020; Rahmani H, 2020; SOLIDARITY, 2020; j. Inconsistency downgraded by 1 level: $I^2=71.2\%$; k. Imprecision downgraded by 2 levels: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm

3.12 Convalescent plasma transfusion

About the treatment under consideration

Convalescent plasma is plasma collected from patients that have recovered from an infectious disease and can be transfused to patients fighting an infection or can be used to manufacture immune globulin concentrates (plasma derived medicinal products). Possible explanations for the efficacy are that the antibodies from convalescent plasma might suppress viraemia and activate the complement system, thus promoting viral elimination. Antibody is most effective when administered shortly after the onset of symptoms, and a sufficient amount of antibody must be administered. Plasma transfusions may be associated with transfusion reactions such as allergic reactions, antibody-mediated enhancement of infection, transfusion-related acute lung injury (TRALI) and circulatory overload [109-111]. Rare complications include the transmission of infectious pathogens and red cell alloimmunization.

The European Commission (EC) and US Food and Drug Administration (FDA) published guidance on convalescent plasma collected from individuals who have recovered from COVID-19 [112, 113]. The EC guidance aims to facilitate a common approach across EU Member States to the donation, collection, testing, processing, storage, distribution and monitoring of convalescent plasma for the treatment of Covid-19 [112]. The FDA guidance provides recommendations on the pathways for use of investigational COVID-19 convalescent plasma; patient eligibility; collection of COVID-19 convalescent plasma, including donor eligibility and donor qualifications; labeling and record keeping. As COVID-19 convalescent plasma is regulated as an investigational product, three pathways for use are available in US: 1. Clinical Trials; 2. Expanded Access; 3. Single Patient Emergency IND [113, 114].

On August 23, 2020 the FDA issued an **emergency use authorization (EUA)** for investigational convalescent plasma for the treatment of COVID-19 in hospitalized patients [115].

Current US **NIH COVID-19 Treatment Guidelines** stated that there are insufficient clinical data to recommend either for or against the use of convalescent plasma for the treatment of COVID-19 (last update October 9, 2020) [116].

Withdrawn, suspended or terminated studies

1 RCT was found as withdrawn in US, NCT04467151 (did not obtain funding to proceed) and 1 RCT found as terminated in Italy, NCT04393727, the Promoter was changed and a new study promoted by AIFA started).

Results of publications

Li et al. 2020 published results from RCT (**ChiCTR200029757**) [117] conducted in 103 patients with COVID-19 (severe to critical) admitted to 7 centers in China. Convalescent plasma therapy added to standard treatment, compared with standard treatment alone, did not result in a statistically significant improvement in time to clinical improvement within 28 days (51.9% (27/52) of the convalescent plasma group vs 43.1% (22/51) in the control group (difference, 8.8% [95% CI, -10.4% to 28.0%]; hazard ratio [HR], 1.40 [95% CI, 0.79-2.49]; $p = 0.26$). Among those with severe disease,

(Re-) Konvaleszenzplasma von covid-19 Patient*innen, die sich von der Erkrankung bereits erholt haben

auch zur Herstellung von Immunglobulin-konzentraten verwendet

EMA & FDA Guidance zu CVP

FDA im August 2020: Emergency Authorization

US NIH COVID-19 Treatment Guidelines: insuffiziente Datenlage weder für noch gegen CVP

Li (China) RCT, 103 Pts (statt 200, wegen Mangel an Pts)

keine Unterschiede bei Endpunkten

the primary outcome was statistically significant in favour of convalescent plasma (91.3% (21/23) vs 68.2% (15/22) of the control group (HR, 2.15 [95% CI, 1.07-4.32]; $p = 0.03$); among those with life-threatening disease the primary outcome occurred in 20.7% (6/29) of the convalescent plasma group vs 24.1% (7/29) of the control group (HR, 0.88 [95% CI, 0.30-2.63]; $p = 0.83$) (P for interaction = 0.17). There was no significant difference in 28-day mortality (15.7% vs 24.0%; OR, 0.65 [95% CI, 0.29-1.46]; $p = 0.30$) or time from randomization to discharge (51.0% vs 36.0% discharged by day 28; HR, 1.61 [95% CI, 0.88-2.93]; $p = 0.12$). Two patients in the convalescent plasma group experienced adverse events within hours after transfusion that improved with supportive care. Interpretation of results is limited by early termination of the trial, which may have been underpowered to detect a clinically important difference.

Gharbharan et al. 2020 [118], published results as **preprint**, from prematurely **halted RCT (NCT04342182)**, performed on 86 patients with COVID-19 (moderate-critical) admitted to 14 centers in the Netherlands [118].

**RCT (Niederlande):
86 Pts.,**

Avendano-Sola et al. 2020 published as **preprint**, results of multi-center RCT (**NCT04345523**) [119]: All patients received standard of care treatment, including off-label use of marketed medicines, and were randomized 1:1 to receive one dose (250-300 mL) of CP from donors with IgG anti-SARS-CoV-2. The trial was stopped after first interim analysis due to the fall in recruitment related to pandemic control. With 81 patients randomized, there were no patients progressing to mechanical ventilation or death among the 38 patients assigned to receive plasma (0%) versus 6 out of 43 patients (14%) progressing in control arm. Mortality rates were 0% vs 9.3% at days 15 and 29 for the active and control groups, respectively. No significant differences were found in secondary endpoints.

**Sept 2020:
Publikation zu RCT
CVP vs. SOC**

**frühzeitiger Abbruch
wegen Mangel an
Rekrutierung: Interim
Analyse von 81 Pts**

Agarwal et al. 2020 [120] [121] reported results from open-label, parallel-arm, phase 2, multicentre, randomized controlled trial in India (**CTRI/2020/04/024775**) conducted on hospitalized, moderately ill confirmed COVID-19 patients (PaO₂/FiO₂: 200-300 or respiratory rate > 24/min and SpO₂ ≤ 93% on room air). 464 participants were enrolled; 235 and 229 in intervention and control arm, respectively. Composite primary outcome (progression to severe disease or all cause mortality at 28 days) was achieved in 44 (19%) participants in the intervention arm and 41 (18%) in the control arm (risk difference 0.008 [95% confidence interval -0.062 to 0.078]; risk ratio 1.04, 95% confidence interval 0.71 to 1.54

**Okt 2020
preprint RCT
(open-label)
Indien
464 Pts**

**kein Unterschied bei
Mortalität oder
Fortschreiten der
Krankheit**

Balcells et al. 2020 [122] reported, as preprint, results from open-label, single-center, randomized clinical trial performed in an academic center in Santiago, Chile, including 58 patients (**NCT04375098**). No benefit was found in the primary outcome (32.1% vs 33.3%, OR 0.95, 95% CI 0.32-2.84, $p > 0.99$) in the early versus deferred CP group. In-hospital mortality rate was 17.9% vs 6.7% (OR 3.04, 95% CI 0.54-17.2, $p = 0.25$), mechanical ventilation 17.9% vs 6.7% (OR 3.04, 95% CI 0.54-17.2, $p = 0.25$), and prolonged hospitalization 21.4% vs 30% (OR 0.64, 95%CI, 0.19-2.1, $p = 0.55$) in early versus deferred CP group, respectively. Viral clearance rate on day 3 (26% vs 8%, $p = 0.20$) and day 7 (38% vs 19%, $p = 0.37$) did not differ between groups. Two patients experienced serious adverse events within 6 or less hours after plasma transfusion.

**preprint RCT
(open-label)
Chile
58 Pts**

**kein Unterschied bei
Mortalität, Dauer des
Krankenhausaufenthalts
und künstlicher
Beatmung**

Simonovich et al 2020 [123] published results from RCT (NCT04383535) in hospitalised adult patients with severe Covid-19 pneumonia. A total of 228 patients were assigned to receive convalescent plasma and 105 to receive

**RCT
228 Patient*innen
kein Unterschied**

placebo. The median time from the onset of symptoms to enrollment in the trial was 8 days (interquartile range, 5 to 10), and hypoxemia was the most frequent severity criterion for enrollment. The infused convalescent plasma had a median titer of 1:3200 of total SARS-CoV-2 antibodies (interquartile range, 1:800 to 1:3200). At day 30 day, no significant difference was noted between the convalescent plasma group and the placebo group in the distribution of clinical outcomes according to the ordinal scale (odds ratio, 0.83 (95% confidence interval [CI], 0.52 to 1.35; P=0.46). Overall mortality was 10.96% in the convalescent plasma group and 11.43% in the placebo group, for a risk difference of -0.46 percentage points (95% CI, -7.8 to 6.8). Adverse events and serious adverse events were similar in the two groups.

Three more RCTs was found as preprint publications: AlQahtani et al. 2020 (NCT04356534); Libster et al. 2020 (NCT04479163; PAEPCC19; Plataforma PRIISA (1421)) and Ray et al. 2020 (CTRI/2020/05/025209) ; results will be presented after peer-review publication. The **Living Systematic Review with meta-analysis**, related to **seven RCTs**: Li et al. 2020 [117], Gharbharan et al. 2020 [118], Avendano-Sola et al. 2020 [141], Agarwal et al. 2020 [120], Simonovich [123], AlQahtani et al. 2020 and Libster et al. 2020 with **Summary of findings** table is provided in Table 3.12-1.

In summary, risk ratio related to all outcomes was not statistically significant different between convalescent plasma and standard care, except for the outcome - Viral negative conversion D7, RR 1.23 (1.04 to 1.46), with very low certainty of evidence.

**3 weitere RCTs in preprint
in SoF Tabelle präsentiert**

**Zusammenfassung der
SoF: kein Unterschied**

Results: Therapeutics

Table 3.12-1: Summary of findings table on Convalescent plasma compared to Standard Care for Mild/Moderate/Severe/Critical COVID-19 –(7 RCTs: Li, Gharbharan, Avendano-Sola, Agarwal, AlQahtani, Simonovich, Libster)

Convalescent plasma compared to Standard Care for Mild/Moderate/Severe/Critical COVID-19

Patient or population: Mild/Moderate/Severe/Critical COVID-19

Setting: Worldwide

Intervention: Convalescent plasma

Comparison: Standard Care

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard Care	Risk with Convalescent plasma				
Viral negative conversion D3	350 per 1,000	610 per 1,000 (270 to 1,000)	RR 1.74 (0.77 to 3.90)	470 (2 RCTs) ^g	⊕○○○ VERY LOW ^{5,6,8}	
Viral negative conversion D7	550 per 1,000	677 per 1,000 (572 to 803)	RR 1.23 (1.04 to 1.46)	342 (1 RCT) ^f	⊕○○○ VERY LOW ^{8,9,1}	
Clinical improvement D7	98 per 1,000	96 per 1,000 (29 to 313)	RR 0.98 (0.30 to 3.19)	103 (1 RCT) ¹	⊕○○○ VERY LOW ^{9,10}	
Clinical improvement D14-D28	570 per 1,000	604 per 1,000 (525 to 701)	RR 1.06 (0.92 to 1.23)	229 (3 RCTs) ¹⁰	⊕⊕○○ LOW ^{8,9}	
WHO progression score (level 6 or above) D7	47 per 1,000	27 per 1,000 (2 to 279)	RR 0.57 (0.05 to 5.99)	81 (1 RCT) ¹¹	⊕○○○ VERY LOW ^{10,12}	
WHO progression score (level 6 or above) D14-28	70 per 1,000	11 per 1,000 (1 to 211)	RR 0.16 (0.01 to 3.02)	81 (1 RCT) ¹¹	⊕○○○ VERY LOW ^{10,12}	zero events in the intervention arm
WHO progression score (level 7 or above) D7	182 per 1,000	184 per 1,000 (124 to 277)	RR 1.01 (0.68 to 1.52)	414 (2 RCTs) ²	⊕⊕○○ LOW ¹	
WHO progression score (level 7 or above) D14-28	155 per 1,000	140 per 1,000 (90 to 221)	RR 0.90 (0.58 to 1.42)	414 (2 RCTs) ²	⊕⊕○○ LOW ¹	
All-cause mortality D7	47 per 1,000	14 per 1,000 (4 to 53)	RR 0.30 (0.08 to 1.11)	414 (2 RCTs) ²	⊕⊕⊕○ MODERATE ⁸	
All-cause mortality D14-D28	148 per 1,000	127 per 1,000 (93 to 174)	RR 0.86 (0.63 to 1.18)	1098 (6 RCTs) ³	⊕⊕○○ LOW ^{8,7}	
Adverse events	280 per 1,000	299 per 1,000 (252 to 355)	RR 1.07 (0.90 to 1.27)	596 (3 RCTs) ³	⊕⊕⊕○ MODERATE ¹	
Serious adverse events	81 per 1,000	102 per 1,000 (67 to 155)	RR 1.26 (0.63 to 1.92)	763 (5 RCTs) ³	⊕○○○ LOW ^{1,1}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
 CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence
 High certainty: We are very confident that the true effect lies close to that of the estimate of the effect
 Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different
 Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect
 Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Results: Therapeutics

Explanations

a. Last update: December 10, 2020; b. Agarwal A, 2020; Li L, 2020; c. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, deviations from intended interventions, missing outcome data and selection of reported results; d. Inconsistency downgraded by 1 level: $I^2=89.9\%$; e. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for no effect and low number of participants; f. Agarwal A, 2020; g. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, missing outcome data and selection of reported results; h. Indirectness downgraded by 1 level: despite a multicentre design this is a single study from a single country, therefore results in this population might not be generalizable to other settings; i. Imprecision downgraded by 1 level: due to low number of participants; j. Li L, 2020; k. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization, deviation from intended intervention and outcome measurement; l. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants; m. AlQahtani M, 2020; Gharbharan A, 2020; Li L, 2020; n. Avendaño-Solà C, 2020; o. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization and outcome measurement; p. Avendaño-Solà C, 2020; Simonovich VA, 2020; q. AlQahtani M, 2020; Avendaño-Solà C; Agarwal A, 2020; Gharbharan A, 2020; Li L, 2020; Simonovich VA, 2020; r. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, deviation from intended intervention and missing data; s. Li L, 2020; Libster R, 2020; Simonovich VA, 2020; t. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for no effect and the possibility for harm and low number of participants; u. Avendaño-Solà C, 2020; Gharbharan A, 2020; Li L, 2020; Libster R, 2020; Simonovich VA, 2020; v. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, deviation from intended intervention, missing data and outcome measurement

3.13 Plasma derived medicinal products

Neutralizing monoclonal antibodies

As Marovich et al. 2020 [124] stated, **neutralizing monoclonal antibodies** to SARS-CoV-2 have the potential to be used for both prevention and treatment of infection. They can help to guide vaccine design and development as well. The main target of SARS-CoV-2 neutralizing monoclonal antibodies is the surface spike glycoprotein that mediates viral entry into host cells. Some products will include of a combination of 2 monoclonal antibodies targeting different sites on the spike protein. Due to long half-life of most monoclonal antibodies (approximately 3 weeks for IgG1), a single infusion should be sufficient. A potential limitation of monoclonal antibodies for treatment of COVID-19 is the unknown bioavailability of passively infused IgG in tissues affected by the disease, especially the lungs, which serve as a key target of SARS-CoV-2 infection. Due to the effect of viral diversity it will be important to monitor for the emergence of resistant viral mutations under selective pressure of monoclonal antibody treatment.

Possible disease enhancement include antibody-mediated enhancement of viral entry and replication in target cells (Fc-bearing monocytes or macrophages) and virus-antibody immune complexes and the associated cytokine release [124].

**neutralisierende
monoklonale Antikörper:
Prävention und
Behandlung**

**Halbwertszeit bis
3 Wochen von Vorteil**

**Nachteil: unbekannte
Bioverfügbarkeit der
infundierten Antikörper**

3.13.1 REGN-COV2 - combination of two monoclonal antibodies (REGN10933 and REGN10987)

REGN-COV2 is combination of two monoclonal antibodies (REGN10933 and REGN10987) which bind non-competitively to the critical receptor binding domain of the virus's spike protein, which diminishes the ability of mutant viruses to escape treatment and protects against spike variants that have arisen in the human population.

A **phase 3 prevention trial** evaluates REGNCOV2's ability to prevent infection among uninfected people who have had close exposure to a COVID-19 patient (such as the patient's housemate) at approximately 100 sites and is expected to enroll 2,000 patients in the U.S.; the trial will assess SARS-CoV-2 infection status.

REGN-COV2 has also moved into the **phase 2/3** portion of **two adaptive phase 1/2/3 trials** testing the cocktail's ability **to treat hospitalized and non-hospitalized (or "ambulatory") patients with COVID-19**. The two phase 2/3 treatment trials in hospitalized (estimated enrollment =1,850) and non-hospitalized (estimated enrollment =1,050) patients are planned to be conducted at approximately 150 sites in the U.S., Brazil, Mexico and Chile, and will evaluate virologic and clinical endpoints, with preliminary data expected later this summer.

On September 14, 2020 the University of Oxford and Regeneron Pharmaceuticals, Inc. announced that **RECOVERY** (Randomised Evaluation of COVID-19 thERapY will evaluate Regeneron's investigational anti-viral antibody cocktail, REGNCOV2, <https://www.recoverytrial.net/news/recovery-covid-19-phase-3-trial-to-evaluate-regeneron2019s-regn-cov2-investigational-antibody-cocktail-in-the-uk>. The phase 3 open-label trial in patients hospitalised with COVID-19 will

**Kombination aus 2
monoklonalen
Antikörpern: Casirivimab
+ Imdevimab**

**Phase 3
REGNCOV2 Studie
NIAID (NIH) Studie mit
2.000 Teilnehmer*innen**

**Behandlung von
hospitalisierten und
ambulanten
Patiente*innen
1.050 Pts.
In Planung**

**Sept 2020:
RECOVERY nimmt
REGNCOV2 als
Studienmedikament auf**

compare the effects of adding REGN-COV2 to the usual standard-of-care versus standard-of-care on its own.

Results of publication

On Oct 28, 2020 Regeneron Pharmaceuticals, Inc. announced **positive results** from an **ongoing phase 2/3 RCT** in the **COVID-19 outpatient setting** (ambulatory patients, n=799) on their website; the trial met the primary and key secondary endpoints. REGN-COV2 significantly reduced viral load and patient medical visits (hospitalizations, emergency room, urgent care visits and/or physician office/telemedicine visits), by 57% through day 29 (2.8% combined dose groups; 6.5% placebo; p=0.024) and by 72% in patients with one or more risk factor (including being over 50 years of age; body mass index greater than 30; cardiovascular, metabolic, lung, liver or kidney disease; or immunocompromised status) (combined dose groups; nominal p = 0.0065). Manufacturer will submit detailed results from this trial for publication, <https://www.prnewswire.com/news-releases/regeneron-covid-19-outpatient-trial-prospectively-demonstrates-that-regn-cov2-antibody-cocktail-significantly-reduced-virus-levels-and-need-for-further-medical-attention-301162255.html>.

Safety issue

On 30 October 2020, Regeneron Pharmaceuticals, Inc. received a recommendation from the independent data monitoring committee (IDMC) for the REGN-COV2 antibody cocktail treatment trials for COVID-19 that the current **hospitalized patient** trial be modified. Specifically, based on a potential safety signal and an unfavorable risk/benefit profile at this time, the IDMC recommends further enrollment of patients requiring high-flow oxygen or mechanical ventilation be placed on hold pending collection and analysis of further data on patients already enrolled. The IDMC also recommends continuing enrollment of hospitalized patients requiring either no or low-flow oxygen as the risk/benefit remains acceptable in these cohorts. Finally, the IDMC recommends continuation of the outpatient trial without modification, <https://investor.regeneron.com/news-releases/news-release-details/regn-cov2-independent-data-monitoring-committee-recommends>.

Regulatory update: On November 21, 2020, the U.S. Food and Drug Administration issued an **emergency use authorization (EUA)** for casirivimab and imdevimab to be administered together for the **treatment of mild to moderate COVID-19** in adults and pediatric patients (12 years of age or older weighing at least 40 kilograms [about 88 pounds]) with positive results of direct SARS-CoV-2 viral testing and who are **at high risk for progressing to severe COVID-19**. This includes those who are 65 years of age or older or who have certain chronic medical conditions. [125]

The **US COVID-19 Treatment Guidelines Panel** issued new recommendations on pharmacological treatment for patients with COVID-19 (as of December 3, 2020) [116]. In summary, related to the anti-SARS-CoV-2 monoclonal antibodies bamlanivimab and casirivimab plus imdevimab, in the earliest stages of infection, before the host has mounted an effective immune response, anti-SARS-CoV-2 antibody-based therapies may have their greatest likelihood of having an effect. In this regard, although there are insufficient data from clinical trials to recommend either for or against the use of any specific therapy in this setting, preliminary data suggests that outpatients may benefit from receiving anti-SARS-CoV-2 monoclonal antibodies early in the course of infection. The anti-SARS-CoV-2 monoclonal antibodies bamlanivimab and casirivimab plus imdevimab are available

Phase 2/3 RCT
799 ambulante Pts.

Firmenankündigung zu
positive Effekten

Endpunkte:
Reduktion der Viruslast
Arzt-/ Notfall-/
Spitalsbesuche

Sicherheitswarnung für
Kohorte hospitalisierte
und künstlich beatmete
Pts.

FDA: Notzulassung von
von REGN-COV2

für milde bis moderate
Erkrankung

Empfehlung des US
COVID-19 Treatment
Guidelines Panel

wenn, dann in sehr
frühem Stadium
aber insuffiziente
Datenlage

through Emergency Use Authorizations for outpatients who are at high risk for disease progression.

3.13.2 LY-CoV555 - neutralizing IgG1 monoclonal antibody (bamlanivimab) and LY-CoV016 - recombinant fully human monoclonal neutralizing antibody (etesevimab)

LY-CoV555 is a neutralizing IgG1 monoclonal antibody (mAb) directed against the spike protein of SARS-CoV-2. It is designed to block viral attachment and entry into human cells, thus neutralizing the virus, potentially preventing and treating COVID-19.

LY-CoV016 (also known as JS016) is a recombinant fully human monoclonal neutralizing antibody, which specifically binds to the SARS-CoV-2 surface spike protein receptor binding domain with high affinity and can effectively block the binding of the virus to the ACE2 host cell surface receptor.

Lilly has successfully completed enrollment and primary safety assessments of LY-CoV555 in a **phase 1** study of hospitalized patients with COVID-19 (NCT04411628) and long-term follow-up is ongoing.

BLAZE-1 (NCT04427501) is ongoing randomized, double-blind, placebo-controlled **phase 2** study designed to assess the efficacy and safety of LY-CoV555 and LY-CoV016 for the treatment of symptomatic COVID-19 in the outpatient setting. Across all treatment arms, the trial will enroll an estimated 800 participants.

A **phase 3** study for the prevention of COVID-19 in residents and staff at long-term care facilities (NCT04497987, BLAZE-2) is recently initiated.

In addition, LY-CoV555 is being tested in the National Institutes of Health-led ACTIV-2 and ACTIV-3 studies of ambulatory and hospitalized COVID-19 patients.

To generate additional efficacy and safety data, a pragmatic, open-label study enrolling patients treated with either monotherapy or combination therapy, with a focus on collecting data regarding hospitalizations, deaths and safety, planned to be initiated in October 2020.

US COVID-19 Treatment Guidelines (see above in casirivimab plus imdevimab section).

**2 weitere mAb:
LY-CoV555
(Bamlanivimab)**

**LY-CoV016
(Etesevimab)**

LY-CoV555: Phase 1

**BLAZE-1: RCT, Phase 2
800 Pts.
LY-CoV555 &
LY-CoV016**

**BLAZE-2: RCT, Phase 3
initiiert**

**NIH-Studien: ACTIV-2 and
ACTIV-3**

pragmatic trial in Planung

Empfehlung wie 3.13.1

Results of publications

Chen et al. 2020 [126] published interim analysis results of BLAZE-1, phase 2 RCT (NCT04427501), in 452 mild or moderate Covid-19 patients. One of three doses of neutralizing antibody LY-CoV555 appeared to accelerate the natural decline in viral load over time, whereas the other doses had not by day 11: 2800-mg dose of LYCoV555, the difference from placebo in the decrease from baseline was -0.53 (95% confidence interval [CI], -0.98 to -0.08; p=0.02. On days 2 to 6, the patients who received LY-CoV555 had a slightly lower severity of symptoms than those who received placebo. The percentage of patients who had a Covid-19–related hospitalization or visit to an emergency department was 1.6% in the LY-CoV555 group and 6.3% in the placebo group. In a post hoc analysis that was focused on high-risk subgroups (an age of ≥ 65 years or a BMI of ≥ 35), the percentage of hospitalization was 4.2% in the LY-CoV555 group and 14.6% in the placebo group. The safety outcomes were similar in intervention and placebo groups.

On October 7, 2020 Eli Lilly and Company **announced** data from an interim analysis of the BLAZE-1 clinical trial showed that combination therapy with two of Lilly's SARS-CoV-2 neutralizing antibodies reduced viral load, symptoms and COVID-related hospitalization and ER visits. The combination cohort enrolled recently diagnosed patients with mild-to-moderate COVID-19, who were assigned to 2800 mg of each antibody (n=112) or placebo (n=156). The combination therapy significantly reduced viral load at day 11 (p=0.011), meeting the primary endpoint of the study.

The combination therapy also met prespecified clinical endpoints, including the time-weighted average change from baseline in total symptom score from day 1 to 11 (p=0.009). The rate of COVID-related hospitalization and ER visits was lower for patients treated with combination therapy (0.9 percent) versus placebo (5.8 percent), a relative risk reduction of 84.5 percent (p=0.049). Combination therapy has been generally well tolerated with no drug-related serious adverse events.

Regulatory update:

On November 9, 2020, the **U.S. Food and Drug Administration** issued an **Emergency Use Authorization (EUA)** for the investigational monoclonal antibody therapy **bamlanivimab (previously LY-CoV555)** for the treatment of mild-to-moderate COVID-19 in adult and pediatric patients. Bamlanivimab is authorized for patients with positive results of direct SARS-CoV-2 viral testing who are 12 years of age and older weighing at least 40 kilograms (about 88 pounds), and who are at high risk for progressing to severe COVID-19 and/or hospitalization. This includes those who are 65 years of age or older, or who have certain chronic medical conditions, <https://www.fda.gov/news-events/press-announcements/coronavirus-covid-19-update-fda-authorizes-monoclonal-antibody-treatment-covid-19>. Bamlanivimab is not authorized for patients who are hospitalized due to COVID-19 or require oxygen therapy due to COVID-19. A benefit of bamlanivimab treatment has not been shown in patients hospitalized due to COVID-19. Monoclonal antibodies, such as bamlanivimab, may be associated with worse clinical outcomes when administered to hospitalized patients with COVID-19 requiring high flow oxygen or mechanical ventilation.

Phase 2 RCT
452 Pts.
milde/moderate
Erkrankung

Vorteil bei Endpunkten:
Reduktion der Viruslast
Artzt-/ Notfall-/
Spitalsbesuche

BLAZE-1
268 Pts,
Zwischenauswertung

Vorteil bei Endpunkten:
Reduktion der Viruslast
Artzt-/ Notfall-/
Spitalsbesuche

Ergebnisse in
Kombinationstherapie
gleich wie in
Monotherapie
LY-CoV555

November: FDA
EUA für bamlanivimab

für ambulante Pts mit
Risiko auf
Verschlechterung

nicht für bereits
hospitalisierte Pts.

3.13.3 AZD7442 - combination of two monoclonal antibodies (AZD8895 + AZD1061)

AZD7442 is a combination of two mAbs (AZD8895 + AZD1061) derived from convalescent patients with SARS-CoV-2 infection. Discovered by Vanderbilt University Medical Center and licensed to AstraZeneca in June 2020, the mAbs were optimised by AstraZeneca with half-life extension and reduced Fc receptor binding. The half-life extended mAbs should afford at least six months of protection from COVID-19.

NCT04507256 is a **phase 1**, first time in human, randomised, double-blind, placebo-controlled, and dose escalation study that aims to evaluate the safety, tolerability and pharmacokinetics of AZD7442 in healthy participants. Estimated study completion date is September 2021.

Should AZD7442 prove to be tolerated and have a favourable safety profile in the trial, AstraZeneca will progress it into larger late-stage **phase 2** and **phase 3** trials to evaluate its efficacy as a potential preventative and treatment approach against COVID-19, <https://www.astrazeneca.com/media-centre/press-releases/2020/phase-1-clinical-trial-initiated-for-monoclonal-antibody-combination-for-the-prevention-and-treatment-of-covid-19.html>.

AZD7442 Kombination aus 2 monoklonalen Antikörpern Vanderbilt University/ AstraZeneca

**Phase 1
Ende Sept 2021**

Phase 2 & 3 in Vorbereitung

3.14 Combination therapy – triple combination of interferon beta-1b, lopinavir–ritonavir and ribavirin vs. lopinavir–ritonavir or other triple combination of interferons

Hung et al. 2020 [103] present the results of the first randomised controlled trial (NCT04276688) on the triple combination of interferon beta-1b, lopinavir–ritonavir, and ribavirin, compared with lopinavir–ritonavir alone, in the treatment of patients admitted to hospital with mild to moderate COVID-19 in Hong-Kong. In this multicentre, prospective, open-label, randomised, phase 2 trial, 127 patients were randomly assigned (2:1) to a 14-day combination of lopinavir 400 mg and ritonavir 100 mg every 12 h, ribavirin 400 mg every 12 h, and three doses of 8 million international units of interferon beta-1b on alternate days (combination group) or to 14 days of lopinavir 400 mg and ritonavir 100 mg every 12 h (control group). Triple therapy was associated with a significant reduction in the duration of viral shedding (time to negative nasopharyngeal swab 7 days [IQR 5–11] in the combination group vs 12 days [8–15] in the control group; hazard ratio [HR] 4.37 [95% CI 1.86–10.24], $p=0.0010$), symptom alleviation (time to NEWS2 0 of 4 days [IQR 3–8] vs 8 days [7–9]; HR 3.92 [1.66–9.23], $p<0.0001$), and duration of hospital stay (9.0 days [7.0–13.0] vs 14.5 days [9.3–16.0]; HR 2.72 [1.2–6.13], $p=0.016$). There was no mortality in either group. The triple combination also suppressed IL-6 levels. Adverse events included self-limited nausea and diarrhoea with no difference between the two groups. No serious adverse events were reported in the combination group. One patient in the control group had a serious adverse event of impaired hepatic enzymes requiring discontinuation of treatment.

The **Living Systematic Review**, related to this RCT mentioned above, with **Summary of finding table** (https://covid-nma.com/living_data/index.php) is provided in Table 3.14-1.

Reduktion der Dauer der Virausscheidung, Symptomverbesserung, Dauer des Krankenhausaufenthalts

**kein Unterschied bei AE
keine Todesfälle in beiden Gruppen**

keine weiteren RCTs publiziert

Huang et al. 2020 [104] reported the results from a single-center, randomized, open-labeled, prospective clinical trial (**ChiCTR2000029387**). 101 eligible patients with mild to moderate COVID-19 were randomized into three groups: ribavirin (RBV) plus interferon- α (IFN- α), lopinavir/ritonavir (LPV/r) plus IFN- α , and RBV plus LPV/r plus IFN- α at a 1:1:1 ratio, with a 28-d follow-up. The median interval from baseline to SARS-CoV-2 nucleic acid negativity was 12 d in the LPV/r+IFN- α -treated group, as compared with 13 and 15 d in the RBV+IFN- α -treated group and in the RBV+LPV/r+IFN- α -treated group, respectively ($p=0.23$). The proportion of patients with SARS-CoV-2 nucleic acid negativity in the LPV/r+IFN- α -treated group (61.1%) was higher than the RBV+IFN- α -treated group (51.5%) and the RBV+LPV/r+IFN- α -treated group (46.9%) at day 14; however, the difference between these groups was calculated to be statistically insignificant. The RBV+LPV/r+IFN- α -treated group developed a significantly higher incidence of gastrointestinal adverse events than the LPV/r+IFN- α -treated group and the RBV+IFN- α -treated group.

RCT: 101 Pts

**3 Gruppen:
RBV+IFN
LPV/r+IFN
RBV+LPV/r**

kein Unterschied

Chinese RCT published by **Zheng et al. 2020** [127, 128] with three arms including 89 patients has evaluated the effect of Novaferon (the pharmaceutical which has similar properties of IFN-I but its antiviral activities has been greatly improved being at least 10 times more potent than human interferon α -2b) ($n=30$), Lopinavir/Ritonavir ($n=29$) and Novaferon + Lopinavir/Ritonavir ($n=30$) in COVID-19 patients. The groups treated with Novaferon alone or in combination with Lopinavir/Ritonavir showed significantly higher clearance rates on day 6 than the group treated with Lopinavir/Ritonavir alone, but the certainty on the evidence is very low. No serious adverse events were reported.

**RCT (China)
89 Pts.**

**3 Gruppen
Novaferon (IFN-I)
LPV/r
Novaferon + LPV/r**

**bessere Ergebnisse in IFN
Gruppen**

The **Living Systematic Review**, related to this RCT mentioned above, with **Summary of findings table** is provided in Table 3.14-1 continued.

Li C et al. 2020 [129] reported, as preprint, results from a multicenter, randomized controlled trial (**ChiCTR2000029638**) with aim to evaluate the efficacy and safety of recombinant super-compound interferon versus traditional interferon alpha added to baseline antiviral agents (lopinavir rSIFN-co-ritonavir or umifenovir) for the treatment of moderate-to-severe COVID-19. Recombinant super-compound interferon (rSIFN-co) is a new genetically engineered interferon. Participants received rSIFN-co (12 million international units [IU], twice daily) or interferon alpha (5 million IU, twice daily) nebulization added to baseline antiviral agents for no more than 28 days.

**Okt 2020:
preprint RCT
China
94 Pts.**

**rSIFN vs. IFN- α
beide in Kombination mit
Lopinavir oder
Ritonavir oder
Umifenovir**

94 patients hospitalized with moderate-to-severe COVID-19 were included in the safety set (46 patients assigned to rSIFN-co group, 48 to interferon alpha group). Individuals in the rSIFN-co group showed shorter time to clinical improvement (11.5 days vs 14.0 days; $p = 0.019$) as compared to those in the interferon alpha group. The overall rate of clinical improvement on day 28 was much higher in the rSIFN-co group than that in the interferon alpha group (93.5% vs 77.1%; difference, 16.4%; 95% condence interval 3% to 30%). The time to radiological improvement and the time to virus nucleic acid negative conversion were also much shorter in the rSIFN-co group (8.0 days vs 10.0 days, $p = 0.002$; 7.0 days vs 10.0 days, $p = 0.018$, respectively). Adverse events were reported in 13 (28.3%) patients in the rSIFN-co group and 18 (37.5%) patients in the interferon alpha group. No patients died during the study.

**signifikanter Unterschied
zugunsten von rSIFN-co
bei klinischer
Verbesserung und bei
Nebenwirkungen**

Table 3.14-1: Summary of findings table on **triple combination of interferon beta-1b, lopinavir-ritonavir and ribavirin** (1 RCT: Hung) - https://covid-nma.com/living_data/index.php

Summary of findings:						
Lopinavir + Ritonavir + Ribavirin + Interferon-b-1b compared to Lopinavir + Ritonavir for Mild/Moderate COVID-19						
Patient or population: Mild/Moderate COVID-19						
Setting: Worldwide						
Intervention: Lopinavir + Ritonavir + Ribavirin + Interferon-b-1b						
Comparison: Lopinavir + Ritonavir						
Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Lopinavir + Ritonavir	Risk with Lopinavir + Ritonavir + Ribavirin + Interferon-b-1b				
Incidence of viral negative conversion D7	902 per 1.000	875 per 1.000 (767 to 993)	RR 0.97 (0.85 to 1.10)	127 (1 RCT)	⊕⊕○○ LOW ^{a,b}	
WHO Clinical Progression Score (decrease in 1 point) (i.e., improvement) - not reported	-	-	-	-	-	outcome not yet measured or reported
Admission in ICU or death - not reported	-	-	-	-	-	outcome not yet measured or reported
Incidence of WHO progression score (level 6 or above) - not reported	-	-	-	-	-	outcome not yet measured or reported
Incidence of WHO progression score (level 7 or above) - not reported	-	-	-	-	-	outcome not yet measured or reported
All-cause mortality D7				127 (1 RCT)	⊕○○○ VERY LOW ^{a,c}	zero events in both groups

Results: Therapeutics

All-cause mortality D14-D28				127 (1 RCT)	⊕○○○ VERY LOW a,c	zero events in both groups
Adverse events D14-D28	488 per 1.000	478 per 1.000 (327 to 698)	RR 0.98 (0.67 to 1.43)	127 (1 RCT)	⊕⊕○○ MODERATE d,e	
Serious adverse events D14-D28	24 per 1.000	4 per 1.000 (0 to 94)	RR 0.16 (0.01 to 3.87)	127 (1 RCT)	⊕⊕○○ LOW ^{d,f}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

- a. Indirectness downgraded by 1 level: single study from a single country, therefore results in this population might not be generalizable to other settings
- b. Imprecision downgraded by 1 level: low number of participants
- c. Imprecision downgraded by 2 levels: no events in both groups and low number of participants
- d. Indirectness not downgraded: we presume that adverse event rate is not specific to a certain setting
- e. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants
- f. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants

Table 3.14-1 continued: Summary of findings tables on *Novaferon*, *Lopinavir/Ritonavir* and *Novaferon + Lopinavir/Ritonavir* (1 RCT: Zheng 2020)

Novaferon versus Lopinavir/Ritonavir

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect (95% CI)	Number of participants (studies)	Certainty of evidence
	Risk with Lopinavir/Ritonavir	Risk with Novaferon				
SARS-CoV-2 clearance	517 per 1000	567 per 1000	RR 1.10 (0.68 to 1.75)	52 more per 1000 (from 166 fewer to 388 more)	59	Very low
Progression of COVID-19 severity	143 per 1000	0 per 1000	RR 0.11 (0.01 to 1.97)	127 fewer per 1000 (from 141 fewer to 139 more)	56	Very low
Number with adverse events	138 per 1000	0 per 1000	RR 0.11 (0.01 to 1.91)	123 fewer per 1000 (from 137 fewer to 126 more)	59	Very low

Explanations of GRADE: Level of certainty was downgraded of one level for high risk of performance bias and unclear risk of selection bias, and further downgraded of two levels for very few events and small sample size

Novaferon versus Novaferon + Lopinavir/Ritonavir

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect (95% CI)	Number of participants (studies)	Certainty of evidence
	Risk with Novaferon + Lopinavir/Ritonavir	Risk with Novaferon				
SARS-CoV-2 clearance	700 per 1000	567 per 1000	RR 1.24 (0.84 to 1.83)	136 more per 1000 (from 91 fewer to 470 more)	60	Very low
Number with adverse events	100 per 1000	0 per 1000	RR 7.00 (0.38 to 129.93)	0 fewer per 1000 (from 0 fewer to 0 fewer)	60	Very low
Number with severe adverse events	Serious adverse events were not reported in either group.					Low
Progression of COVID-19 severity	None of the patients, with a moderate disease severity, had worsened disease.					Low

Explanations of GRADE: For the outcomes “SARS-CoV-2 clearance” and “Number with adverse events”, the level of certainty was downgraded of two levels for very few events and small sample size, and further downgraded of one level for small sample size. For the outcomes “Number with severe adverse events” and “Progression of COVID-19 severity”, the level of certainty was downgraded of one level for high risk of performance bias and unclear risk of selection bias, and further downgraded of one level for small sample size

Novaferon + Lopinavir/Ritonavir versus Lopinavir/Ritonavir

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect (95% CI)	Number of participants (studies)	Certainty of evidence
	Risk with Lopinavir/Ritonavir	Risk with Novaferon + Lopinavir/Ritonavir				
SARS-CoV-2 clearance	517 per 1000	700 per 1000	RR 1.35 (0.89 to 2.06)	181 more per 1000 (from 57 fewer to 548 more)	59	Very low
Progression of COVID-19 severity	143 per 1000	0 per 1000	RR 0.11 (0.18 to 2.96)	127 fewer per 1000 (from 141 fewer to 139 more)	56	Very low
Number with severe adverse events	138 per 1000	100 per 1000	RR 0.72 (0.18 to 2.96)	39 fewer per 1000 (from 113 fewer to 270 more)	59	Low

Explanations of GRADE: For the outcomes “SARS-CoV-2 clearance” and “Number with adverse events”, the level of certainty was downgraded of two levels for very few events and small sample size, and further downgraded of one level for small sample size. For the outcomes “Number with severe adverse events” and “Progression of COVID-19 severity”, the level of certainty was downgraded of one level for high risk of performance bias and unclear risk of selection bias, and further downgraded of one level for small sample size

Novaferon + Lopinavir/Ritonavir versus Lopinavir/Ritonavir

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Absolute effect (95% CI)	Number of participants (studies)	Certainty of evidence
	Risk with Lopinavir/Ritonavir	Risk with Novaferon + Lopinavir/Ritonavir				
SARS-CoV-2 clearance	517 per 1000	700 per 1000	RR 1.35 (0.89 to 2.06)	181 more per 1000 (from 57 fewer to 548 more)	59	Very low
Progression of COVID-19 severity	143 per 1000	0 per 1000	RR 0.11 (0.18 to 2.96)	127 fewer per 1000 (from 141 fewer to 139 more)	56	Very low
Number with severe adverse events	138 per 1000	100 per 1000	RR 0.72 (0.18 to 2.96)	39 fewer per 1000 (from 113 fewer to 270 more)	59	Low

Explanations of GRADE: For the outcomes “SARS-CoV-2 clearance” and “Progression of COVID-19 severity”, the level of certainty was downgraded of two levels for very few events and small sample size, and further downgraded of one level for small sample size. For the outcome “Number with severe adverse events” the level of certainty was downgraded of one level for high risk of performance bias and unclear risk of selection bias, and further downgraded of one level for small sample size.

3.15 Solnatide

About the treatment under consideration

The therapeutic molecule solnatide (INN) has been designed by APEPTICO (a privately-held biotechnology company from Vienna/Austria) for the therapeutic treatment of patients with Acute Respiratory Distress Syndrome (ARDS) and various forms of life-threatening Pulmonary Oedema (PPO). Solnatide is a synthetic peptide of less than 20 amino acids applied directly in the lower airways in the form of a liquid aerosol, aims to accelerate the dissolution of alveolar oedema and reduce barrier damage caused by Covid-19 in the lungs.

In April 2020, solnatide has been approved for Compassionate Use by the Austrian Federal Office for Safety in Health Care (BASG) for the treatment of patients infected by the novel coronavirus SARS-CoV-2 and subsequently developing severe pulmonary dysfunction (severe COVID-19), as well as by the Italian Medicines Agency and the Ethics Committee of the National Institute for Infectious Diseases (Lazzaro Spallanzani-Rome), within the compassionate use program of drugs undergoing clinical trials for the treatment of COVID-19 patients suffering from pulmonary oedema and acute respiratory distress syndrome.

APEPTICO Forschung und Entwicklung GmbH has signed, together with the “solnatide consortium”, the Grant Agreement ID: 101003595 with the European Commission to accelerate the process of making APEPTICO’s proprietary investigational medicinal product (IMP) solnatide available for medical treatment of patients severely affected by the novel coronavirus 2019 (SARS-CoV-2) disease, COVID-19; the Grant Agreement was made available via the Horizon2020 programme “Advancing knowledge for the clinical and public health response to the 2019-nCoV epidemic” (https://ec.europa.eu/commission/presscorner/detail/en/ip_20_386). Project started on 1 April 2020 and will end on 31 December 2021.

One ongoing randomised, double-blind, placebo controlled, parallel assignment trial with aim to assess efficacy and safety of 7 days orally inhaled 100 mg solnatide to treat pulmonary permeability oedema of 40 SARS-Cov-2 positive patients with moderate-to-severe ARDS is registered in EUdraCT register (EudraCT number 2020-001244-26), <https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001244-26/AT> [130].

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated studies related to solnatide in COVID-19 patients were found in ClinicalTrials.gov and EUdraCT registers [130].

Results of publications

No publications related to the RCTs of solnatide in COVID-19 patients were found [130].

Medikament gegen akutes Atemnotsyndrom

Verabreichung: Inhalation

April: BASG, AIFA lassen Solnatide für “Compassionate Use” zu

EC-Grant seit April für covid-19

bis Dezember 2021

1 laufender RCT mit 40 moderat bis schwer Covid-19 Erkrankten

ClinicalTrials.gov & EUdraCT: keine klinischen Studien registriert,

keine Publikation von RCT

3.16 Umifenovir (Arbidol®)

About the treatment under consideration

Umifenovir (Arbidol), an indole-derivative is a broad-spectrum drug against a wide range of enveloped and non-enveloped viruses: it interacts preferentially with aromatic amino acids, and it affects multiple stages of the virus life cycle, either by direct targeting viral proteins or virus-associated host factors. Umifenovir is currently being investigated as a potential treatment and prophylactic agent for COVID-19 caused by SARS-CoV2 infections in combination with both currently available and investigational HIV therapies (<https://pubchem.ncbi.nlm.nih.gov/compound/Arbidol>). Its use is only in China and Russia, since not approved by neither the FDA nor the EMA.

As Wang et al. 2020 recently published, arbidol efficiently inhibited SARS-CoV-2 infection in vitro (it appears to block virus entry by impeding viral attachment and release from the EIs) [131].

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated studies related to umifenovir were found in ClinicalTrials.gov and EUdraCT registers.

Results of publications

RCT published by **Yueping et al. 2020** (NCT04252885) [132] was an exploratory randomised (2:2:1) controlled trial, conducted in China, with the aim to assess the efficacy and safety of lopinavir/ritonavir or arbidol monotherapy in 86 patients with mild/moderate COVID-19. 34 of them assigned to lopinavir/ritonavir; 35 to arbidol and 17 with no antiviral medication as control, with follow-up of 21 days. The rate of positive-to-negative conversion of SARS-CoV-2 nucleic acid, as the primary endpoint, was similar between groups (all $p > 0.05$) and there were no differences between groups in the secondary endpoints, the rates of antipyresis, cough alleviation, or improvement of chest CT at days 7 or 14 (all $p > 0.05$). At day 7, eight (23.5%) patients in the LPV/r group, 3 (8.6%) in the arbidol group and 2 (11.8%) in the control group showed a deterioration in clinical status from moderate to severe/critical ($p = 0.206$). Related to adverse events, 12 (35.3%) patients in the lopinavir/ritonavir group and 5 (14.3%) in the arbidol group experienced adverse events during the follow-up period, and no AE occurred in the control group [132].

One publication [71] on the completed RCT (ChiCTR2000030254) about the efficacy and safety of favipiravir, in comparison with umifenovir, to treat Covid-19 patients was identified; Summary of findings table can be found in Section related to favipiravir.

RCT (IRCT20180725040596N2) published by **Nojomi et al. 2020**, as preliminary report in the format of preprints [133], is an open label randomized controlled trial, on effectiveness of umifenovir on 100 patients with COVID-19, assigned randomly to two groups of either hydroxychloroquine just on the 1st day followed by Kaletra (lopinavir-ritonavir) or hydroxychloroquine just on the 1st day followed by umifenovir 7-14 days based on severity of disease. The duration of hospitalization in umifenovir group was less than lopinavir-ritonavir arm significantly (7.2 versus 9.6 days; $p = 0.02$). Time to relief fever was similar across two groups (2.7 versus 3.1 days in umifenovir and lopinavir-ritonavir arms respectively). Peripheral oxygen saturation rate was different

**antivirales Medikament
zugelassen in China,
Russland, aber nicht EMA/
FDA**

1 in vitro Publikation

**ClinicalTrials.gov &
EudraCT: keine Studien
registriert**

**Yueping (China)
RCT, 86 Pts.
leichte/ moderate
Erkrankung**

**kein Unterschied
zwischen den Gruppen in
einigen
Surrogatendpunkten**

mehr AE

**1 RCT nur im preprint
(nicht peer-reviewed)**

**Okt 2020:
RCT (Iran)
100 Pts.**

**in Kombinationstherapie
kleine Vorteile**

after seven days of admission across two groups significantly (94% versus 92% in umifenovir and lopinavir-ritonavir groups respectively) ($p=0.02$).

Yethindra et al. 2020 [134] published results from exploratory randomized controlled study recruited 30 mild and moderate COVID-19 patients in Kyrgyzstan. No patient progressed toward severe and critical illness in either category. Pneumonia was ameliorated in 76.6% (23/30) of the patients, with moderate and potential amelioration in 36.6% and 40% of the patients, respectively. Many patients were observed to have significantly ameliorated pneumonia in the umifenovir category (86.6%, 13 of 15) compared to the control category (66.6%, 10 of 15). In addition, 66.6% of patients in the umifenovir category had potential pneumonia absorption. Only one patient presented with mild side effects in the umifenovir category, while one patient had cephalalgia; notably, no patient experienced severe side effects.

**November 2020
RCT, 30 Pts. Kirgistan**

The **Living Systematic Review**, related to these two RCTs mentioned above, with Summary of findings table (https://covid-nma.com/living_data/index.php) is presented in Table 3.16-1.

Results: Therapeutics

Table 3.16-1. Summary of findings table, on **umifenovir vs standard care** (2 RCTs: Yueping, Yethindra)

Umifenovir compared to Standard Care for Mild/Moderate COVID-19

Patient or population: Mild/Moderate COVID-19

Setting: Worldwide

Intervention: Umifenovir

Comparison: Standard Care

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard Care	Risk with Umifenovir				
Viral negative conversion D3 - not reported	-	-	-	-	-	outcome not yet measured or reported
Viral negative conversion D7	412 per 1,000	371 per 1,000 (181 to 756)	RR 0.90 (0.44 to 1.84)	52 (1 RCT) ^b	⊕○○○ VERY LOW ^d	
Clinical improvement D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement D14-D28 - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO progression score (level 6 or above) D7	63 per 1,000	46 per 1,000 (8 to 240)	RR 0.73 (0.13 to 3.98)	82 (2 RCT) ^b	⊕○○○ VERY LOW ^{d,f,g}	
WHO progression score (level 6 or above) D14-D28	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	30 (1 RCT) ^b	⊕○○○ VERY LOW ^{d,j}	zero events in both groups
WHO progression score (level 7 or above) D7	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	82 (2 RCT) ^b	⊕○○○ VERY LOW ^{d,k}	zero events in both groups
WHO progression score (level 7 or above) D14-D28	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	30 (1 RCT) ^b	⊕○○○ VERY LOW ^{d,j}	zero events in both groups
All-cause mortality D7	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	82 (2 RCT) ^b	⊕○○○ VERY LOW ^{d,k,m}	zero events in both groups
All-cause mortality D14-D28	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	82 (2 RCT) ^b	⊕○○○ VERY LOW ^{d,k,m}	zero events in both groups
Adverse events	0 per 1,000	0 per 1,000 (0 to 0)	RR 5.50 (0.32 to 94.06)	52 (1 RCT) ^b	⊕⊕○○ LOW ^{d,n}	zero events in control group
Serious adverse events	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	82 (2 RCT) ^b	⊕○○○ VERY LOW ^{d,k,n}	zero events in both groups

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
 CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence
 High certainty: We are very confident that the true effect lies close to that of the estimate of the effect
 Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different
 Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect
 Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Last update: November 13, 2020; b. Yueping L, 2020; c. Indirectness downgraded by 1 level: single study from a single institution, therefore results in this population might not be generalizable to other settings; d. Imprecision downgraded by 2 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for harm and very low number of participants; e. Yethindra V, 2020; Yueping L, 2020; f. Risk of bias downgraded by 1 level: some concerns around deviation from intended intervention in both studies, some concerns in one study

Results: Therapeutics

regarding randomization, outcome measurement, and selection of reported result; g. Indirectness downgraded by 1 level: results are mainly from a single study from a single institution, therefore results in this population might not be generalizable to other settings.; h. Yethindra, 2020; i. Risk of bias downgraded by 1 level: some concerns regarding randomization, deviations from intended intervention, outcome measurement, and selection of the reported results; j. Imprecision downgraded by 2 levels: no events in both groups and very low number of participants; k. Risk of bias downgraded by 1 level: some concerns regarding deviations from intended intervention in both studies, some concerns regarding randomization and selection of reported result in one study; l. Risk of bias downgraded by 1 level: some concerns regarding randomization, deviations from intended intervention, and selection of the reported results; m. Indirectness downgraded by 1 level: results from two single-institution studies, therefore results in the population might not be generalizable to other settings.; n. We presume that the adverse event rates, and the corresponding relative risks, is similar across diverse settings; therefore not downgraded for indirectness

3.17 Dexamethasone and other corticosteroids

About the drug under consideration

Dexamethasone is a long-acting glucocorticoid which is used principally as an anti-inflammatory or immunosuppressant agent. Daily regimen of dexamethasone 6 mg once daily is equivalent to 160 mg of hydrocortisone, 40 mg of prednisone, and 32 mg of methylprednisolone. The proposed mechanism of glucocorticoids in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) involves the mitigation of an excessive immune response that can lead to acute respiratory distress syndrome (ARDS) and multi-organ failure. ARDS develops in approximately 20% of COVID-19 patients and is linked to multi-organ failure through cytokine release syndrome [135, 136].

Dexamethasone is authorised at national level in the EU and is used in a wide range of conditions, including rheumatic problems, skin diseases, severe allergies, asthma and chronic obstructive lung disease. The UK has approved dexamethasone for the treatment of Covid-19 on June 16, 2020 [137].

CHMP is currently evaluating Dexamethasone Taw for a marketing authorisation for the treatment of hospitalised adult patients with COVID-19 [138].

On September 18, 2020 EMA announced that CHMP has completed its review of results from the RECOVERY dexamethasone study arm. EMA is endorsing the use of dexamethasone in adults and adolescents (from 12 years of age and weighing at least 40 kg) who require supplemental oxygen therapy. In all cases, the recommended dose in adults and adolescents is 6 milligrams once a day for up to 10 days. Companies that market dexamethasone medicines can request this new use to be added to their product's license by submitting an application to national medicines agencies or to EMA [139].

Based on results of the RECOVERY Trial described below, the US COVID-19 Treatment Guidelines Panel **recommends using dexamethasone** (at a dose of 6 mg per day for up to 10 days) in patients with COVID-19 who are mechanically ventilated (**AI**) and in patients with COVID-19 who require supplemental oxygen but who are not mechanically ventilated (**BI**). The Panel **recommends against** using dexamethasone in patients with COVID-19 who do not require supplemental oxygen (**AI**) [61]. If dexamethasone is not available, the Panel **recommends using** alternative glucocorticoids such as **prednisone, methylprednisolone, or hydrocortisone (AIII)** [61]. For more details, see also section on remdesivir.

The WHO panel made two recommendations: a **strong recommendation** (based on moderate certainty evidence) **for systemic** (i.e. intravenous or oral) **corticosteroid therapy** (e.g. 6 mg of dexamethasone orally or intravenously daily or 50 mg of hydrocortisone intravenously every 8 hours) for 7 to 10 days in **patients with severe and critical COVID-19**, and a **conditional recommendation** (based on low certainty evidence) **not to use corticosteroid therapy** in **patients with non-severe COVID-19** [140, 141].

**Glukokortikoide:
entzündungshemmend**

**nationale, nicht EMA
Zulassung, UK: Zulassung
im Juni für Covid-19**

**EMA- CHMP:
Zulassungsantrag von
Taw Pharma (Sept 2020)**

**Sept 2020:
basierend auf Ergebnissen
aus RECOVERY
EMA (Rasch-)Zulassung für
Pts mit (künstlicher)
Beatmung oder Sauerstoff
Supplementierung**

**Empfehlungen des US
COVID-19 Treatment
Guidelines Panel: bei
künstlich beatmeten
Patient*innen, nicht
jedoch bei nicht
beatmeten Pts.**

**WHO-Empfehlung für Pts.
mit schwerer oder
kritischer Erkrankung**

Withdrawn, suspended or terminated studies

Two RCTs were found as terminated: RCT - NCT04327401 (CoDEX), related to dexamethasone, in 299 COVID-19 patients with moderate and severe ARDS in Brazil, the Data Monitoring Committee recommended to stop the trial based on the Recovery Trial results, which was accepted by the CoDEX Steering Committee. NCT04344288 (CORTI-Covid) on prednisone in France, terminated due Competent Authority decision. DEXA-COVID trial (NCT04325061, EudraCT 2020-001278-31) on dexamethasone, is written as suspended (lack of enrollment) in ClinicalTrials.gov, but as ongoing in EudraCT register. The results of this RCT are not yet published [33]. 1 RCT in US (NCT04360876) is withdrawn because funding not received.

2 abgeschlossene RCTs
1 abgebrochener RCT
wegen (besseren
Ergebnissen in) Rovey
Trial in Brasilien

1 eingestellter RCT –
wegen Mangel an
Rekrutierung

Results of publications

The RCT with the largest number of included COVID-19 patients is RCTs of dexamethasone arm of the **RECOVERY trial** in Covid-19 patients (NCT04381936, EudraCT 2020-001113-21) [142]. The primary outcome was all-cause mortality within 28 days after randomization; further analyses were specified at 6 months.

größter RCT: RECOVERY

2.104 Pts

Results from preliminary report of the RECOVERY trial are related to the comparison of oral or intravenous dexamethasone 6 mg given once daily for up to ten days (2104 patients) plus the usual standard of care vs. usual care alone (4321 patients). Authors showed that overall, 482 (22.9%) patients allocated dexamethasone and 1110 (25.7%) patients allocated usual care died within 28 days (age adjusted rate ratio [RR] 0.83; 95% confidence interval [CI] 0.75 to 0.93; P<0.001). The proportional and absolute mortality rate reductions varied significantly depending on level of respiratory support at randomization (test for trend p<0.001): dexamethasone reduced deaths by one-third in patients receiving invasive mechanical ventilation (29.3% vs. 41.4%, RR 0.64 [95% CI 0.51 to 0.81]), by one-fifth in patients receiving oxygen without invasive mechanical ventilation (23.3% vs. 26.2%, RR 0.82 [95% CI 0.72 to 0.94]), but did not reduce mortality in patients not receiving respiratory support at randomization (17.8% vs. 14.0%, RR 1.19 [95% CI 0.91 to 1.55]). Allocation to dexamethasone was associated with a shorter duration of hospitalization than usual care (median 12 days vs. 13 days) and a greater probability of discharge within 28 days (rate ratio 1.10 [95% CI 1.03 to 1.17]) with the greatest effect seen among those receiving invasive mechanical ventilation at baseline (11.5 by chi-square test for trend). The risk of progression to invasive mechanical ventilation was lower among those allocated dexamethasone vs. usual care (risk ratio 0.92 [95% CI 0.84 to 1.01]). Analyses are ongoing regarding cause-specific mortality, the need for renal dialysis or hemofiltration, and the duration of ventilation [142, 143].

Reduktion der Mortalität
RR -30% bei Pts. mit
künstlicher Beatmung

RR -20% bei Pts. mit
Sauerstoff ohne invasive
Beatmung

ohne Effekt auf Mortalität
bei Pts ohne Untestützung
bei Beatmung

zusätzlich: kürzere
Hospitalisierung

The **CoDEX trial (NCT04327401)** randomized 299 patients in 41 ICUs in Brazil with moderate or severe ARDS and COVID-19 to open-label high-dose dexamethasone (20 mg/d for 5 days, then 10 mg/d for 5 days) vs usual care alone, with the primary outcome ventilator-free days through day 28, which were greater in patients randomized to dexamethasone (6.6 vs 4.0, p=0.04). 28-day mortality was not significantly different between patients randomized to corticosteroids vs usual care (56.3% vs 61.5%, p=0.83); stopping the study early when RECOVERY results were announced resulted in a sample size that was underpowered to adequately evaluate the effect of corticosteroids on mortality and other secondary outcomes [144, 145].

CoDEX
299 Pt (Brasilien)

kein signifikanter
Unterschied, aber wegen
Abbruch "underpowered"
für valide Ergebnisse

The **CAPE COVID trial (NCT02517489)** was blinded, placebo-controlled trial randomized 149 patients in 9 ICUs in France with severe respiratory disease from COVID-19 to low-dose hydrocortisone (200 mg/d infusion, tapered per protocol) vs placebo. The primary outcome of 21-day treatment failure, defined as death or ongoing respiratory support with mechanical ventilation or high-flow oxygen, occurred in 42.1% of patients randomized to hydrocortisone vs 50.7% of those randomized to placebo ($p=0.29$) [145, 146].

The **REMAP-CAP trial (NCT02735707)**, an existing multicenter, multinational adaptive platform trial for pneumonia, randomized 403 patients with severe COVID-19 (in the intensive care unit and receiving respiratory or cardiovascular organ support) to 1 of 3 open-label groups: fixed low-dose hydrocortisone, shock-dependent hydrocortisone, or no hydrocortisone. The primary study outcome was days patients remained alive and free of organ support to day 21. The Bayesian model found that fixed-dose hydrocortisone (93% probability), as well as shock-dependent hydrocortisone (80% probability), were both likely superior to no hydrocortisone, but data were insufficient to confirm a single optimal regimen. In addition, the probabilities did not meet the prespecified probabilities to define success [145, 147].

MetCOVID trial (NCT04343729) was parallel, double-blind, placebo-controlled, randomized, phase IIb clinical trial, performed with hospitalized patients aged ≥ 18 years with clinical, epidemiological and/or radiological suspected COVID-19, at a tertiary care facility in Brazil. 416 patients were randomly allocated (1:1 ratio) to receive either intravenous methylprednisolone (0.5 mg/kg) or placebo (saline solution), twice daily, for 5 days. Mortality at day 28 was not different between groups. A subgroup analysis showed that patients over 60 years in the methylprednisolone group had a lower mortality rate at day 28. Patients in the methylprednisolone arm tended to need more insulin therapy, and no difference was seen in virus clearance in respiratory secretion until day 7 [148].

GLUCOCOVID trial (EudraCT 2020-001934-37) was multicentric, partially randomized, preference, open-label trial, including adults with COVID-19 pneumonia, impaired gas exchange and biochemical evidence of hyperinflammation, aimed to determine whether a 6-day course of intravenous methylprednisolone improves outcome in patients with SARS CoV-2 infection at risk of developing Acute Respiratory Distress Syndrome (ARDS). Patients were assigned to standard of care (SOC), or SOC plus intravenous methylprednisolone (40mg/12h 3 days, then 20mg/12h 3 days). The use of methylprednisolone was associated with a reduced risk of the composite endpoint in the intention-to-treat, age-stratified analysis (combined risk ratio -RR- 0.55 [95% CI 0.33-0.91]; $p=0.024$). In the per-protocol analysis, RR was 0.11 (0.01-0.83) in patients aged 72 yr or less, 0.61 (0.32-1.17) in those over 72 yr, and 0.37 (0.19-0.74, $p=0.0037$) in the whole group after age-adjustment by stratification. The decrease in C-reactive protein levels was more pronounced in the methylprednisolone group ($p=0.0003$). Hyperglycaemia was more frequent in the methylprednisolone group [148].

CAPE COVID
149 Pts (Frankreich)
bessere Ergebnisse mit
hydrocortisone

REMAP-CAP
403 Pts (UK, CA, USA)
bessere Ergebnisse mit
hydrocortisone

MetCOVID
418 Pts (Brasilien)
methylprednisolone
kein Unterschied
zwischen Gruppen bei
Mortalität
methylprednisolone
Subgruppenanalyse:
>60 Jahre bessere
Ergebnisse

GLUCOCOVID
85 Pts (Spanien)
Methylprednisolone
bessere Ergebnisse bei
„composite“ Endpunkten

Ergebnisse sind ebenfalls
alters-abhängig

Edalatifard et al. 2020 [149] published results of a single-blind, randomized, controlled, clinical trial involving severe hospitalized patients with confirmed COVID-19 at the early pulmonary phase of the illness in Iran (IRCT20200404046947N1). Sixty-eight eligible patients underwent randomization (34 patients in each group) The percentage of improved patients was significantly higher in the methylprednisolone group than in the standard care group (32 (94.1%) vs 16 (57.1%); $P = 0.001$) and the mortality rate was significantly lower in the methylprednisolone group (2 (5.9%) vs 12 (42.9%); $P < 0.001$). Patients in the methylprednisolone intervention group had a significantly increased survival time compared with the patients in the standard care group [Log rank test: $P < 0.001$; Hazard ratio: 0.293; 95% CI: 0.154-0.555]. A total of two patients in each group (5.8% and 7.1% respectively) showed severe adverse events between initiation of treatment and the end of the study. There were one infection and one edema adverse event in the methylprednisolone group and two shock adverse events in the standard care group. Following the use of high dose of corticosteroids, most of the patients required insulin due to their known or hidden diabetes, and the insulin requirement was increased in the intervention group especially in diabetic and overweight patients.

Farahani et al. 2020 [150] reported, as preprint, results from phase 2, double-blind, randomized, clinical trial in 29 adults with intermediate or severe COVID-19 with PaO₂/FiO₂ less than 300 and progressive disease unresponsive to standard treatments admitted to the intensive care unit (ICU) (IRCT20200406046963N1): The investigation group received the recommended regimen plus methylprednisolone (1000mg/day for three days) and oral prednisolone 1mg/kg with tapering of dose within ten days. There was no mortality among the patients receiving the methylprednisolone treatment, but the mortality was high in patients without methylprednisolone therapy. In addition to improvement of respiratory outcome, Glasgow Coma Scale (GCS) of methylprednisolone group significantly ($p < 0.001$) improved also.

Results from three unpublished studies were found related to hydrocortisone (NCT04348305), methylprednisolone (NCT04244591) and dexamethasone (NCT04325061), which included small number of COVID-19 patients (from 19 to 47), in comparisons to placebo or standard care. RCTs results, the meta-analysis results and SoF table will be updated after results are published in peer-review journals.

Meta-analysis data on high, low and very low certainty of evidence, related to effectiveness and safety of dexamethasone and other corticosteroids reported in 7 RCTs, can be found in the Summary of Findings Table 3.17-1. In summary, according to the results of six RCTs with high certainty of evidence, corticosteroids reduce the risk of all-cause mortality D14-28 in COVID-19 patients /RR 0.90 (95% CI 0.83 to 0.97); absolute effect estimate 25 fewer per 1000 (95% CI from 23 fewer to 27 fewer). The same is true for outcomes WHO progression score level 6 or above D14-28 (RR 0.87, 95% CI 0.78 to 0.97, low certainty of evidence, 3 RCTs) and WHO progression score level 7 or above D14-28 RR 0.88, 95% CI 0.79 to 0.98, high certainty of evidence, 4 RCTs).

**Okt 2020:
RCT (Iran)
68 Pts.**

schwere Erkrankung

**signifikante Ergebnisse
bei klinischer
Verbesserung und bei
Mortalität**

**Phase 2 RCT
(Iran)
29 Pts.**

**signifikante Vorteile
bei Mortalität**

3 weitere kleine Studien

Metaanalyse von 7 RCTs

**Reduktion von
Gesamtmortalität
Verbesserung der
klinischen Symptomatik**

Results: Therapeutics

Table 3.17-1: Summary of findings table, on **dexamethasone and other corticosteroids** (7 RCTs: Horbey, Tomazini, Dequin, REMAP-CAP Investigators, Jeronimo, Corral, Edalatifard)

Corticosteroids compared to Standard Care/Placebo for Mild/Moderate/Severe/Critical COVID-19

Patient or population: Mild/Moderate/Severe/Critical COVID-19

Setting: Worldwide

Intervention: Corticosteroids

Comparison: Standard Care/Placebo

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard Care/Placebo	Risk with Corticosteroids				
Viral negative conversion D3 - not reported	-	-	-	-	-	Outcome not yet measured or reported
Viral negative conversion D7	474 per 1,000	476 per 1,000 (360 to 635)	RR 1.01 (0.78 to 1.34)	212 (1 RCT) ^b	⊕○○○ VERY LOW ^{5,6,8}	
Clinical improvement D7 - not reported	-	-	-	-	-	Outcome not yet measured or reported
Clinical improvement D14-28	620 per 1,000	775 per 1,000 (506 to 1,000)	RR 1.25 (0.82 to 1.90)	6724 (2 RCTs) ^f	⊕○○○ VERY LOW ^{9,10}	
WHO progression score level 6 or above D7 - not reported	-	-	-	-	-	Outcome not yet measured or reported
WHO progression score level 6 or above D14-28	720 per 1,000	626 per 1,000 (562 to 698)	RR 0.87 (0.78 to 0.97)	512 (3 RCTs) ⁱ	⊕⊕○○ LOW ⁶	
WHO progression score level 7 or above D7 - not reported	-	-	-	-	-	Outcome not yet measured or reported
WHO progression score level 7 or above D14-28	254 per 1,000	224 per 1,000 (201 to 249)	RR 0.88 (0.79 to 0.98)	6937 (4 RCTs) ^{im}	⊕⊕⊕⊕ HIGH	
All-cause mortality D7	246 per 1,000	187 per 1,000 (128 to 271)	RR 0.76 (0.52 to 1.10)	416 (1 RCT) ^b	⊕⊕○○ LOW ^{5,8}	
All-cause mortality D14-28	27 per 100	25 per 100 (23 to 27)	RR 0.90 (0.83 to 0.97)	7591 (6 RCTs) ^h	⊕⊕⊕⊕ HIGH	
Adverse events	68 per 1,000	101 per 1,000 (7 to 1,000)	RR 1.49 (0.11 to 20.63)	363 (2 RCTs) ^b	⊕○○○ VERY LOW ^{5,6,8}	
Serious adverse events	66 per 1,000	75 per 1,000 (41 to 137)	RR 0.88 (0.48 to 1.60)	617 (5 RCTs) ^f	⊕○○○ VERY LOW ^{5,8}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Results: Therapeutics

Explanations: a. Last update: November 10, 2020; b. Prado Jeronimo CM, 2020; c. Risk of bias downgraded by 1 level: high risk due to missing data; d. Indirectness downgraded by 1 level: single study from a single institution, therefore results in this population might not be generalizable to other settings; e. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for no effect and low number of participants; f. Horby P (RECOVERY Trial), 2020; Tomazini BM, 2020; g. Risk of bias downgraded by 1 level: some concerns regarding deviations from intended intervention and outcome measurement; h. Inconsistency downgraded by 1 level: $I^2=74.1\%$; i. Imprecision downgraded by 1 level: due to wide confidence interval consistent with the possibility for benefit and the possibility for no effect; j. Corral-Gudino L, 2020; Dequin P-F, 2020; Tomazini BM, 2020; k. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, deviations from intended interventions and outcome measurement; l. Imprecision downgraded by 1 level: due to low number of events and/or participants; m. Corral-Gudino L, 2020; Dequin P-F, 2020; Horby P (RECOVERY Trial), 2020; Tomazini BM, 2020; n. Angus DC, 2020; Corral-Gudino L, 2020; Dequin P-F, 2020; Horby P (RECOVERY Trial), 2020; Prado Jeronimo CM, 2020; Tomazini BM, 2020; o. Corral-Gudino L, 2020; Tomazini BM, 2020; p. Inconsistency downgraded by 1 level: $I^2=81.6\%$; q. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants; r. Angus DC, 2020; Corral-Gudino L, 2020; Edalatifard M, 2020; Dequin P-F, 2020; Tomazini BM, 2020; s. Risk of bias downgraded by 1 level: some concerns or high risk regarding adequate randomization, deviations from intended interventions, missing data and outcome measurement

3.18 Anakinra (Kineret®)

About the drug under consideration

Anakinra (Kineret®) is an immunosuppressive medicine, a copy of a natural human protein - 'human interleukin 1 receptor antagonist' (r-metHuIL-1ra, produced in Escherichia coli cells by recombinant DNA technology). Anakinra neutralises the biologic activity of interleukin-1 α (IL-1 α) and interleukin-1 β (IL-1 β) by competitively inhibiting their binding to interleukin-1 type I receptor (IL-1RI). Interleukin-1 (IL-1) is a pivotal pro-inflammatory cytokine mediating many cellular responses including those important in synovial inflammation. Anakinra is not authorised in Covid-19 patients (EMA, FDA).

The US COVID-19 Treatment Guidelines Panel stated that there are insufficient data to recommend either for or against Interleukin-1 inhibitors (e.g., anakinra) therapy in patients with COVID-19 disease [61].

Withdrawn, suspended or terminated studies

One RCT was found as suspended – ANACONDA (NCT04364009) –due to efficiency and safety reasons, after enrolment of 71 hospitalized COVID-19 patients in France. The intermediate review of data from this clinical trial showed early excess mortality in the group of patients treated with anakinra combined with standard optimized care, compared to the group of patients treated with standard optimized care alone. On October 29, 2020, the French National Agency for Medicines and Health Products Safety (ANSM) announced that inclusions in clinical trials evaluating anakinra in the treatment of COVID-19 are suspended due to safety information regarding the ANACONDA-COVID-19 clinical trial, <https://ansm.sante.fr/S-informer/Actualite/Suspension-des-inclusions-en-France-dans-les-essais-clinique-évaluant-l-anakinra-dans-la-prise-en-charge-de-la-COVID-19-Point-d-information>.

One RCT was found as terminated: NCT04366232 (JAKINCOV), due investigator decision in France, on anakinra alone and in combination with ruxolitinib.

Results of publications

Until now no scientific publication on RCTs of anakinra (Kineret®) in Covid-19 patients could be identified.

3.19 Colchicine

About the drug under consideration

Colchicine is an alkaloid isolated from the autumn crocus, Colchicum autumnale, with anti-gout and anti-inflammatory activities. Colchicine is available throughout the world in a generic form [151].

Colchicine is not authorised in Covid-19 patients (EMA, FDA).

**Immunsuppressivum,
humaner Interleukin-1
Rezeptorantagonist**

**EMA-Zulassung für
Rheumatoide Arthritis seit
2002**

**mehrere laufende Studien,
Empfehlung des US COVID-
19 Treatment Guidelines
Panel: insuffiziente
Datenlage**

**ANACONDA (Frankreich)
71 hospitalisierte Pts**

**wegen Sicherheit
abgebrochen**

**JAKINCOV (Frankreich)
abgebrochen**

**keine Publikation
eines RCTs**

**toxisches Alkaloid
wirkt als Zellgift
(Mitosehemmung)**

generisch

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on colchicine in ClinicalTrials.gov and EUdraCT registers.

keine Studien

Results of publications

Deftereos et al. 2020 [152] reported results from open-label, randomized controlled trial (NCT04326790) on 105 patients hospitalized with COVID-19 in 16 tertiary hospitals in Greece (randomization in a 1:1 allocation to either standard medical treatment or colchicine with standard medical treatment). Patient recruitment was terminated on April 27, 2020, because of slow enrollment as a result of the rapid flattening of the curve of COVID-19 cases in Greece. The clinical primary end point rate was 14.0% in the control group (7 of 50 patients) and 1.8% in the colchicine group (1 of 55 patients) (odds ratio, 0.11; 95% CI, 0.01-0.96; $p=0.02$). Mean (SD) event-free survival time was 18.6 (0.83) days in the control group vs 20.7 (0.31) in the colchicine group (log rank $p=0.03$). Adverse events were similar in the 2 groups, except for diarrhea, which was more frequent with colchicine group than the control group (25 patients [45.5%] vs 9 patients [18.0%]; $p=0.003$).

**1 publizierter RCT
(Griechenland):
105 Pts.**

**klinisch gering-relevanter
Unterschied bei
Verbesserung der
Erkrankung**

**viele Surrogatendpunkte
niedrige Evidenz**

Salehzadeh et al. 2020 [153] reported results (as preprint) from prospective, open-label, randomized and double blind clinical trial, in 100 patients hospitalized with COVID-19 in Iran (IRCT20200418047126N1). Patients were randomized in a 1:1 allocation, to either standard medical treatment (hydroxychloroquine) or colchicine with standard medical treatment. Colchicine group were received 1 mg tablet of colchicine daily alongside the hydroxychloroquine for 6 days. Duration of hospitalisation and duration of fever were significantly different between patients groups, in favour of colchicine ($p<0.05$). Although in colchicine group dyspnea was improved more rapid than the placebo group, difference was not statistically significant. None of the patients died or were readmitted.

**RCT preprint (Iran)
100 Pts.**

kein Unterschied

Lopes et al. 2020 [154], reported (as preprint) interim results of a single-center, randomized, double-blinded, placebo controlled clinical trial of colchicine for the treatment of 38 moderate to severe COVID-19 patients in Brazil. Thirty-five patients (18 for placebo and 17 for colchicine) completed the study. Median (and interquartile range) time of need for supplemental oxygen was 3.0 (1.5- 6.5) days for the colchicine group and 7.0 (3.0-8.5) days for placebo group ($p=0.02$). Median (IQR) time of hospitalization was 6.0 (4.0-8.5) days for the colchicine group and 8.5 (5.5-11.0) days for placebo group ($p=0.03$). At day 2, 53% vs 83% of patients maintained the need for supplemental oxygen, while at day 7 the values were 6% vs 39%, in the colchicine and placebo groups, respectively (log rank; $p=0.01$). Hospitalization was maintained for 53% vs 78% of patients at day 5 and 6% vs 17% at day 10, for the colchicine and placebo groups, respectively (log rank; $p=0.01$). One patient per group needed admission to ICU. No recruited patient died. At day 4, patients of colchicine group presented significant reduction of serum C-reactive protein compared to baseline ($p<0.001$). The majority of adverse events were mild and did not lead to patient withdrawal. Diarrhea was more frequent in the colchicine group ($p=0.17$). Cardiac adverse events were absent.

**RCT preprint
(Brasilien)
38 Pt.**

**Reduktion von Sauerstoff
Supplementierung und
von Hospitalisierung**

Summary of Finding table related to colchicine compared to standard care for moderate/severe COVID-19 patients, related to 3 RCTs mentioned above, is presented in Table 3.19-1 below. No statistically significant difference was found in any outcome listed, in favour of colchicine.

Results: Therapeutics

Table 3.19-1: Summary of findings table on **colchicine compared to standard care** (3 RCT: Deftereos, Lopes, Salehzadeh) - https://covid-nma.com/living_data/index.php

Colchicine compared to Standard care or Placebo for Moderate/Severe/Critical COVID-19

Patient or population: Moderate/Severe/Critical COVID-19

Setting: Worldwide

Intervention: Colchicine

Comparison: Standard care or Placebo

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard care or Placebo	Risk with Colchicine				
Incidence viral negative conversion D7 - not measured	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement D7	632 per 1,000	640 per 1,000 (568 to 1,000)	RR 1.33 (0.90 to 1.98)	38 (1 RCT) ^b	⊕○○○ VERY LOW ^{c,d}	
Clinical improvement D24-D28	1,000 per 1,000	0 per 1,000 (0 to 0)	not estimable	38 (1 RCT) ^b	⊕○○○ VERY LOW ^{e,f}	
WHO progression score (level 6 or above) D7	158 per 1,000	106 per 1,000 (21 to 561)	RR 0.67 (0.13 to 3.55)	38 (1 RCT) ^b	⊕○○○ VERY LOW ^{c,d}	
WHO progression score (level 6 or above) D14-D28	96 per 1,000	13 per 1,000 (2 to 104)	RR 0.14 (0.02 to 1.08)	148 (2 RCT) ^h	⊕○○○ VERY LOW ^{g,i}	
WHO progression score (level 7 or above) D7	53 per 1,000	105 per 1,000 (11 to 1,000)	RR 2.00 (0.20 to 20.24)	38 (1 RCT) ^b	⊕○○○ VERY LOW ^{c,d}	
WHO progression score (level 7 or above) D14-D28	82 per 1,000	13 per 1,000 (2 to 106)	RR 0.16 (0.02 to 1.29)	148 (2 RCT) ^h	⊕○○○ VERY LOW ^{g,i}	
All-cause mortality D7	0 per 1,000	0 per 1,000 (0 to 0)	not estimable	38 (1 RCT) ^b	⊕○○○ VERY LOW ^{c,k}	
All-cause mortality D14-D28	33 per 1,000	8 per 1,000 (1 to 68)	RR 0.24 (0.03 to 2.09)	258 (3 RCT) ^l	⊕○○○ VERY LOW ^{g,i}	
Adverse events	421 per 1,000	526 per 1,000 (265 to 1,000)	RR 1.25 (0.63 to 2.46)	38 (1 RCT) ^b	⊕○○○ VERY LOW ^{c,m}	
Serious adverse events	27 per 1,000	27 per 1,000 (4 to 175)	RR 1.00 (0.16 to 6.38)	148 (2 RCT) ^h	⊕○○○ VERY LOW ^{g,i,n}	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
 CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Last update: November 10, 2020; b. Lopes MIF, 2020; c. Risk of bias downgraded by 1 level: some concerns regarding outcome measurement and selection of the reported result; d. Indirectness downgraded by 1 level: single study from a single country, therefore results in this population might not be generalizable to other settings; e. Imprecision downgraded by 1 level: due to few events; f. Imprecision downgraded by 2 levels: all participants had the event, no relative effect calculated; g. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of participants; h. Deftereos S, 2020; Lopes MIF, 2020; i. Risk of bias downgraded by 1 level: some concerns regarding deviation from intended intervention and outcome measurement; j. Risk of bias downgraded by 1 level: some concerns with deviation from intended interventions and selection of reported result; k. Imprecision downgraded by 2 levels: no events in both groups; l. Deftereos S, 2020; Lopes MIF, 2020; Salehzadeh F, 2020; m. We presume that the adverse event rates, and the corresponding relative risks, is similar across diverse settings; therefore not downgraded for indirectness

3.20 Nafamostat (Futhan©)

About the drug under consideration

Nafamostat mesilate (FUT-175, Futhan®, Nichi-Iko Pharmaceutical) is (with implications on coagulation, fibrinolysis, complement system, inflammatory cytokine release) and is quickly hydrolysed, the reason why it is typically administered as an intravenous drip. Nafamostat is not approved for any use by EMA or FDA.

Futhan®

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on nafamostat in ClinicalTrials.gov and EUdraCT registers.

keine abgeschlossenen, abgebrochenen Studien

Results of publications

Until now, no scientific publication on randomized clinical trials of nafamostat in Covid-19 patients could be identified.

keine veröffentlichten Studien

3.21 Gimsilumab

About the drug under consideration

Gimsilumab is a fully human monoclonal antibody that acts on granulocyte-macrophage colony-stimulating factor (GM-CSF) [1]; it is manufactured by Roivant Sciences Ltd. /Altasciences. Gimsilumab – ATC-code not assigned yet. Gimsilumab belongs to anti-inflammatories, antirheumatics, monoclonal antibodies drug class and has no approval for any indication by EMA or FDA yet.

monoklonaler Antikörper in Entwicklung

EMA/ FDA: keine Zulassung

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on gimsilumab in ClinicalTrials.gov and EUdraCT registers.

keine abgeschlossenen, abgebrochenen Studien

Results of publications

There are no published results from RCTs related to effectiveness and safety of gimsilumab for Covid-19 treatment; one Phase II study of gimsilumab is ongoing, estimated study completion date is March 2021 [155, 156].

keine veröffentlichten Studien

1 Phase 2 Studie läuft

3.22 Canakinumab

About the drug under consideration

Canakinumab is a human monoclonal anti-human interleukin-1 beta (IL-1 beta) antibody of the IgG1/κ isotype manufactured by Novartis Pharma AG. Canakinumab binds with high affinity specifically to human IL-1 beta and neutralises the biological activity of human IL-1 beta by blocking its

monoklonaler Antikörper

EMA Orphan Drug Zulassung für diverse Indikationen

interaction with IL-1 receptors, thereby preventing IL-1 beta-induced gene activation and the production of inflammatory mediators [157].

Canakinumab is not authorised in Covid-19 patients (EMA, FDA).

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on canakinumab in ClinicalTrials.gov and EUdRACT registers.

keine abgeschlossenen, abgebrochenen Studien

Results of publications

There are no published RCTs related to effectiveness and safety of canakinumab for Covid-19. Two studies of canakinumab are still ongoing: one Phase III study, estimated study completion date on December 2020 and one Phase II study, estimated completion date on December 2020 [158-160].

**keine veröffentlichten Studien
1 Phase 3 Studie läuft**

Manufacturer recently **announced preliminary interim results** from the CAN-COVID trial: the CAN-COVID trial failed to meet its primary endpoint showing that treatment with canakinumab plus standard of care (SoC) did not demonstrate a significantly greater chance of survival for patients without the need for invasive mechanical ventilation, compared with placebo plus SoC up to Day 29. The trial did not meet its key secondary endpoint of reducing the COVID-19-related death rate during the 4-week period after treatment. The safety profiles of canakinumab plus SoC and placebo plus SoC were comparable (<https://www.novartis.com/coronavirus/can-covid-clinical-trial>).

**CAN-COVID
negative Ergebnisse
kein Unterschied**

3.23 Lenzilumab

About the drug under consideration

Lenzilumab is a first-in-class Humaneered® recombinant monoclonal antibody targeting human GM-CSF, with potential immunomodulatory activity, high binding affinity in the picomolar range, 94% homology to human germline, and has low immunogenicity. Following intravenous administration, lenzilumab binds to and neutralizes GM-CSF, preventing GM-CSF binding to its receptor, thereby preventing GM-CSF-mediated signaling to myeloid progenitor cells. The inhibition of GM-CSF signaling may be beneficial in improving the hyperinflammation-related lung damage in the most severe cases of COVID-19. This blockade can be achieved through antagonism of the GM-CSF receptor or the direct binding of circulating GM-CSF [161, 162].

monoklonaler Antikörper

**für keine Indikation
bislang zugelassen**

**FDA: für
Einzelanwendungen im
Notfall – compassionate
use zur Verhinderung von
akutem Lungenversagen**

Lenzilumab is not authorised in Covid-19 patients (EMA, FDA). FDA has approved the administration of lenzilumab for COVID-19 patients under individual patient emergency IND applications to patients under the company's compassionate use program.

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on lenzilumab in ClinicalTrials.gov and EUdRACT registers.

Okt 2020: keine weiteren Studien

Results of publications

There are no published RCTs related to effectiveness and safety of lenzilumab for Covid-19.

A multicenter, phase 3, randomized, double-blinded, controlled, clinical trial with lenzilumab for the prevention of ARDS and/or death in hospitalized patients with pneumonia associated with coronavirus 2 (SARS-CoV-2) infection in COVID-19 patients is ongoing in US (NCT04351152). The primary objective of this study is to assess whether the use of lenzilumab in addition to current standard of care can alleviate the immune-mediated cytokine release syndrome (CRS) and reduce the time to recovery in 300 hospitalized patients with severe or critical COVID-19 pneumonia, with estimated completion date on September 2020 [33].

**Phase 3 RCT an
hospitalisierten Pts mit
Lungenentzündung
300 Pts.**

3.24 Vitamin D

About the drug under consideration

Vitamin D (ergocalciferol-D2, cholecalciferol-D3) is a fat-soluble vitamin increases the intestinal absorption of calcium and phosphate. Vitamin D is absorbed from the intestine and transported by protein binding in the blood to the liver (first hydroxylation to 25-hydroxycholecalciferol) and to the kidney (2nd hydroxylation to 1,25- dihydroxycholecalciferol, active metabolite responsible for increasing calcium absorption). It has been claimed as potentially protective against the infection since it may be associated with immunocompetence, inflammation, aging, and those diseases involved in determining the outcomes of COVID-19 [163]. VIOLET RCT (NCT03096314) of early high-dose enteral vitamin D3 supplementation in critically ill, vitamin D-deficient patients who were at high risk for death did not provide an advantage over placebo with respect to 90-day mortality or other, nonfatal outcomes among critically ill, vitamin D-deficient patients [164]. RCTs to assess efficacy and safety of vitamin D in COVID-19 patients are underway.

**protektive Wirkung gegen
Infekte bekannt**

**assoziiert mit guter
Immunantwort**

**VIOLET
RCT zu hoch-dosiertem Vit
D3 zur Supplementierung
kein Vorteil**

Vitamin D is not authorised in Covid-19 patients (EMA, FDA).

**mehrere klinische Studien
laufend**

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on Vitamin D in ClinicalTrials.gov and EUdraCT registers.

Results of publications

Entrenas Castillo et al. 2020 [165] published results from parallel pilot randomized open label, double-masked clinical trial on 76 consecutive patients hospitalized with COVID-19 infection in Spain (NCT04366908). Eligible patients were allocated at a 2 calcifediol:1 no calcifediol ratio, through electronic randomization on the day of admission to take oral calcifediol (0.532 mg), or not. Patients in the calcifediol treatment group continued with oral calcifediol (0.266 mg) on day 3 and 7, and then weekly until discharge or ICU admission. Of 50 patients treated with calcifediol, one required admission to the ICU (2%), while of 26 untreated patients, 13 required admission (50 %), $p < 0.001$. Calcifediol or 25-hydroxyvitamin D, a main metabolite of vitamin D, significantly reduced the need for ICU treatment of patients requiring hospitalization due to proven

**RCT
76 hospitalisierte Pts**

**Vorteil bei
Verhinderung von ICU
Verschlechterung der
Erkrankung**

COVID-19: Univariate Risk Estimate Odds Ratio for ICU in patients with Calcifediol treatment versus without Calcifediol treatment: 0.02 (95 %CI 0.002- 0.17). Multivariate Risk Estimate Odds Ratio for ICU in patients with Calcifediol treatment vs Without Calcifediol treatment ICU (adjusting by Hypertension and T2DM): 0.03 (95 %CI: 0.003-0.25). Of the patients treated with calcifediol, none died, and all were discharged, without complications. The 13 patients not treated with calcifediol, who were not admitted to the ICU, were discharged. Of the 13 patients admitted to the ICU, two died and the remaining 11 were discharged.

Rastogi et al. 2020 [166] published results from randomized, placebo-controlled trial (NCT04459247, SHADE) on 40 COVID-19 adult asymptomatic or mildly symptomatic SARS-CoV-2 RNA positive vitamin D deficient individuals (intervention (n=16) or control (n=24) group), with outcomes measure: Proportion of patients with SARSCoV-2 RNA negative before day-21 and change in inflammatory markers. 10 (62.5%) participants in the intervention group and 5 (20.8%) participants in the control arm (p<0.018) became SARS-CoV-2 RNA negative. Fibrinogen levels significantly decreased with cholecalciferol supplementation (intergroup difference 0.70 ng/ml; P=0.007) unlike other inflammatory biomarkers.

Murai et al. 2020 [167] presented as pre-print results from double-blind, randomised, placebo-controlled trial involving 240 hospitalised patients with severe COVID-19, in Brasil (NCT04449718). A single dose of 200,000 IU of vitamin D3 supplementation was safe and effective in increasing 25-hydroxyvitamin D levels, but did not significantly reduce hospital length of stay (hazard ratio, 1.12) or any other 10 clinically-relevant outcomes compared with placebo.

Summary of Finding table related to Vitamin D compared to Standard care/Placebo for mild/moderate/severe COVID-19 patients, related to 3 RCTs mentioned above, is presented in Table 3.24-1 below. No statistically significant difference was found in favour to Vitamin D in outcomes All-cause mortality D14-D28, AEs and SAEs. On outcome – WHO progression score (level 7 or above) D14-D28, statistically significant difference was found in favour to Vitamin D, RR 0.04 (0.01 to 0.29), absolute effect (95% CI) 480 fewer per 1.000 (from 495 fewer to 355 fewer), based on very low certainty of evidence.

RCT
40 Patient*innen
asymptomatisch oder
mild symptomatisch

Reduktion
Entzündungsmarker
Fibrinogen

RCT
240 hospitalisierte
Patient*innen
kein Unterschied bei
Dauer des
Krankenhausaufenthalts

SoF von 3 RCTs
kein Unterschied bei
Gesamtmortalität
ev. Verhinderung von
Verschlechterung

Results: Therapeutics

Table 3.24-1: Summary of findings table on **colchicine compared to standard care** (3 RCT:Entrenas Castillo, Rastogi, Murai) - https://covid-nma.com/living_data/index.php

Vitamin D compared to Standard care/Placebo for Mild/Moderate/Severe COVID-19

Patient or population: Mild/Moderate/Severe COVID-19

Setting: Worldwide

Intervention: Vitamin D

Comparison: Standard care/Placebo

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with Standard care/Placebo	Risk with Vitamin D				
Viral negative conversion D3 - not reported	-	-	-	-	-	outcome not yet measured or reported
Viral negative conversion D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
Clinical improvement D14-D28 - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO Progression Score (level 6 or above) D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO Progression Score (level 6 or above) D14-D28 - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO progression score (level 7 or above) D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
WHO progression score (level 7 or above) D14-D28	500 per 1,000	20 per 1,000 (5 to 145)	RR 0.04 (0.01 to 0.29)	76 (1 RCT) ^b	⊕⊕⊕⊕ VERY LOW ^{c,d,e}	
All-cause mortality D7 - not reported	-	-	-	-	-	outcome not yet measured or reported
All-cause mortality D14-D28	56 per 1,000	31 per 1,000 (3 to 325)	RR 0.56 (0.05 to 5.95)	313 (2 RCTe) ^f	⊕⊕⊕⊕ VERY LOW ^{g,h}	
Adverse events	0 per 1,000	0 per 1,000 (0 to 0)	RR 2.98 (0.12 to 72.30)	237 (1 RCT) ⁱ	⊕⊕⊕⊕ LOW ^j	
Serious adverse events - not reported	-	-	-	-	-	outcome not yet measured or reported

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).
CI: Confidence interval; RR: Risk ratio

GRADE Working Group grades of evidence: High certainty: We are very confident that the true effect lies close to that of the estimate of the effect; **Moderate certainty:** We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different; **Low certainty:** Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect; **Very low certainty:** We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations: a. Last updated: 06 December, 2020; b. Entrenas Castillo M, J Steroid Biochem Mo, 2020; c. Risk of bias downgraded by 1 level: some concerns regarding adequate randomization and deviations from intended interventions.; d. Indirectness downgraded by 1 level: results are from a single study from a single institution, therefore results in this population might not be generalizable to other settings.; e. Imprecision downgraded by 1 level: due to low number of events and participants.; f. Entrenas Castillo M, J Steroid Biochem Mo, 2020; Murai I, medRxiv, 2020; g. Inconsistency downgraded by 1 level: I²=58.9%; h. Imprecision downgraded by 2 levels: due to very wide confidence interval consistent with the possibility for benefit and the possibility for harm and low number of events and participants.; i. Murai I, medRxiv, 2020; j. We presume that the adverse event rates and the corresponding relative risks, are similar across diverse settings; therefore not downgraded for indirectness.

3.25 Baricitinib

About the drug under consideration

Baricitinib is a selective and reversible inhibitor of Janus kinase (JAK)1 and JAK2. Janus kinases (JAKs) are enzymes that transduce intracellular signals from cell surface receptors for a number of cytokines and growth factors involved in haematopoiesis, inflammation and immune function. Baricitinib (Olumiant) is indicated in EU for the treatment of moderate to severe active rheumatoid arthritis in adult patients who have responded inadequately to, or who are intolerant to one or more disease-modifying anti-rheumatic drugs and for the treatment of moderate to severe atopic dermatitis in adult patients who are candidates for systemic therapy [168, 169].

Baricitinib (Olumiant) has not been approved by the European Medicines Agency (EMA) or the American Food and Drug Administration (FDA) for COVID-19.

On November 19, 2020, the U.S. Food and Drug Administration (FDA) issued an Emergency Use Authorization (EUA) for the distribution and emergency use of baricitinib to be used in combination with remdesivir in hospitalised adult and pediatric patients two years of age or older with suspected or laboratory confirmed COVID-19 who require supplemental oxygen, invasive mechanical ventilation, or extracorporeal membrane oxygenation (ECMO) [170].

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on baricitinib in ClinicalTrials.gov and EUdraCT registers.

Results of publications

On December 11, 2020, **Kalil et al.** [171] published results from the Adaptive COVID-19 Treatment Trial (ACTT-2) (NCT04401579), multicentre, double-blind, randomized, placebo-controlled trial evaluating baricitinib plus remdesivir with remdesivir alone in hospitalised adults with Covid-19 in eight countries. Patients treated with baricitinib in combination with remdesivir had a significant reduction in median time to recovery from 8 to 7 days compared to remdesivir. Patients receiving high-flow oxygen or noninvasive ventilation at enrollment had a time to recovery of 10 days with combination treatment and 18 days with remdesivir alone (rate ratio for recovery, 1.51; 95% CI, 1.10 to 2.08). Patients treated with baricitinib in combination with remdesivir were more likely to have a better clinical status at day 15 compared to patients treated with remdesivir. Patients with a baseline ordinal score of 6 who received combination treatment were most likely to have clinical improvement at day 15 (odds ratio, 2.2; 95% CI, 1.4 to 3.6). The proportion of patients who died by Day 29 was not statistically significant different between groups: the 28-day mortality was 5.1% in the combination group and 7.8% in the remdesivir group (hazard ratio for death, 0.65; 95% CI, 0.39 to 1.09). The incidence of new use of oxygen was statistically significant lower in the combination group than in the remdesivir group (22.9% vs. 40.3%; difference, -17.4 percentage points; 95% CI, -31.6 to -2.1), as was the incidence of new use of mechanical ventilation or ECMO (10.0% vs. 15.2%; difference, -5.2 percentage points; 95% CI, -9.5 to -0.9). The incidence of progression to death

Januskinase-Inhibitor

Baricitinib (Olumiant) in EU für moderate bis schwere rheumatoide Arthritis zugelassen

weder von EMA noch FDA für covid-19 zugelassen

Nov 2020: Notzulassung durch FDA in Kombination mit Remdesivir hospitalisierte Patient*innen mit bedarf zur Beatmung

ACTT-2 Baricitinib + Remdesivir vs. Remdesivir

Verkürzung der Erkrankung unter Kombinationstherapie

besser in Verhinderung von Fortschreiten der Erkrankung und bei Bedarf nach Beatmung

kein Unterschied bei Mortalität

Results: Therapeutics

or noninvasive or invasive ventilation was lower in the combination group than in the remdesivir group (22.5% vs. 28.4%; rate ratio, 0.77; 95% CI, 0.60 to 0.98), as was the incidence of progression to death or invasive ventilation (12.2% vs. 17.2%; rate ratio, 0.69; 95% CI, 0.50 to 0.95).

The most common grade 3 or 4 adverse events occurring in at least 5% of all patients were hyperglycemia, anemia, decreased lymphocyte count, and acute kidney injury. The incidence of these adverse events was similar in the two treatment groups. Serious adverse events were statistically significant less frequent in the combination group than in the remdesivir group (16.0% vs. 21.0%; difference, -5.0 percentage points; 95% CI, -9.8 to -0.3; p=0.03), as were new infections (5.9% vs. 11.2%; difference, -5.3 percentage points; 95% CI, -8.7 to -1.9; p=0.003).

5% der Patient*innen haben ≥ 3 Nebenwirkungen – in IG und KG

weniger SAE unter Kombinationstherapie

3.26 Molnupiravir

About the drug under consideration

Molnupiravir is the orally available pro-drug of the nucleoside analogue N4-hydroxycytidine (NHC), which has shown potent anti-influenza virus activity in mice, guinea pigs, ferrets and human airway epithelium organoids. Animal study in ferrets showed that therapeutic treatment of infected animals with molnupiravir (MK-4482/EIDD-2801) twice a day significantly reduced the SARS-CoV-2 load in the upper respiratory tract and completely suppressed spread to untreated contact animals [172, 173].

antivirales Medikament ähnlich Remdesivir aber orale Verabreichung

Molnupiravir attacks the same viral enzyme as Gilead's Remdesivir, but it can be taken orally. This would allow an administration at home and, therefore, earlier in the course of the disease. According to Ridgeback Biotherapeutics, molnupiravir has an extremely high barrier to resistance. According to Merck Sharp & Dohme/ MSD, molnupiravir is aimed at the treatment of Covid-19 in patients hospitalised due to mild, moderate or severe disease, and non-hospitalized patients with mild or moderate disease. [173].

frühere Verabreichung zu Hause daher möglich

hospitalisierte, aber auch milde und moderate Erkrankung

Molnupiravir is not approved by the European Medicines Agency (EMA) or the American Food and Drug Administration (FDA). [173].

weder von EMA noch FDA zugelassen

Withdrawn, suspended or terminated studies

No withdrawn, suspended or terminated interventional studies were found on molnupiravir in ClinicalTrials.gov and EUdraCT registers.

Results of publications

There are no published RCTs related to effectiveness and safety of molnupiravir for Covid-19. It is currently investigated in phase 1/2, 2 and 2/3 clinical trials (NCT04405570, NCT04405739, NCT04575584, NCT04575597, ISRCTN27106947), in hospitalised and non-hospitalised adults with COVID-19.

keine RCTs derzeit in Phase 1/2, 2 und 1/3 Studien mit verschiedenen Pts. Populationen

References

- [1] Pang J., Wang M. X., Ang I. Y. H., Tan S. H. X., Lewis R. F., Chen J. I., et al. Potential Rapid Diagnostics, Vaccine and Therapeutics for 2019 Novel Coronavirus (2019-nCoV): A Systematic Review. *J Clin Med*. 2020;9(3). Epub 2020/03/01. DOI: 10.3390/jcm9030623
- [2] Martin R., Löchel H., Welzel M., Hattab G., Hauschild A. and Heider D. CORDITE: The Curated CORona Drug InTERactions Database for SARS-CoV-2. *iScience*. 2020;23(7):101297-101297. DOI: 10.1016/j.isci.2020.101297.
- [3] Boutron I. and al. e. Interventions for preventing and treating COVID-19: protocol for a living mapping of research and a living systematic review Zenodo. 2020;April 8(<http://doi.org/10.5281/zenodo.3744600>).
- [4] Thorlund K., Dron L., Park J., Hsu G., Forrest J. and Mills E. A real-time dashboard of clinical trials for COVID-19. *Lancet*. 2020;April 24 (DOI:[https://doi.org/10.1016/S2589-7500\(20\)30086-8](https://doi.org/10.1016/S2589-7500(20)30086-8)).
- [5] Chen Q., Allot A. and Lu Z. Keep up with the latest coronavirus research. *Nature Communications*. 2020;579(7798):193.
- [6] Jackson L., Anderson E., Roupheal N., Roberts P., Makhene M., Coler R., et al. An mRNA Vaccine against SARS-CoV-2 — Preliminary Report. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2022483.
- [7] Anderson E., Roupheal N., Widge A., Jackson L., Roberts P., Makhene M., et al. Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2028436.
- [8] Zhu F., Li Y., Guan X., Hou L., Wang W., Li J., et al. Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial. *The Lancet*. 2020;395(10240):1845-1854. DOI: 10.1016/S0140-6736(20)31208-3.
- [9] Zhu F., Guan X., Li Y., Huang J., Jiang T., Hou L., et al. Immunogenicity and safety of a recombinant adenovirus type-5-vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised, double-blind, placebo-controlled, phase 2 trial. *The Lancet*. 2020;396(10249):479-488. DOI: 10.1016/S0140-6736(20)31605-6.
- [10] Keech C., Albert G., Cho I., Robertson A., Reed P., Neal S., et al. Phase 1–2 Trial of a SARS-CoV-2 Recombinant Spike Protein Nanoparticle Vaccine. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2026920.
- [11] Folegatti P., Ewer K., Aley P., Angus B., Becker S., Belij-Rammerstorfer S., et al. Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial. *The Lancet*. 2020;396(10249):467-478. DOI: 10.1016/S0140-6736(20)31604-4.
- [12] Voysey M., Clemens S., Madhi S., Weckx L., Folegatti P., Aley P., et al. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *The Lancet*. DOI: 10.1016/S0140-6736(20)32661-1.

References

- [13] Ramasamy M., Minassian A., Ewer K., Flaxman A., Folegatti P., Owens D., et al. Safety and immunogenicity of ChAdOx1 nCoV-19 vaccine administered in a prime-boost regimen in young and old adults (COV002): a single-blind, randomised, controlled, phase 2/3 trial. *The Lancet*. DOI: 10.1016/S0140-6736(20)32466-1.
- [14] Logunov D., Dolzikhova I., Zubkova O., Tukhvatullin A., Shcheblyakov D., Dzharullaeva A., et al. Safety and immunogenicity of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine in two formulations: two open, non-randomised phase 1/2 studies from Russia. *The Lancet*. 2020. DOI: 10.1016/S0140-6736(20)31866-3.
- [15] Mulligan M., Lyke K., Kitchin N., Absalon J., Gurtman A., Lockhart S., et al. Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. *Nature*. 2020. DOI: 10.1038/s41586-020-2639-4.
- [16] Sahin U., Muik A., Derhovanessian E., Vogler I., Kranz L., Vormehr M., et al. COVID-19 vaccine BNT162b1 elicits human antibody and TH1 T-cell responses. *Nature*. 2020. DOI: 10.1038/s41586-020-2814-7.
- [17] Polack F., Thomas S., Kitchin N., Absalon J., Gurtman A., Lockhart S., et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2034577.
- [18] Xia S., Duan K., Zhang Y., Zhao D., Zhang H., Xie Z., et al. Effect of an Inactivated Vaccine Against SARS-CoV-2 on Safety and Immunogenicity Outcomes: Interim Analysis of 2 Randomized Clinical Trials. *JAMA*. 2020. DOI: 10.1001/jama.2020.15543.
- [19] Mulligan M. An Inactivated Virus Candidate Vaccine to Prevent COVID-19. *JAMA*. 2020. DOI: 10.1001/jama.2020.15539.
- [20] Xia S., Zhang Y., Wang Y., Wang H., Yang Y., Gao G., et al. Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *The Lancet Infectious Diseases*. 2020. DOI: 10.1016/S1473-3099(20)30831-8.
- [21] Zhang Y., Zeng G., Pan H., Li C., Hu Y., Chu K., et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18–59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. *The Lancet Infectious Diseases*. DOI: 10.1016/S1473-3099(20)30843-4.
- [22] Phillips N., Cyranoski D. and Mallapaty S. A leading coronavirus vaccine trial is on hold: scientists react. *Nature*. 2020. Epub 2020/09/11. DOI: 10.1038/d41586-020-02594-w.
- [23] European Medicines Agency (EMA). International regulators align positions on phase 3 COVID-19 vaccine trials.: 09/07/2020. Available from: <https://www.ema.europa.eu/en/news/international-regulators-align-positions-phase-3-covid-19-vaccine-trials>.
- [24] European Medicines Agency (EMA). EMA starts second rolling review of a COVID-19 vaccine. 2020 [cited 06/10/2020]. Available from: <https://www.ema.europa.eu/en/news/ema-starts-second-rolling-review-covid-19-vaccine>.
- [25] European Medicines Agency (EMA). EMA receives application for conditional marketing authorisation of COVID-19 mRNA vaccine BTN162b2. 2020 [cited 01/12/2020]. Available from: <https://www.ema.europa.eu/en/news/ema-receives-application-conditional-marketing-authorisation-covid-19-mrna-vaccine-bnt162b2>.

References

- [26] European Medicines Agency (EMA). EMA receives application for conditional marketing authorisation of Moderna COVID-19 vaccine. 2020 [cited 01/12/2020]. Available from: <https://www.ema.europa.eu/en/news/ema-receives-application-conditional-marketing-authorisation-moderna-covid-19-vaccine>.
- [27] European Medicines Agency (EMA). EMA starts rolling review of Janssen's COVID-19 vaccine Ad26.COV2.S. 2020 [cited 01/12/2020]. Available from: <https://www.ema.europa.eu/en/news/ema-starts-rolling-review-janssens-covid-19-vaccine-ad26cov2s>.
- [28] Jackson L. A. Safety and Immunogenicity Study of 2019-nCoV Vaccine (mRNA-1273) for Prophylaxis SARS CoV-2 Infection. 2020 [cited 07.04.]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04283461>. Jackson.
- [29] National Institute of Health (NIH). NIH clinical trial of investigational vaccine for COVID-19 begins. 2020 [cited 07.04.]. Available from: <https://www.nih.gov/news-events/news-releases/nih-clinical-trial-investigational-vaccine-covid-19-begins>.
- [30] World Health Organisation (WHO). DRAFT landscape of COVID-19 candidate vaccines –20 March 2020. 2020 [cited 31.03.2020]. Available from: <https://www.who.int/blueprint/priority-diseases/key-action/novel-coronavirus-landscape-ncov-21march2020.PDF?ua=1>.
- [31] BioWorld. China approves first homegrown COVID-19 vaccine to enter clinical trials. [cited 02.03.2020]. Available from: <https://www.bioworld.com/articles/433791-china-approves-first-homegrown-covid-19-vaccine-to-enter-clinical-trials>.
- [32] U.S. National Library of Medicine. A Phase I Clinical Trial in 18-60 Adults (APICTH). 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04313127>.
- [33] U.S. National Library of Medicine. ClinicalTrials.gov. Available from: <https://clinicaltrials.gov/>.
- [34] University of Oxford. Oxford team to begin novel coronavirus vaccine research. 2020 [cited 03.04.2020]. Available from: <http://www.ox.ac.uk/news/2020-02-07-oxford-team-begin-novel-coronavirus-vaccine-research>.
- [35] Denis M., Vanderweerd V., Verbeke R. and Van der Vliet D. Overview of information available to support the development of medical countermeasures and interventions against COVID-19. Living document. 2020 [cited 03.03.2020]. Available from: https://rega.kuleuven.be/if/pdf_corona.
- [36] U.S. National Library of Medicine. A Study of a Candidate COVID-19 Vaccine (COV001). 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04324606>.
- [37] Mahase E. Covid-19: Oxford team begins vaccine trials in Brazil and South Africa to determine efficacy. BMJ. 2020;369:m2612. DOI: 10.1136/bmj.m2612.
- [38] ISRCTNregistry. ISRCTN89951424. A phase III study to investigate a vaccine against COVID-19. 2020 [cited 13/07/2020]. Available from: <https://doi.org/10.1186/ISRCTN89951424>.
- [39] Hodgson J. The pandemic pipeline. 2020 [cited 03.04.]. Available from: <https://www.nature.com/articles/d41587-020-00005-z>.
- [40] FierceBiotech. Pfizer, BioNTech strike COVID-19 deal, commit multiple R&D sites to vaccine development. 2020 [cited 07.04.]. Available from: <https://www.fiercebiotech.com/biotech/pfizer-biontech-strike-covid-19-deal-commit-multiple-r-d-sites-to-vaccine-development>.

References

- [41] Walsh E., Frenck R., Falsey A., Kitchin N., Absalon J., Gurtman A., et al. RNA-Based COVID-19 Vaccine BNT162b2 Selected for a Pivotal Efficacy Study. medRxiv. 2020:2020.2008.2017.20176651. DOI: 10.1101/2020.08.17.20176651.
- [42] Krammer F. SARS-CoV-2 vaccines in development. Nature. 2020. DOI: 10.1038/s41586-020-2798-3.
- [43] Sinovac. Sinovac Announces Positive Preliminary Results of Phase I/II Clinical Trials for Inactivated Vaccine Candidate Against COVID-19. 2020. Available from: http://www.sinovac.com/?optionid=754&auto_id=904.
- [44] Isakova-Sivak I. and Rudenko L. A promising inactivated whole-virion SARS-CoV-2 vaccine. The Lancet Infectious Diseases. 2020. DOI: 10.1016/S1473-3099(20)30832-X.
- [45] Sadoff J., Le Gars M., Shukarev G., Heerwegh D., Truyers C., de Groot A., et al. Safety and immunogenicity of the Ad26.COV2.S COVID-19 vaccine candidate: interim results of a phase 1/2a, double-blind, randomized, placebo-controlled trial. medRxiv. 2020:2020.2009.2023.20199604. DOI: 10.1101/2020.09.23.20199604.
- [46] Novavax. Novavax Awarded Funding from CEPI for COVID-19 Vaccine Development. 2020 [cited 06.04.]. Available from: <https://ir.novavax.com/news-releases/news-release-details/novavax-awarded-funding-cepi-covid-19-vaccine-development>.
- [47] Nature. A surprising player in the race for a SARS-CoV-2 vaccine. 2020 [cited 06.04.]. Available from: <https://www.nature.com/articles/d42473-020-00032-z>.
- [48] Drug Development and Delivery. Novavax Advances Development of Novel COVID-19 Vaccine. 2020 [cited 06.04.]. Available from: <https://drug-dev.com/novavax-advances-development-of-novel-covid-19-vaccine/>.
- [49] Novavax. MATRIX-M™ ADJUVANT TECHNOLOGY. 2020 [cited 06.04.]. Available from: <https://novavax.com/page/10/matrix-m-adjuvant-technology.html>.
- [50] Mahase E. Covid-19: What do we know so far about vaccine? . BMJ. 2020;369:m1679.
- [51] Callaway E. The race for coronavirus vaccines. Nature. 2020;580(April 30).
- [52] Le T. and al. e. The COVID-19 vaccine development landscape. 2020. Available from: <https://www.nature.com/articles/d41573-020-00073-5>.
- [53] European Medicines Agency (EMA). Update on remdesivir. Meeting highlights from the Committee for Medicinal Products for Human Use (CHMP) 25-28 May 2020.: 2020 [cited 14/06/2020]. Available from: <https://www.ema.europa.eu/en/news/meeting-highlights-committee-medicinal-products-human-use-chmp-25-28-may-2020> <https://www.ema.europa.eu/en/human-regulatory/overview/public-health-threats/coronavirus-disease-covid-19/treatments-vaccines-covid-19#remdesivir-section>
- [54] European Medicines Agency (EMA). Summary of opinion (initial authorisation. Veklury (remdesivir). 25/06/2020. Available from: https://www.ema.europa.eu/en/documents/smop-initial/chmp-summary-positive-opinion-veklury_en.pdf.
- [55] Beigel J., Tomashek K., Dodd L., Mehta A., Zingman B., Kalil A., et al. Remdesivir for the Treatment of Covid-19 — Preliminary Report. New England Journal of Medicine. 2020. DOI: 10.1056/NEJMoa2007764.

References

- [56] The European public assessment report (EPAR). Veklury: Product information. 2020 [cited 06/07/2020]. Available from: https://www.ema.europa.eu/en/documents/product-information/veklury-epar-product-information_en.pdf.
- [57] European Medicines Agency (EMA). Meeting highlights from the Pharmacovigilance Risk Assessment Committee (PRAC) 28 September - 1 October 2020. 2020 [cited 02/10/2020]. Available from: <https://www.ema.europa.eu/en/news/meeting-highlights-pharmacovigilance-risk-assessment-committee-prac-28-september-1-october-2020>.
- [58] Federal Drug Administration (FDA). Fact sheet for healthcare providers: Emergency use authorization (EUA) of Baricitinib. 2020. Available from: <https://www.fda.gov/media/143823/download>.
- [59] Rochwerg B., Agoritsas T., Lamontagne F., Leo Y., Macdonald H., Agarwal A., et al. A living WHO guideline on drugs for covid-19. *BMJ*. 2020;370:m3379. DOI: 10.1136/bmj.m3379.
- [60] World Health Organisation (WHO). Repurposed Antiviral Drugs for Covid-19 — Interim WHO Solidarity Trial Results. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2023184.
- [61] National Institutes of Health (NIH). COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. 2020 [cited 13/07/2020]. Available from: <https://www.covid19treatmentguidelines.nih.gov/>.
- [62] Wang Y., Zhang D., Du G. and al. e. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet*. 2020;published online April 29([https://doi.org/10.1016/S0140-6736\(20\)31022-9](https://doi.org/10.1016/S0140-6736(20)31022-9)).
- [63] Beigel J., Tomashek K., Dodd L., Mehta A., Zingman B., Kalil A., et al. Remdesivir for the Treatment of Covid-19 — Final Report. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2007764.
- [64] Goldman D., Lye D. C., Hui D., Marks K., Bruno R., Montejano R., et al. Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2015301.
- [65] Spinner C., Gottlieb R., Criner G., Arribas López J., Cattelan A., Soriano Viladomiu A., et al. Effect of Remdesivir vs Standard Care on Clinical Status at 11 Days in Patients With Moderate COVID-19: A Randomized Clinical Trial. *JAMA*. 2020. Epub 2020/08/22. DOI: 10.1001/jama.2020.16349.
- [66] Pan H., Peto R., Karim Q., Alejandria M., Henao-Restrepo A., García C., et al. Repurposed antiviral drugs for COVID-19 –interim WHO SOLIDARITY trial results. *medRxiv*. 2020:2020.2010.2015.20209817. DOI: 10.1101/2020.10.15.20209817.
- [67] European network for health technology assessment (EUnetHTA). PTRCR15 Authoring Team. Remdesivir for the treatment of hospitalised patients with COVID-19 - Update 1. Rapid Collaborative. Diemen (The Netherlands): 2020. Available from: <https://eunetha.eu/wp-content/uploads/2020/12/PTRCR15-update-1-December-2020-v1.0.pdf>.
- [68] Wang Y., Zhang D., Du G., Du R., Zhao J., Jin Y., et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet*. 2020;395(10236):1569-1578. Epub 2020/05/20. DOI: 10.1016/s0140-6736(20)31022-9.

References

- [69] Should favipiravir be used for COVID-19? Ministry of Health Singapore and Agency for Care Effectiveness: 2020 [cited 06/04/2020]. Available from: [https://www.moh.gov.sg/docs/librariesprovider5/clinical-evidence-summaries/favipiravir-for-covid-19-\(26-march-2020\).pdf](https://www.moh.gov.sg/docs/librariesprovider5/clinical-evidence-summaries/favipiravir-for-covid-19-(26-march-2020).pdf).
- [70] Dong L., Hu S. and Gao J. Discovering drugs to treat coronavirus disease 2019 (COVID-19). *Drug Discoveries & Therapeutics*. 2020;14(1):58-60. DOI: 10.5582/ddt.2020.01012.
- [71] Chen C., Huang J., Cheng Z., Wu J., Chen S., Zhang Y., et al. Favipiravir versus Arbidol for COVID-19: A Randomized Clinical Trial. 2020.
- [72] Lou Y., Liu L. and Qiu Y. Clinical Outcomes and Plasma Concentrations of Baloxavir Marboxil and Favipiravir in COVID-19 Patients: an Exploratory Randomized, Controlled Trial. *medRxiv*. 2020:2020.2004.2029.20085761. DOI: 10.1101/2020.04.29.20085761.
- [73] Ivashchenko A., Dmitriev K., Vostokova N., Azarova V., Blinow A., Egorova A., et al. AVIFAVIR for Treatment of Patients with Moderate COVID-19: Interim Results of a Phase II/III Multicenter Randomized Clinical Trial. *Clinical Infectious Diseases*. 2020. DOI: 10.1093/cid/ciaa1176.
- [74] Dabbous H., El-Sayed M., El Assal G., Elghazaly H., Ebeid F. F., Sherief A., et al. Research Square. 2020. DOI: 10.21203/rs.3.rs-83677/v1.
- [75] Balykova L., Granovskaya M., Zaslavskaya K., Simakina E., Agafina A., Ivanova A., et al. New possibilities for targeted antiviral therapy for COVID-19. Results of a multi center clinical study of the efficacy and safety of using the drug Areplivir. *Infectious diseases: News, Opinions, Training*. 2020;9:16-29. DOI: 10.33029/2305-3496-2020-9-3-16-29.
- [76] Ruzhentsova T., Chukhliaev P., Khavkina D., Garbuzov A., Oseshnyuk R. and Soluyanova T. Phase 3 Trial of Coronavir (Favipiravir) in Patients with Mild to Moderate COVID-19. *The Lancet*. 2020;(Preprint). DOI: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3696907.
- [77] Udwardia Z., Singh P., Barkate H., Patil S., Rangwala S., Pendse A., et al. Safety of Favipiravir, an Oral RNA-Dependent RNA Polymerase Inhibitor, in Mild-to-Moderate COVID-19: A Randomized, Comparative, Open-Label, Multicenter, Phase 3 Clinical TEfficacy and safety of favipiravir, an oral RNA-dependent RNA polymerase inhibitor, in mild-to-moderate COVID-19: A randomized, comparative, open-label, multicenter, phase 3 clinical trial. *International Journal of Infectious Diseases*. DOI: 10.1016/j.ijid.2020.11.142.
- [78] Doi Y., Hibino M., Hase R., Yamamoto M., Kasamatsu Y., Hirose M., et al. A prospective, randomized, open-label trial of early versus late favipiravir in hospitalized patients with COVID-19. *Antimicrobial Agents and Chemotherapy*. 2020:AAC.01897-01820. DOI: 10.1128/aac.01897-20.
- [79] Zhao H., Zhu Q., Zhang C., Li J., Wei M., Qin Y., et al. Tocilizumab combined with favipiravir in the treatment of COVID-19: A multicenter trial in a small sample size. *Biomedicine & Pharmacotherapy*. 2020:110825. DOI: <https://doi.org/10.1016/j.biopha.2020.110825>.
- [80] McKeage K., Perry C. M. and Keam S. J. Darunavir: a review of its use in the management of HIV infection in adults. *Drugs*. 2009;69(4):477-503.
- [81] Chen J., Xia L., Liu L., Xu Q., Ling Y., Huang D., et al. Antiviral Activity and Safety of Darunavir/Cobicistat for the Treatment of COVID-19. *Open Forum Infectious Diseases*. 2020;7(7). DOI: 10.1093/ofid/ofaa241.

References

- [82] Fujii S. and Hitomi Y. New synthetic inhibitors of Clr, C1 esterase, thrombin, plasmin, kallikrein and trypsin. *Biochim Biophys Acta*. 1981;661(2):342-345. Epub 1981/10/13. DOI: 10.1016/0005-2744(81)90023-1.
- [83] Hoffmann M., Kleine-Weber H., Schroeder S., Kruger N., Herrler T., Erichsen S., et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. 2020;181. Epub 2020/03/07. DOI: 10.1016/j.cell.2020.02.052.
- [84] Kawase M., Shirato K., van der Hoek L., Taguchi F. and Matsuyama S. Simultaneous treatment of human bronchial epithelial cells with serine and cysteine protease inhibitors prevents severe acute respiratory syndrome coronavirus entry. *J Virol*. 2012;86(12):6537-6545. Epub 2012/04/13. DOI: 10.1128/JVI.00094-12.
- [85] Zhou Y., Vedantham P., Lu K., Agudelo J., Carrion R., Jr., Nunneley J. W., et al. Protease inhibitors targeting coronavirus and filovirus entry. *Antiviral Res*. 2015;116:76-84. Epub 2015/02/11. DOI: 10.1016/j.antiviral.2015.01.011.
- [86] European network for health technology assessment (EUnetHTA). EUnetHTA Rolling Collaborative Review (RCR04). Authoring Team. Camostat for the treatment of Covid-19. Diemen (The Netherlands). 2020 [cited 15/08/2020]. Available from: https://eunetha.eu/wp-content/uploads/2020/08/EUnetHTA-Covid-19_RCR04_Camostat-v1.0.pdf.
- [87] Apeiron Biologics. APN01. 2020 [cited 07.04.2020]. Available from: <https://www.apeiron-biologics.com/project-overview/#APN01>.
- [88] Kuba K., Imai Y., Rao S., Gao H., Guo F., Guan B., et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nature medicine*. 2005;11(8):875-879. Epub 2005/07/10. DOI: 10.1038/nm1267.
- [89] Monteil V., Hyesoo Kwon, Patricia Prado, Astrid Hagelkrüys, Reiner A. Wimmer, Martin Stahl, et al. Inhibition of SARS-CoV-2 infections in engineered human tissues using clinical-grade soluble human ACE2. 2020 [cited 07.04.2020]. Available from: https://www.cell.com/pb-assets/products/coronavirus/CELL_CELL-D-20-00739.pdf.
- [90] European network for health technology assessment (EUnetHTA). EUnetHTA Rolling Collaborative Review (RCR09). Authoring Team. APN01 for the Treatment of Covid19. Diemen (The Netherlands). 2020 [cited 15/08/2020]. Available from: https://eunetha.eu/wp-content/uploads/2020/08/EUnetHTA-Covid-19_RCR09_APN01_August2020_FINAL.pdf.
- [91] European Medicines Agency. RoActemra (tocilizumab). Amsterdam: 2020. Available from: <https://www.ema.europa.eu/en/medicines/human/EPAR/roactemra>.
- [92] Rosas I., Bräu N., Waters M., Go R., Hunter B., Bhagani S., et al. Tocilizumab in Hospitalized Patients With COVID-19 Pneumonia. *medRxiv*. 2020:2020.2008.2027.20183442. DOI: 10.1101/2020.08.27.20183442.
- [93] Wang D., Fu B., Peng Z., Yang D. and Han M. Tocilizumab Ameliorates the Hypoxia in COVID-19 Moderate Patients with Bilateral Pulmonary Lesions: A Randomized, Controlled, Open-Label, Multicenter Trial. Preprints with The Lancet. 2020.
- [94] Salama C., Han J., Yau L., Reiss W., Kramer B., Neidhart J., et al. Tocilizumab in nonventilated patients hospitalized with Covid-19 pneumonia. *medRxiv*. 2020:2020.2010.2021.20210203. DOI: 10.1101/2020.10.21.20210203.

References

- [95] Hermine O., Mariette X., Tharaux P., Resche-Rigon M., Porcher R., Ravaud P., et al. Effect of Tocilizumab vs Usual Care in Adults Hospitalized With COVID-19 and Moderate or Severe Pneumonia: A Randomized Clinical Trial. *JAMA Internal Medicine*. 2020. DOI: 10.1001/jamainternmed.2020.6820.
- [96] Salvarani C., Dolci G., Massari M., Merlo D., Cavuto S., Savoldi L., et al. Effect of Tocilizumab vs Standard Care on Clinical Worsening in Patients Hospitalized With COVID-19 Pneumonia: A Randomized Clinical Trial. *JAMA Internal Medicine*. 2020. DOI: 10.1001/jamainternmed.2020.6615.
- [97] Stone J., Frigault M., Serling-Boyd N., Fernandes A., Harvey L., Foulkes A., et al. Efficacy of Tocilizumab in Patients Hospitalized with Covid-19. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2028836.
- [98] Furlow B. COVACTA trial raises questions about tocilizumab's benefit in COVID-19. *The Lancet Rheumatology*. DOI: 10.1016/S2665-9913(20)30313-1.
- [99] European Medicines Agency (EMA). EPAR summary for the public: Kevzara (sarilumab). 2017. Available from: <https://www.ema.europa.eu/en/medicines/human/EPAR/kevzara>.
- [100] Institut national d'excellence en santé et en services sociaux (INESSS). COVID-19 et interférons. Québec, Qc: 2020. Available from: https://www.inesss.qc.ca/fileadmin/doc/INESSS/COVID-19/COVID-19_interferons.pdf.
- [101] The European public assessment report (EPAR). Betaferon – Product information. Available from: https://www.ema.europa.eu/documents/product-information/betaferon-epar-product-information_en.pdf.
- [102] The European public assessment report (EPAR). Extavia – Product information. Available from: https://www.ema.europa.eu/documents/product-information/extavia-epar-product-information_en.pdf.
- [103] Hung I., Lung K., Tso E., Liu R., Chung T., Chu M., et al. Triple combination of interferon beta-1b, lopinavir–ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *The Lancet*. 2020. DOI: 10.1016/S0140-6736(20)31042-4.
- [104] Huang Y., Tang S., Xu X., Zeng Y., He X., Li Y., et al. No Statistically Apparent Difference in Antiviral Effectiveness Observed Among Ribavirin Plus Interferon-Alpha, Lopinavir/Ritonavir Plus Interferon-Alpha, and Ribavirin Plus Lopinavir/Ritonavir Plus Interferon-Alpha in Patients With Mild to Moderate Coronavirus Disease 2019: Results of a Randomized, Open-Labeled Prospective Study. *Frontiers in Pharmacology*. 2020;11(1071). DOI: 10.3389/fphar.2020.01071.
- [105] Esquivel-Moynelo I., Perez-Escribano J., Duncan-Robert Y., Vazquez-Blonquist D., Bequet-Romero M., Baez-Rodriguez L., et al. Effect and safety of combination of interferon alpha-2b and gamma or interferon alpha-2b for negativization of SARS-CoV-2 viral RNA. Preliminary results of a randomized controlled clinical trial. *medRxiv*. 2020:2020.2007.2029.20164251. DOI: 10.1101/2020.07.29.20164251.
- [106] Monk P., Marsden R., Tear V., Brookes J., Batten T., Mankowski M., et al. Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001) for treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase 2 trial. *The Lancet Respiratory Medicine*. DOI: 10.1016/S2213-2600(20)30511-7.

References

- [107] Davoudi-Monfared E., Rahmani H., Khalili H., Hajiabdolbaghi M., Salehi M., Abbasian L., et al. A Randomized Clinical Trial of the Efficacy and Safety of Interferon β -1a in Treatment of Severe COVID-19. *Antimicrob Agents Chemother.* 2020;64(9). Epub 2020/07/15. DOI: 10.1128/aac.01061-20.
- [108] Rahmani H., Davoudi-Monfared E., Nourian A., Khalili H., Hajizadeh N., Jalalabadi N., et al. Interferon β -1b in treatment of severe COVID-19: A randomized clinical trial. *International Immunopharmacology.* 2020;88:106903. DOI: <https://doi.org/10.1016/j.intimp.2020.106903>.
- [109] Casadevall A. and Pirofski L. The convalescent sera option for containing COVID-19. *J Clin Invest.* 2020;Mar 13(pii: 138003. doi: 10.1172/JCI138003. [Epub ahead of print]).
- [110] Roback J. and Guarner J. Convalescent Plasma to Treat COVID-19 Possibilities and Challenges. *JAMA.* 2020;Mar 27(doi: 10.1001/jama.2020.4940. [Epub ahead of print]).
- [111] Chen L., Xiong J., Bao L. and Shi Y. Convalescent plasma as a potential therapy for COVID-19. *Lancet Infect Dis.* 2020;Apr; 20(4):398–400. Published online 2020 Feb 2027. doi: 10.1016/S1473-3099(2020)30141-30149.
- [112] European Commission (EC). An EU programme of COVID-19 convalescent plasma collection and transfusion; Guidance on collection, testing, processing, storage, distribution and monitored use, Version 1.0 2020 [cited April 4]. Available from: https://ec.europa.eu/health/sites/health/files/blood_tissues_organs/docs/guidance_plasma_covid19_en.pdf.
- [113] Food and Drug Administration (FDA). Recommendations for Investigational COVID-19 Convalescent Plasma. 2020 [cited April 13]. Available from: <https://www.fda.gov/vaccines-blood-biologics/investigational-new-drug-ind-or-device-exemption-ide-process-cber/recommendations-investigational-covid-19-convalescent-plasma>.
- [114] Tanne J. Covid-19: FDA approves use of convalescent plasma to treat critically ill patients *BMJ.* 2020; Mar 26(368:m1256. doi: 10.1136/bmj.m1256.).
- [115] Food and Drug Administration (FDA). FDA Issues Emergency Use Authorization for Convalescent Plasma as Potential Promising COVID-19 Treatment, Another Achievement in Administration's Fight Against Pandemic. 2020 [cited 23/08/2020]. Available from: <https://www.fda.gov/news-events/press-announcements/fda-issues-emergency-use-authorization-convalescent-plasma-potential-promising-covid-19-treatment>.
- [116] National Institute of Health (NIH). COVID-19 Treatment Guidelines. 2020. Available from: <https://covid19treatmentguidelines.nih.gov/introduction/>.
- [117] Li L., Zhang W., Hu Y., Tong X., Zheng S., Yang J., et al. Effect of Convalescent Plasma Therapy on Time to Clinical Improvement in Patients With Severe and Life-threatening COVID-19: A Randomized Clinical Trial. *JAMA.* 2020. DOI: 10.1001/jama.2020.10044.
- [118] Gharbharan A., Jordans C., GeurtsvanKessel C., den Hollander J., Karim F., Mollema F., et al. Convalescent Plasma for COVID-19. A randomized clinical trial. *medRxiv.* 2020:2020.2007.2001.20139857. DOI: 10.1101/2020.07.01.20139857.
- [119] Avendano-Sola C., Ramos-Martinez A., Munez-Rubio E., Ruiz-Antoran B., Malo de Molina R., Torres F., et al. Convalescent Plasma for COVID-19: A multicenter, randomized clinical trial. *medRxiv.* 2020:2020.2008.2026.20182444. DOI: 10.1101/2020.08.26.20182444.

References

- [120] Agarwal A., Mukherjee A., Kumar G., Chatterjee P., Bhatnagar T., Malhotra P., et al. Convalescent plasma in the management of moderate COVID-19 in India: An open-label parallel-arm phase II multicentre randomized controlled trial (PLACID Trial). medRxiv. 2020:2020.2009.2003.20187252. DOI: 10.1101/2020.09.03.20187252.
- [121] Agarwal A., Mukherjee A., Kumar G., Chatterjee P., Bhatnagar T. and Malhotra P. Convalescent plasma in the management of moderate covid-19 in adults in India: open label phase II multicentre randomised controlled trial (PLACID Trial). BMJ. 2020;371:m3939. DOI: 10.1136/bmj.m3939.
- [122] Balcells M., Rojas L., Le Corre N., Martínez-Valdebenito C., Ceballos M., Ferrés M., et al. Early Anti-SARS-CoV-2 Convalescent Plasma in Patients Admitted for COVID-19: A Randomized Phase II Clinical Trial. medRxiv. 2020:2020.2009.2017.20196212. DOI: 10.1101/2020.09.17.20196212.
- [123] Simonovich V., Burgos Pratz L., Scibona P., Beruto M., Vallone M., Vázquez C., et al. A Randomized Trial of Convalescent Plasma in Covid-19 Severe Pneumonia. N Engl J Med. 2020. Epub 2020/11/25. DOI: 10.1056/NEJMoa2031304.
- [124] Marovich M., Mascola J. and Cohen M. Monoclonal Antibodies for Prevention and Treatment of COVID-19. JAMA. 2020. DOI: 10.1001/jama.2020.10245.
- [125] Federal Drug Administration (FDA). Fact Sheet for Health Care Providers: Emergency Use Authorization (EUA) of Casirivimab and Imdevimab. 2020. Available from: <https://www.fda.gov/newsevents/press-announcements/coronavirus-covid-19-update-fda-authorizes-monoclonalantibodies-treatment-covid-19>.
- [126] Chen P., Nirula A., Heller B., Gottlieb R., Boscia J., Morris J., et al. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. New England Journal of Medicine. 2020. DOI: 10.1056/NEJMoa2029849.
- [127] Zheng F., Zhou Y., Zhou Z., Ye F., Huang B., Huang Y., et al. A Novel Protein Drug, Novaferon, as the Potential Antiviral Drug for COVID-19. medRxiv. 2020:2020.2004.2024.20077735. DOI: 10.1101/2020.04.24.20077735.
- [128] European network for health technology assessment (EUnetHTA). Rolling Collaborative Review (RCR13) Authoring Team. Interferon beta-1a (IFN β -1a) and Novaferon (Nova) for the treatment of COVID-19. Diemen (The Netherlands): EUnetHTA. 2020. Available from: <https://www.eunethta.eu>.
- [129] Li C., Luo F., Liu C., Xiong N., Xu Z. and Zhang W. Engineered interferon alpha effectively improves clinical outcomes of COVID-19 patients. Research Square. 2020. DOI: 10.21203/rs.3.rs-65224/v1.
- [130] European network for health technology assessment (EUnetHTA). Rolling Collaborative Review (RCR06) Authoring Team. Solnatide for the treatment of COVID-19. Diemen (The Netherlands). 2020 [cited 15/08/2020]. Available from: https://eunethta.eu/wp-content/uploads/2020/08/EUnetHTA-Covid-19_RCR06_SOLNATIDE_August2020_FINAL.pdf.
- [131] Wang X., Cao R., Zhang H., Liu J., Xu M., Hu H., et al. The anti-influenza virus drug, arbidol is an efficient inhibitor of SARS-CoV-2 in vitro. Cell Discovery. 2020;6(1):28. DOI: 10.1038/s41421-020-0169-8.
- [132] Li Y., Xie Z., Lin W., Cai W., Wen C., Guan Y., et al. Efficacy and safety of lopinavir/ritonavir or arbidol in adult patients with mild/moderate COVID-19: an exploratory randomized controlled trial. Med. 2020. DOI: <https://doi.org/10.1016/j.medj.2020.04.001>.

References

- [133] Nojomi M., Yasin Z., Keyvani H., Makiani M., Roham M., Laali A., et al. Effect of Arbidol on COVID-19: A Randomized Controlled Trial. Research Square. 2020. DOI: 10.21203/rs.3.rs-78316/v1.
- [134] Yethindra V., Tagaev T., Uulu M. and Parihar Y. Efficacy of umifenovir in the treatment of mild and moderate COVID-19 patients. International Journal of Research in Pharmaceutical Sciences. 2020;11(SPL1):506-509. DOI: <https://doi.org/10.26452/ijrps.v11iSPL1.2839>.
- [135] Solinas C., Perra L., Aiello M., Migliori E. and Petrosillo N. A critical evaluation of glucocorticoids in the management of severe COVID-19. Cytokine & growth factor reviews. 2020:S1359-6101(1320)30161-30161. DOI: 10.1016/j.cytogfr.2020.06.012.
- [136] Canadian Agency for Drugs and Technologies in Health (CADTH). Dexamethasone in the Treatment of Hospitalized Patients With COVID-19: A Critical Appraisal of the RECOVERY Trial. 2020. Available from: <https://cadth.ca/sites/default/files/covid-19/ha0005-dexamethasone-fca-of-recovery.pdf>.
- [137] Government UK. World first coronavirus treatment approved for NHS use by government. [cited 16/06/2020]. Available from: <https://www.gov.uk/government/news/world-first-coronavirus-treatment-approved-for-nhs-use-by-government>.
- [138] European Medicines Agency (EMA). Treatments and vaccine for COVID-19. Dexamethasone. 2020 [cited 04/09/2020]. Available from: <https://www.ema.europa.eu/en/human-regulatory/overview/public-health-threats/coronavirus-disease-covid-19/treatments-vaccines-covid-19#dexamethasone--section>.
- [139] European Medicines Agency (EMA). EMA endorses use of dexamethasone in COVID-19 patients on oxygen or mechanical ventilation. EMA/483739/2020. 2020 [cited 18/09/2020]. Available from: <https://www.ema.europa.eu/en/news/ema-endorses-use-dexamethasone-covid-19-patients-oxygen-mechanical-ventilation>.
- [140] Bani-Sadr F., Hentzien M., Pascard M., N'Guyen Y., Servettaz A., Andreoletti L., et al. Corticosteroid therapy for patients with COVID-19 pneumonia: a before-after study. Int J Antimicrob Agents. 2020;56(2):106077. Epub 2020/07/08. DOI: 10.1016/j.ijantimicag.2020.106077.
- [141] Sterne J., Murthy S., Diaz J., Slutsky A., Villar J., Angus D., et al. Association Between Administration of Systemic Corticosteroids and Mortality Among Critically Ill Patients With COVID-19: A Meta-analysis. JAMA. 2020. Epub 2020/09/03. DOI: 10.1001/jama.2020.17023.
- [142] The RECOVERY Collaborative Group. Dexamethasone in Hospitalized Patients with Covid-19 — Preliminary Report. New England Journal of Medicine. 2020. DOI: 10.1056/NEJMoa2021436.
- [143] European network for health technology assessment (EUnetHTA). Rolling Collaborative Review (RCR08) Authoring Team. Dexamethasone for the treatment of COVID-19. Diemen (The Netherlands). 2020 [cited 15/08/2020]. Available from: https://eunetha.eu/wp-content/uploads/2020/08/EUnetHTA-Covid-19_RCR08_DEXAMETHASONE_August2020_FINAL.docx.pdf.
- [144] Tomazini B., Maia I., Cavalcanti A., Berwanger O., Rosa R., Veiga V., et al. Effect of Dexamethasone on Days Alive and Ventilator-Free in Patients With Moderate or Severe Acute Respiratory Distress Syndrome and COVID-19: The CoDEX Randomized Clinical Trial. JAMA. 2020. DOI: 10.1001/jama.2020.17021.

References

- [145] Prescott H. and Rice T. Corticosteroids in COVID-19 ARDS: Evidence and Hope During the Pandemic. *JAMA*. 2020. Epub 2020/09/03. DOI: 10.1001/jama.2020.16747.
- [146] Dequin P., Heming N., Meziani F., Plantefève G., Voiriot G., Badié J., et al. Effect of Hydrocortisone on 21-Day Mortality or Respiratory Support Among Critically Ill Patients With COVID-19: A Randomized Clinical Trial. *JAMA*. 2020. Epub 2020/09/03. DOI: 10.1001/jama.2020.16761.
- [147] Angus D., Derde L., Al-Beidh F., Annane D., Arabi Y., Beane A., et al. Effect of Hydrocortisone on Mortality and Organ Support in Patients With Severe COVID-19: The REMAP-CAP COVID-19 Corticosteroid Domain Randomized Clinical Trial. *JAMA*. 2020. Epub 2020/09/03. DOI: 10.1001/jama.2020.17022.
- [148] Jeronimo C., Farias M., Val F., Sampaio V., Alexandre M., Melo G., et al. Methylprednisolone as Adjunctive Therapy for Patients Hospitalized With COVID-19 (Metcovid): A Randomised, Double-Blind, Phase IIB, Placebo-Controlled Trial. *Clin Infect Dis*. 2020. Epub 2020/08/14. DOI: 10.1093/cid/ciaa1177.
- [149] Edalatifard M., Akhtari M., Salehi M., Naderi Z., Jamshidi A., Mostafaei S., et al. Intravenous methylprednisolone pulse as a treatment for hospitalised severe COVID-19 patients: results from a randomised controlled clinical trial. *Eur Respir J*. 2020. Epub 2020/09/19. DOI: 10.1183/13993003.02808-2020.
- [150] Farahani R., Mosaed R., Nezami-Asl A., Chamanara M., Soleiman-Meigooni S., Kalantar S., et al. Evaluation of the Efficacy of Methylprednisolone Pulse Therapy in Treatment of Covid-19 Adult Patients with Severe Respiratory Failure: Randomized, Clinical Trial. *Research Square*. 2020. DOI: 10.21203/rs.3.rs-66909/v1.
- [151] Borazan N. and Furst D. Nonsteroidal Anti-Inflammatory Drugs, Disease-Modifying Antirheumatic Drugs, Nonopioid Analgesics, & Drugs Used in Gout. In: B. Katzung, S. Masters and A. Trevor, editors. *Basic and Clinical Pharmacology*. New York: McGrawHill; 2012. p. 635-657.
- [152] Deftereos S., Giannopoulos G., Vrachatis D., Siasos G., Giotaki S., Gargalianos P., et al. Effect of Colchicine vs Standard Care on Cardiac and Inflammatory Biomarkers and Clinical Outcomes in Patients Hospitalized With Coronavirus Disease 2019: The GRECCO-19 Randomized Clinical Trial. *JAMA Network Open*. 2020;3(6):e2013136-e2013136. DOI: 10.1001/jamanetworkopen.2020.13136.
- [153] Salehzadeh F., Pourfarzi F. and Ataei S. The Impact of Colchicine on The COVID-19 Patients; A Clinical Trial Study. *BMC infectious diseases*. 2020. DOI: 10.21203/rs.3.rs-69374/v1.
- [154] Lopes M., Bonjorno L., Giannini M., Amaral N., Benatti M., Rezek U., et al. Beneficial effects of colchicine for moderate to severe COVID-19: an interim analysis of a randomized, double-blinded, placebo controlled clinical trial. *medRxiv*. 2020:2020.2008.2006.20169573. DOI: 10.1101/2020.08.06.20169573.
- [155] European network for health technology assessment (EUnetHTA). Rolling Collaborative Review (RCR14) Authoring Team. Gimsilumab for the treatment of COVID-19. Diemen (The Netherlands): EUnetHTA; 2020. Available from: https://eunetha.eu/wp-content/uploads/2020/08/EUnetHTA-Covid-19_RCR15_Canakinumab_August2020_FINAL.pdf.
- [156] A Study to Assess the Efficacy and Safety of Gimsilumab in Subjects With Lung Injury or Acute Respiratory Distress Syndrome Secondary to COVID-19 (BREATHE). 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04351243?term=gimsilumab&draw=2&rank=1>.

References

- [157] European Medicines Agency (EMA). Ilaris. Summary of Product Characteristics. 2020. Available from: https://www.ema.europa.eu/en/documents/product-information/ilaris-epar-product-information_en.pdf.
- [158] European network for health technology assessment (EUnetHTA). Rolling Collaborative Procedure (RCR15). Authoring Team. Canakinumab for the treatment of Covid-19. Diemen (The Netherlands): EUnetHTA. 2020. Available from: https://eunetha.eu/wp-content/uploads/2020/08/EUnetHTA-Covid-19_RCR15_Canakinumab_August2020_FINAL.pdf.
- [159] Study of Efficacy and Safety of Canakinumab Treatment for CRS in Participants With COVID-19-induced Pneumonia (CAN-COVID). 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04362813?term=canakinumab&draw=2&rank=4>.
- [160] Canakinumab in Covid-19 Cardiac Injury (The Three C Study). 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04365153?term=canakinumab&draw=4>.
- [161] Rizk J., Kalantar-Zadeh K., Mehra M., Lavie C., Rizk Y. and Forthal D. Pharmaco-Immunomodulatory Therapy in COVID-19. *Drugs*. 2020;80(13):1267-1292. Epub 2020/07/23. DOI: 10.1007/s40265-020-01367-z.
- [162] Bonaventura A., Vecchié A., Wang T., Lee E., Cremer P., Carey B., et al. Targeting GM-CSF in COVID-19 Pneumonia: Rationale and Strategies. *Front Immunol*. 2020;11:1625. Epub 2020/07/29. DOI: 10.3389/fimmu.2020.01625.
- [163] Ferrari D., Locatelli M., Briguglio M. and Lombardi G. Is there a link between vitamin D status, SARS-CoV-2 infection risk and COVID-19 severity? *Cell Biochemistry and Function*. 2020:1-13. DOI: <https://doi.org/10.1002/cbf.3597>.
- [164] The National Heart Lung and Blood Institute PETAL Clinical Trials Network. Early High-Dose Vitamin D3 for Critically Ill, Vitamin D-Deficient Patients. *New England Journal of Medicine*. 2019;381(26):2529-2540. DOI: 10.1056/NEJMoa1911124.
- [165] Entrenas Castillo M., Entrenas Costa L., Vaquero Barrios J., Alcalá Díaz J., López Miranda J., Bouillon R., et al. Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study. *The Journal of Steroid Biochemistry and Molecular Biology*. 2020;203:105751. DOI: <https://doi.org/10.1016/j.jsbmb.2020.105751>.
- [166] Rastogi A., Bhansali A., Khare N., Suri V., Yaddanapudi N., Sachdeva N., et al. Short term, high-dose vitamin D supplementation for COVID-19 disease: a randomised, placebo-controlled, study (SHADE study). *Postgraduate Medical Journal*. 2020:postgradmedj-2020-139065. DOI: 10.1136/postgradmedj-2020-139065.
- [167] Murai I., Fernandes A., Sales L., Pinto A., Goessler K., Duran C., et al. Effect of Vitamin D₃ Supplementation vs Placebo on Hospital Length of Stay in Patients with Severe COVID-19: A Multicenter, Double-blind, Randomized Controlled Trial. *medRxiv*. 2020:2020.2011.2016.20232397. DOI: 10.1101/2020.11.16.20232397.
- [168] European Medicines Agency (EMA). Olimiant (baricitinib): EPAR – Medicine overview. 2020. Available from: <https://www.ema.europa.eu/en/medicines/human/EPAR/olumiant>.
- [169] European network for health technology assessment (EUnetHTA). Rolling Collaborative Review (RCR18) Authoring Team. Baricitinib for the treatment of COVID-19. Diemen (The Netherlands):

References

2020. Available from: https://eunetha.eu/wp-content/uploads/2020/12/EUnetHTA-Covid-19_RCR18_Baricitinib_v1.0_FINAL.pdf.
- [170] Federal Drug Administration (FDA). Fact sheet for healthcare providers emergency use authorization (EUA) of Baricitinib. 2020. Available from: <https://www.fda.gov/media/143823/download>.
- [171] Kalil A., Patterson T., Mehta A., Tomashek K., Wolfe C., Ghazaryan V., et al. Baricitinib plus Remdesivir for Hospitalized Adults with Covid-19. *New England Journal of Medicine*. 2020. DOI: 10.1056/NEJMoa2031994.
- [172] Cox R., Wolf J. and Plemper R. Therapeutic MK-4482/EIDD-2801 Blocks SARS-CoV-2 Transmission in Ferrets. *Research Square*. 2020:rs.3.rs-89433. DOI: 10.21203/rs.3.rs-89433/v1.
- [173] European network for health technology assessment (EUnetHTA). Rolling Collaborative Review (RCR19) Authoring Team. Molnupiravir for the treatment of COVID-19. Diemen (The Netherlands): 2020. Available from: https://eunetha.eu/wp-content/uploads/2020/12/EUnetHTA-Covid-19_RCR19_Molnupiravir_v1.0_FINAL.pdf.



HTA Austria

Austrian Institute for
Health Technology Assessment
GmbH